STRONG UNIQUE CONTINUATION FOR m -TH POWERS OF A LAPLACIAN OPERATOR WITH SINGULAR COEFFICIENTS

CHING-LUNG LIN

(Communicated by David S. Tartakoff)

ABSTRACT. In this paper we prove strong unique continuation for u satisfying an inequality of the form $|\Delta^m u| \leq f(x, u, Du, \dots, D^k u)$, where k is up to $[3m/2]$. This result gives an improvement of a work by Colombini and Grammatico (1999) in some sense. The proof of the main theorem is based on Carleman estimates with three-parameter weights $|x|^{2\sigma_1}(\log|x|)^{2\sigma_2}\exp(\frac{\beta}{2}(\log|x|)^2)$.

1. INTRODUCTION

Let Ω be a connected open subset of $\mathbb{R}^n$ ($n \geq 2$) containing 0. In [6], Colombini and Grammatico prove some results of strong unique continuation for u satisfying an inequality of the form

$$(1.1) \quad |\Delta^m u| \leq f(x, u, Du, \dots, D^k u) \quad \text{for } k = m.$$

In this paper we are interested in u satisfying the following forms:

$$(1.2) \quad |\Delta^m u| \leq C_0 \sum_{|\alpha| \leq m-1} |x|^{-2m+|\alpha|} |D^\alpha u| + C_0 \sum_{|\alpha|=m}^{[3m/2]} |x|^{-2m+|\alpha|+\epsilon} |D^\alpha u|,$$

where $0 < \epsilon < 1/2$ and the orders of lower order terms are up to $[3m/2]$.

Theorem 1.1. *Let $u \in H_{loc}^{2m}(\Omega)$ be a solution of (1.2), and for all $N > 0$ let*

$$\int_{|x| \leq R} |u|^2 dx = O(R^N), \quad R \rightarrow 0.$$

Then u is identically zero in Ω .

This problem for $m = 1$ has drawn a lot of attention in partial differential equations and mathematical physics. For the development of this problem, the author refers the readers to [15]. In particular, their work was focused on second order equations in which the coefficients of the lower order terms are allowed to be singular (see [2], [3], [11] and [5]). Here we mention two articles which are closely related to our work in this paper.

Hörmander [8] proved that if $u \in H_{loc}^1(\Omega)$ satisfies

$$|\Delta u| \leq C_1 |x|^{-2+\epsilon} |u| + C_2 |x|^{-1+\epsilon} |\nabla u|, \quad \epsilon > 0,$$

and u vanishes of infinite order at 0, then u is identically zero in Ω .

Received by the editors August 23, 2005.

2000 *Mathematics Subject Classification.* Primary 54C40, 14E20; Secondary 46E25, 20C20.

The author was supported in part by the Taiwan National Science Council, NSC 93-2119-M-194-007.

Regbaoui [13] extended Hörmander's result to the sharp case $\epsilon = 0$ with small C_2 . His proof was also based on suitable Carleman estimates, but he worked with the strictly convex weights $\varphi_\beta(x) = \exp(\frac{\beta}{2}(\log|x|)^2)$ rather than the usual polynomial weights. As for negative results, some counterexamples were given when C_2 is not small (such as [1] and [16]).

For the case $m = 2$, Borgne [4] got some results about strong unique continuation when $k = 3$ in (1.1). On the other hand, Watanabe [14] established some uniqueness results for $m = 3$ with $k = 5$. For $m = 4$, Ōkaji [10] solved unique continuation when $k = 7$. For the higher orders, Protter [12] got the unique continuation for inequalities of the form (1.1) in which k is up to $[3m/2]$. According to the counterexamples for $n = 2$ in [7], Colombini and Grammatico [6] proved the sharp results about the coefficients of the m -th lower order terms for strong unique continuation with some cases when $k = m$. In this paper, we consider the cases (1.2).

We will follow Regbaoui's approach in [13] and organize this paper as follows. In Section 2, we shall study the asymptotic behavior of the function u near 0 which guarantees the use of the singular weights $|x|^{2\sigma_1}(\log|x|)^{2\sigma_2}\varphi_\beta^2$. Some of the key Carleman estimates with the weights $|x|^{2\sigma_1}(\log|x|)^{2\sigma_2}\varphi_\beta^2$ will be derived in Section 3. Using these Carleman estimates, we prove Theorem 1.1 in Section 4.

2. CARLEMAN ESTIMATES WITH POLYNOMIAL WEIGHTS

Lemma 2.1. *There exists a positive constant C_m such that for any $u \in C_0^\infty(\mathbb{R}^n \setminus \{0\})$ and for any $\tau \in \{k + 1/2, k \in \mathbb{N}\}$, we have the estimate*

$$(2.1) \quad \sum_{|\alpha| \leq 2m} \int \tau^{2m-2|\alpha|} |x|^{-2\tau+2|\alpha|-n} |D^\alpha u|^2 dx \leq C_m \int |x|^{-2\tau+4m-n} |\Delta^m u|^2 dx.$$

Proof. In [13], Regbaoui proved in Lemma 2.1 for any $u \in C_0^\infty(\mathbb{R}^n \setminus \{0\})$ and for any $\tau \in \{k + 1/2, k \in \mathbb{N}\}$ that

$$(2.2) \quad \sum_{|\alpha| \leq 2} \int \tau^{2-2|\alpha|} |x|^{-2\tau+2|\alpha|-n} |D^\alpha u|^2 dx \leq C \int |x|^{-2\tau+4-n} |\Delta u|^2 dx.$$

By the repeated use of (2.2), we can get (2.1).

Remark 2.2. The estimate (2.1) in Lemma 2.1 remains valid if we assume $u \in H_{loc}^{2m}(\Omega)$ with compact support and satisfying for all $|\alpha| \leq 2m$ and all $N > 0$, $\int_{|x| \leq R} |D^\alpha u|^2 dx = O(R^N)$ as $R \rightarrow 0$. This can be easily obtained by cutting u off for small $|x|$ and regularizing.

Motivated by [8] and [13], we get the following theorem. The proof of this theorem is based on Lemma 2.1. From now on, c stands for a generic constant and its value may vary from line to line.

Theorem 2.3. *Let $u \in H_{loc}^{2m}(\Omega)$ be a solution to (1.2). If $|\alpha| \leq 2m$, then*

$$(2.3) \quad \int_{|x| \leq R} |D^\alpha u|^2 dx = O(e^{-BR^{-\epsilon/m}}), \quad R \rightarrow 0,$$

for some positive constant B .

Proof. Following Hörmander's argument in [9] (Corollary 17.1.4., p. 8), we show that if $u \in H_{loc}^{2m}(\Omega)$ is a solution of (1.2), then for all $|\alpha| \leq 2m$ and for all $N > 0$

$$(2.4) \quad \int_{|x| \leq R} |D^\alpha u|^2 dx = O(R^N), \quad R \rightarrow 0.$$

In view of Remark 2.2, we can apply (2.1) to the function ξu , where $\xi(x) \in C_0^\infty(\mathbb{R}^n)$ such that $\xi(x) = 1$ for $|x| \leq R$ and $\xi(x) = 0$ for $|x| \geq 2R$ ($R > 0$ sufficiently small). Here the number R is not yet fixed and is given by $R = (\gamma\tau)^{-m/\epsilon}$, where $\gamma > 0$ is a large constant which will be chosen later. Using the estimate (2.1) and the equation (1.2), we can derive that

$$\begin{aligned}
 & \sum_{|\alpha| \leq m-1} \tau^{2m-2|\alpha|} \int_{|x| \leq R} |x|^{-2\tau+2|\alpha|-n} |D^\alpha u|^2 dx \\
 & + \sum_{|\alpha|=m}^{2m} \gamma^{2m} \int_{|x| \leq R} |x|^{-2\tau+2|\alpha|-n+2\epsilon} |D^\alpha u|^2 dx \\
 = & \sum_{|\alpha| \leq m-1} \tau^{2m-2|\alpha|} \int_{|x| \leq R} |x|^{-2\tau+2|\alpha|-n} |D^\alpha u|^2 dx \\
 & + \sum_{|\alpha|=m}^{2m} R^{-2\epsilon} \tau^{-2m} \int_{|x| \leq R} |x|^{-2\tau+2|\alpha|-n+2\epsilon} |D^\alpha u|^2 dx \\
 \leq & \sum_{|\alpha| \leq m-1} \tau^{2m-2|\alpha|} \int_{|x| \leq R} |x|^{-2\tau+2|\alpha|-n} |D^\alpha u|^2 dx \\
 & + \sum_{|\alpha|=m}^{2m} \tau^{2m-2|\alpha|} \int_{|x| \leq R} |x|^{-2\tau+2|\alpha|-n} |D^\alpha u|^2 dx \\
 (2.5) \quad & \leq \sum_{|\alpha| \leq 2m} \tau^{2m-2|\alpha|} \int |x|^{-2\tau+2|\alpha|-n} |D^\alpha(\chi u)|^2 dx \\
 & \leq C_m \int |x|^{-2\tau+4m-n} |\Delta^m(\chi u)|^2 dx \\
 & \leq cC_m \int_{|x| \leq R} |x|^{-2\tau+4m-n} |\Delta^m u|^2 dx \\
 & \quad + cC_m \int_{|x| > R} |x|^{-2\tau+4m-n} [\chi, \Delta^m] u|^2 dx \\
 & \leq cC_m C_0 \sum_{|\alpha| \leq m-1} \int_{|x| \leq R} |x|^{-2\tau+2|\alpha|-n} |D^\alpha u|^2 dx \\
 & \quad + cC_m C_0 \sum_{|\alpha|=m}^{[3m/2]} \int_{|x| \leq R} |x|^{-2\tau+2|\alpha|-n+2\epsilon} |D^\alpha u|^2 dx \\
 & \quad + cC_m \int_{|x| > R} |x|^{-2\tau+4m-n} [\chi, \Delta^m] u|^2 dx,
 \end{aligned}$$

where $[\cdot, \cdot]$ denotes the commutator.

Therefore, carefully checking terms on both sides of (2.5), we can choose γ and let τ be large enough such that all terms with $\int_{|x| \leq R} |x|^{-2\tau+2|\alpha|-n} |D^\alpha u|^2 dx$ and

$\int_{|x| \leq R} |x|^{-2\tau+2|\alpha|-n+2\epsilon} |D^\alpha u|^2 dx$ on the right side of (2.5) are absorbed by the left-hand side. We now fix such γ . By construction of χ we have $|D^\alpha \chi| \leq c_1 R^{-|\alpha|}$, where c_1 is a positive constant. Consequently, it follows for $R < 1/2$ and (2.5) that

$$\begin{aligned}
 & \tau^{-2m} (R/2)^{-2\tau+4m-n+2\epsilon} \sum_{|\alpha| \leq 2m} \int_{|x| \leq R/2} |D^\alpha u|^2 dx \\
 & \leq \sum_{|\alpha| \leq 2m} \tau^{2m-2|\alpha|} (R/2)^{-2\tau+2|\alpha|-n+2\epsilon} \int_{|x| \leq R/2} |D^\alpha u|^2 dx \\
 & \leq \sum_{|\alpha| \leq 2m} \tau^{2m-2|\alpha|} \int_{|x| \leq R/2} |x|^{-2\tau+2|\alpha|-n+2\epsilon} |D^\alpha u|^2 dx \\
 (2.6) \quad & \leq \sum_{|\alpha| \leq m-1} \tau^{2m-2|\alpha|} \int_{|x| \leq R} |x|^{-2\tau+2|\alpha|-n} |D^\alpha u|^2 dx \\
 & \quad + \sum_{|\alpha|=m}^{2m} \gamma^{2m} \int_{|x| \leq R} |x|^{-2\tau+2|\alpha|-n+2\epsilon} |D^\alpha u|^2 dx \\
 & \leq c \int_{|x| > R} |x|^{-2\tau+4m-n} |[\chi, \Delta^m] u|^2 dx \\
 & \leq c R^{-4m} R^{-2\tau+4m-n} \|u\|_{H^{2m}}^2 \\
 & = c \|u\|_{H^{2m}}^2 R^{-2\tau-n},
 \end{aligned}$$

where $\|u\|_{H^{2m}}^2$ is the H^{2m} norm of u in the ball $B(0, 2R)$.

Recall that for $R = (\gamma\tau)^{-m/\epsilon}$, we have

$$\begin{aligned}
 (2.7) \quad & \sum_{|\alpha| \leq 2m} \int_{|x| \leq R/2} |D^\alpha u|^2 dx \leq c R^{-4m-4\epsilon} 2^{-2\gamma^{-1} R^{-\epsilon/m}} \|u\|_{H^{2m}}^2 \\
 & \leq c e^{-\tilde{B} R^{-\epsilon/m}}.
 \end{aligned}$$

It should be noted that (2.7) is valid for $\tau \in \mathbb{N} + \frac{1}{2}$ and $R = (\gamma\tau)^{-m/\epsilon}$. Therefore, if we choose $\tau \in \{j + \frac{1}{2} : j \in \mathbb{N}\}$, then (2.7) only holds for $R_j = (\gamma(j + \frac{1}{2}))^{-m/\epsilon}$. Nevertheless, we can see that

$$R_{j+1} < R_j < 2R_{j+1} \quad \text{for } j \text{ large enough} \quad \text{and} \quad R_j \rightarrow 0 \text{ as } j \rightarrow \infty.$$

Thus, we can conclude that

$$\sum_{|\alpha| \leq 2m} \int_{|x| \leq R} |D^\alpha u|^2 dx \leq c e^{-B R^{-\epsilon/m}},$$

for all sufficiently small $R > 0$ with $B = 2^{-2\epsilon/m} \tilde{B}$.

3. CARLEMAN ESTIMATES WITH MORE SINGULAR WEIGHTS

To prove the following Carleman estimates, we introduce polar coordinates in $\mathbb{R}^n \setminus \{0\}$ by setting $x = r\omega$, with $r = |x|$, $\omega = (\omega_1, \dots, \omega_n) \in S^{n-1}$ when $x \neq 0$. Furthermore, setting $t = \log r$, we can substitute a new coordinate t for r such that

$$\frac{\partial}{\partial x_j} = e^{-t} (\omega_j \partial_t + \Omega_j),$$

where Ω_j is a vector field in S^{n-1} . Then the Laplacian becomes

$$(3.1) \quad e^{2t}\Delta = \partial_t^2 + (n-2)\partial_t + \Delta_\omega,$$

where $\Delta_\omega = \sum_{j=1}^n \Omega_j^2$ is the Laplace-Beltrami operator in S^{n-1} . The vector field Ω_j has the properties

$$\sum_{j=1}^n \omega_j \Omega_j = 0, \quad \sum_{j=1}^n \Omega_j \omega_j = n-1.$$

The adjoint of Ω_j as an operator in $L^2(S^2)$ is

$$(3.2) \quad \Omega_j^* = (n-1)\omega_j - \Omega_j$$

and

$$(3.3) \quad \sum_{j=1}^n \Omega_j^* \Omega_j = -\Delta_\omega.$$

Hereafter we shall use the following notation:

$$(3.4) \quad \begin{aligned} D_0 &= (1/i)\partial_t; \\ D_k &= (1/i)\Omega_k, \quad k = 1, \dots, n; \\ D^\alpha &= D_0^{\alpha_1} \dots D_n^{\alpha_n}, \quad \alpha = (\alpha_0, \dots, \alpha_n) \in \mathbb{N}^{n+1}. \end{aligned}$$

When $r \rightarrow 0$, $t \rightarrow -\infty$, we will be interested in values of t in a neighborhood of $-\infty$. Motivated by [13], we derive the following Carleman estimate with weights $\varphi_\beta = \varphi_\beta(x) = \exp(\frac{\beta}{2}(\log|x|)^2)$.

Theorem 3.1. *Given $\sigma_1 \in \mathbb{Z}$ and $\sigma_2 \in \mathbb{Z}$, there exist a sufficiently large number $\beta_0 > 0$ and a sufficiently small number $r_0 > 0$ depending on n , σ_1 and σ_2 such that for all $u \in U_{r_0}$ with $0 < r_0 < e^{-1}$, $\beta \geq \beta_0$, we have that*

$$(3.5) \quad \begin{aligned} & C \sum_{|\alpha| \leq 2} \beta^{3-2|\alpha|} \int \varphi_\beta^2 |x|^{2\sigma_1+2|\alpha|-n} (\log|x|)^{2\sigma_2+2-2|\alpha|} |D^\alpha u|^2 dx \\ & \leq \int \varphi_\beta^2 |x|^{2\sigma_1+4-n} (\log|x|)^{2\sigma_2} |\Delta u|^2 dx, \end{aligned}$$

where $U_{r_0} = \{u \in C_0^\infty(\mathbb{R}^n \setminus \{0\}) : \text{supp}(u) \subset B_{r_0}\}$ and C is a positive constant.

Proof. By the polar coordinate system, we have

$$(3.6) \quad \begin{aligned} & \int \varphi_\beta^2 |x|^{2\sigma_1+4-n} (\log|x|)^{2\sigma_2} |\Delta u|^2 dx \\ & = \int \int e^{\beta t^2} e^{2\sigma_1 t + 4t} t^{2\sigma_2} |\Delta u|^2 dt d\omega \\ & = \int \int |e^{\beta t^2/2} e^{\sigma_1 t} t^{\sigma_2} e^{2t} \Delta u|^2 dt d\omega. \end{aligned}$$

If we set $u = e^{-\beta t^2/2} e^{-\sigma_1 t} t^{-\sigma_2} v$ and use (3.1), then

$$(3.7) \quad e^{\beta t^2/2} e^{\sigma_1 t} t^{\sigma_2} e^{2t} \Delta u = (\partial_t - \beta t)^2 v + (n-2)(\partial_t - \beta t)v + \Delta_\omega v + a\partial_t v + bv,$$

where $b = 2\beta\sigma_1 t + 2\beta\sigma_2 + \sigma_1^2 + 2\sigma_1\sigma_2 t^{-1} - \sigma_2 t^{-2} + \sigma_2^2 t^{-2} - (n-2)(\sigma_1 + \sigma_2 t^{-1})$ and $a = -2\sigma_1 - 2\sigma_2 t^{-1}$.

By (3.6), (3.7) and β large enough, (3.5) holds if we have

$$(3.8) \quad \sum_{|\alpha| \leq 2} \beta^{3-2|\alpha|} \int \int t^{2-2|\alpha|} |D^\alpha v|^2 dt d\omega \leq C \int \int |\Delta_\beta v|^2 dt d\omega,$$

where $\Delta_\beta v = (\partial_t - \beta t)^2 v + (n-2)(\partial_t - \beta t)v + \Delta_\omega v$ and C is a positive constant.

Note that

$$\Delta_\beta v = \partial_t^2 v + (n-2)\partial_t v - 2\beta t \partial_t v - (n-2)\beta t v - \beta v + \beta^2 t^2 v + \Delta_\omega v.$$

Denote

$$\Delta_{\bar{\beta}} v = \partial_t^2 v - (n-2)\partial_t v + 2\beta t \partial_t v - (n-2)\beta t v - \beta v + \beta^2 t^2 v + \Delta_\omega v.$$

Using integration by parts, we get

$$(3.9) \quad \begin{aligned} I &:= \int \int |\Delta_\beta v|^2 dt d\omega - \int \int |\Delta_{\bar{\beta}} v|^2 dt d\omega \\ &= \int \int (12\beta^3 t^2 + \beta^2 O(t)) |v|^2 dt d\omega + 4\beta \int \int |\partial_t v|^2 dt d\omega \\ &\quad - 4\beta \int \int \sum_j |\Omega_j v|^2 dt d\omega. \end{aligned}$$

Similarly, we can get

$$(3.10) \quad \begin{aligned} J &:= \int \int t^{-2} |\Delta_\beta v|^2 dt d\omega + \int \int t^{-2} |\Delta_{\bar{\beta}} v|^2 dt d\omega \\ &= \int \int (2\beta^4 t^2 + \beta^3 O(t)) |v|^2 dt d\omega + \int (8\beta^2 + \beta O(t^{-1})) |\partial_t v|^2 dt d\omega \\ &\quad + 2 \int \int t^{-2} |\partial_t^2 v|^2 dt d\omega - \int \int \sum_j (4\beta^2 + \beta O(t^{-1})) |\Omega_j v|^2 dt d\omega \\ &\quad + 2 \int \int t^{-2} |\Delta_\omega v|^2 dt d\omega + 4 \int \int t^{-2} \sum_j |\partial_t \Omega_j v|^2 dt d\omega. \end{aligned}$$

If $v \in C_0^\infty(B(0, e^{T_0}))$ and $\beta \gg |T_0|$ is large enough, then it follows from (3.9) and (3.10) that

$$(3.11) \quad \begin{aligned} \beta I + J &\geq 13\beta^4 \int \int t^2 |v|^2 dt d\omega + 2 \int \int t^{-2} |\Delta_\omega v|^2 dt d\omega \\ &\quad - 9\beta^2 \int \int \sum_j |\Omega_j v|^2 dt d\omega + 11\beta^2 \int \int |\partial_t v|^2 dt d\omega \\ &\quad + 2 \int \int t^{-2} |\partial_t^2 v|^2 dt d\omega + 4 \int \int t^{-2} \sum_j |\partial_t \Omega_j v|^2 dt d\omega \\ &\geq K + \frac{13}{39}\beta^4 \int \int t^2 |v|^2 dt d\omega + \frac{1}{20} \int \int t^{-2} |\Delta_\omega v|^2 dt d\omega \\ &\quad + \beta^2 \int \int \sum_j |\Omega_j v|^2 dt d\omega + 11\beta^2 \int \int |\partial_t v|^2 dt d\omega \\ &\quad + 2 \int \int t^{-2} |\partial_t^2 v|^2 dt d\omega + 4 \int \int t^{-2} \sum_j |\partial_t \Omega_j v|^2 dt d\omega, \end{aligned}$$

where

$$K = \frac{500}{39}\beta^4 \int \int t^2 |v|^2 dt d\omega - 10\beta^2 \int \int \sum_j |\Omega_j v|^2 dt d\omega + \frac{39}{20} \int \int t^{-2} |\Delta_\omega v|^2 dt d\omega.$$

Now we shall get a lower bound of K . Then, by (3.3), we have

$$(3.12) \quad -\beta^2 \int \int \sum_j |\Omega_j v|^2 dt d\omega = \beta^2 \int \int \bar{v} \Delta_\omega v dt d\omega.$$

Since $10|ab| \leq 500a^2/39 + 39b^2/20$, it follows that

$$(3.13) \quad 10|\beta^2 v \Delta_\omega v| \leq \frac{500}{39}\beta^4 t^2 |v|^2 + \frac{39}{20}t^{-2} |\Delta_\omega v|^2.$$

Combining (3.12) and (3.13), we get a lower bound of K :

$$(3.14) \quad K = \frac{500}{39}\beta^4 \int \int t^2 |v|^2 dt d\omega - 10\beta^2 \int \int \sum_j |\Omega_j v|^2 dt d\omega + \frac{39}{20} \int \int t^{-2} |\Delta_\omega v|^2 dt d\omega \geq 0.$$

In addition, the ellipticity of Δ_ω implies that there exists a new positive constant C such that

$$(3.15) \quad \sum_{|\alpha|=2} \int \int t^{-2} |\Omega^\alpha v|^2 dt d\omega \leq C \int \int t^{-2} |\Delta_\omega v|^2 dt d\omega.$$

Thus, it follows from (3.11), (3.14) and (3.15) that

$$\begin{aligned} C(\beta I + J) &\geq \beta^4 \int \int t^2 |v|^2 dt d\omega + \int \int t^{-2} |\Delta_\omega v|^2 dt d\omega \\ &\quad + \beta^2 \int \int \sum_j |\Omega_j v|^2 dt d\omega + \beta^2 \int \int |\partial_t v|^2 dt d\omega \\ &\quad + \int \int t^{-2} |\partial_t^2 v|^2 dt d\omega + \int \int t^{-2} \sum_j |\partial_t \Omega_j v|^2 dt d\omega. \end{aligned}$$

Together with

$$\beta I + J \leq (\beta + 1) \int \int |\Delta_\beta v|^2 dt d\omega,$$

(3.8) holds. So we have the result.

Corollary 3.1. *Given $\sigma_1 \in \mathbb{Z}$ and $\sigma_2 \in \mathbb{Z}$, there exist a sufficiently large number $\beta_0 > 0$ and a sufficiently small number $r_0 > 0$ depending on n, m, σ_1 and σ_2 such that for all $u \in U_{r_0}$ with $0 < r_0 < e^{-1}$, $\beta \geq \beta_0$, we have that*

$$\begin{aligned} C \sum_{|\alpha| \leq 2m} \beta^{3m-2|\alpha|} \int \varphi_\beta^2 |x|^{2\sigma_1+2|\alpha|-n} (\log |x|)^{2\sigma_2+2m-2|\alpha|} |D^\alpha u|^2 dx \\ \leq \int \varphi_\beta^2 |x|^{2\sigma_1+4m-n} (\log |x|)^{2\sigma_2} |\Delta^m u|^2 dx, \end{aligned}$$

where C is a positive constant.

Corollary 3.2. *There exist a sufficiently large number $\beta_0 > 0$ and a sufficiently small number $r_0 > 0$ depending on n and m such that for all $u \in U_{r_0}$ with $0 < r_0 < e^{-1}$, $\beta \geq \beta_0$, we have that*

$$(3.16) \quad \begin{aligned} & C \sum_{|\alpha| \leq 2m} \beta^{3m-2|\alpha|} \int \varphi_\beta^2 |x|^{2|\alpha|-n} (\log |x|)^{2m-2|\alpha|} |D^\alpha u|^2 dx \\ & \leq \int \varphi_\beta^2 |x|^{4m-n} |\Delta^m u|^2 dx, \end{aligned}$$

where C is a positive constant.

Remark 3.2. The estimate (3.16) in Corollary 3.2 remains valid if we assume $u \in H_{loc}^{2m}(\Omega)$ with compact support and satisfies for all $|\alpha| \leq 2m$, $\int_{|x| \leq R} |D^\alpha u|^2 dx = O(e^{-BR^{-\epsilon/m}})$ as $R \rightarrow 0$, $B > 0$.

4. PROOF OF THEOREM 1.1

Let $u \in H_{loc}^{2m}(\Omega)$ be a solution of (1.2). By Lemma 2.1 and Theorem 2.3, u is in $H_{loc}^{2m}(\Omega)$ and satisfies (2.3). Thus by Remark 3.2 we can apply (3.16) with the function ξu , where $\xi(x) \in C_0^\infty(\mathbb{R}^n)$ such that $\xi(x) = 1$ for $|x| \leq R$ and $\xi(x) = 0$ for $|x| \geq 2R$ ($R > 0$ small enough). Then

$$(4.1) \quad \begin{aligned} & \sum_{|\alpha| \leq [\frac{3m}{2}]} \beta^{3m-2|\alpha|} \int_{|x| < R} \varphi_\beta^2 |x|^{2|\alpha|-n} (\log |x|)^{2m-2|\alpha|} |D^\alpha u|^2 dx \\ & \leq \sum_{|\alpha| \leq 2m} \beta^{3m-2|\alpha|} \int_{|x| < R} \varphi_\beta^2 |x|^{2|\alpha|-n} (\log |x|)^{2m-2|\alpha|} |D^\alpha u|^2 dx \\ & \leq \sum_{|\alpha| \leq 2m} \beta^{3m-2|\alpha|} \int \varphi_\beta^2 |x|^{2|\alpha|-n} (\log |x|)^{2m-2|\alpha|} |D^\alpha(\xi u)|^2 dx \\ & \leq c \int \varphi_\beta^2 |x|^{4m-n} |\Delta^m(\xi u)|^2 dx \\ & \leq c \int_{|x| \leq R} \varphi_\beta^2 |x|^{4m-n} |\Delta^m u|^2 dx + c \int_{|x| > R} \varphi_\beta^2 |x|^{4m-n} |[\xi, \Delta^m] u|^2 dx \\ & \leq c \sum_{|\alpha| \leq m-1} \int_{|x| \leq R} \varphi_\beta^2 |x|^{2|\alpha|-n} |D^\alpha u|^2 dx \\ & \quad + c \sum_{|\alpha|=m}^{[3m/2]} \int_{|x| \leq R} \varphi_\beta^2 |x|^{2|\alpha|-n+2\epsilon} |D^\alpha u|^2 dx \\ & \quad + c \int_{|x| > R} \varphi_\beta^2 |x|^{4m-n} |[\xi, \Delta^m] u|^2 dx, \end{aligned}$$

because $\lim_{|x|=r \rightarrow 0} r^{2\epsilon} (\log r)^{-2m} = 0$ which implies $c|x|^{2\epsilon} \leq (\log |x|)^{2m}$ as $R \leq R_0$ for some small $R_0 > 0$. Let $\beta > \beta_0$ (β_0 large enough); the first two terms on the right-hand side will be swallowed by the left-hand side. Recall that $\varphi_\beta = \varphi_\beta(x) =$

$\exp(\frac{\beta}{2}(\log |x|)^2)$. Therefore, we obtain that

$$\begin{aligned}
 & \beta^{3m} \exp(\beta(\log R_0)^2) \int_{|x| < R_0} |x|^{-n} (\log |x|)^{2m} |u|^2 dx \\
 & \leq \sum_{|\alpha| \leq [\frac{3m}{2}]} \beta^{3m-2|\alpha|} \int_{|x| < R_0} \varphi_\beta^2 |x|^{2|\alpha|-n} (\log |x|)^{2m-2|\alpha|} |D^\alpha u|^2 dx \\
 (4.2) \quad & \leq c \int_{|x| > R_0} \varphi_\beta^2 |x|^{4m-n} |[\xi, \Delta^m] u|^2 dx \\
 & \leq c \exp(\beta(\log R_0)^2) \int_{|x| > R_0} |x|^{4m-n} |[\xi, \Delta^m] u|^2 dx.
 \end{aligned}$$

Divide $\exp(\beta(\log R_0)^2)$ into both sides of (4.2) and then let $\beta \rightarrow \infty$; this implies that $u = 0$ in $B(0, R_0)$. By standard arguments, we can get $u = 0$ in Ω . This completes the proof.

ACKNOWLEDGEMENT

I would like to thank Professor Jenn-Nan Wang for his many constructive suggestions and comments.

REFERENCES

1. S. Alinhac and M.S. Baouendi, *A counterexample to strong uniqueness for the partial differential equations of Schrödinger's type*, Comm. in P.D.E. **19** (1994), 1723-1733. MR1294476 (95g:35043)
2. N. Aronszajn, A. Krzywicki and J. Szarski, *A unique continuation theorem for exterior differential forms on Riemannian manifolds*, Ark. Mat. **4** (1962), 417-453. MR0140031 (25:3455)
3. B. Barcelo, C.E. Kenig, A. Ruiz and C.D. Sogge, *Weighted Sobolev inequalities for the Laplacian plus lower order terms*, Illinois J. Math. **32** (1988), 230-245. MR0945861 (89h:35048)
4. P. Le Borgne, *Strong uniqueness for fourth order elliptic differential operators*, Indiana Univ. Math. J. **50**, No. 1, (2001), 353-381. MR1857040 (2003i:35064)
5. C. Grammatico, *A result on strong unique continuation for the Laplace operator*, Comm. in P.D.E. **22** (1997), 1475-1491. MR1469579 (98j:35034)
6. F. Colombini and C. Grammatico, *Some remarks on strong unique continuation for the Laplace operator and its power*, Comm. in P.D.E. **24** (1999), 1079-1094. MR1680873 (2000c:35263)
7. F. Colombini and C. Grammatico, *A counterexample to strong unique continuation for all powers of the Laplace operator*, Comm. in P.D.E. **25** (2000), 586-600. MR1748356 (2001c:35067)
8. L. Hörmander, *Uniqueness theorems for second order elliptic differential equations*, Comm. in P.D.E. **8**, No. 1, (1983), 21-64. MR0686819 (85c:35018)
9. L. Hörmander, "The analysis of linear partial differential operators", Vol. 3, Springer-Verlag, Berlin, New York, 1985. MR0781536 (87d:35002a)
10. T. Ōkaji, *Uniqueness of the Cauchy problem for elliptic operators with fourfold characteristics of constant multiplicity*, Comm. in P.D.E. **22** (1997), 269-290. MR1434146 (97k:35004)
11. Y. Pan, *Unique continuation for Schrödinger operators with singular potentials*, Comm. in P.D.E. **17** (1992), 953-965. MR1177300 (93j:35064)
12. M.H. Protter, *Unique continuation for elliptic equations*, Trans. Amer. Math. Soc. **95** (1960), 81-91. MR0113030 (22:3871)
13. R. Regbaoui, *Strong uniqueness for second order differential operators*, J. Diff. Eq. **141** (1997), 201-217. MR1488350 (98i:35002)
14. K. Watanabe, *On the uniqueness of the Cauchy problem for certain elliptic equations with triple characteristics*, Tohoku Math. J. **23** (1971), 473-490. MR0308584 (46:7698)

15. T.H. Wolff, "Fourier analysis and partial differential equations", Miraflores de la Sierra (1992), 99-128. MR1330234 (96c:35068)
16. T.H. Wolff, *A counterexample in a unique continuation problem*, Comm. Anal. Geom. **2** (1994), 79-102. MR1312679 (96a:35018)

DEPARTMENT OF MATHEMATICS, NATIONAL CHUNG-CHENG UNIVERSITY, CHIA-YI 62117,
TAIWAN

E-mail address: cllin@math.ccu.edu.tw