HARMONIC HOMEOMORPHISMS OF THE CLOSED DISC TO ITSELF NEED BE IN $W^{1,p}$, $p < 2$, BUT NOT $W^{1,2}$

GREGORY C. VERCHOTA

(Communicated by Juha M. Heinonen)

ABSTRACT. Harmonic maps u from the closed disc onto bounded convex sets of the plane obey $u \in W^{1,p}$, $p < 2$.

The (complex-valued) Poisson extension $\mathcal{P}F$ of a homeomorphism F of the circle $\mathbb{T}$ onto the boundary of a bounded convex domain $\Omega \subset \mathbb{C}$ extends F to a homeomorphism of the closed disc $\mathbb{D}$ onto $\overline{\Omega}$. This theorem was first proved by Hellmuth Kneser [Kne26]. See Chapter 3 of [Dur04].

Theorem 0.1 (H. Kneser). *Let $\Omega \subset \mathbb{C}$ be a bounded convex domain. Let $F : \mathbb{T} \rightarrow \partial\Omega$ be a homeomorphism. Then*

$$u(z) = \mathcal{P}F(z) = \frac{1}{2\pi} \int_0^{2\pi} \frac{1 - |z|^2}{|e^{it} - z|^2} F(e^{it}) dt, \quad z \in \mathbb{D},$$

is a homeomorphism of $\mathbb{D}$ onto Ω .

The question of Sobolev regularity in $\mathbb{D}$ for such harmonic homeomorphisms has recently arisen [AIMO]. There the authors study homeomorphisms of Ω onto $\mathbb{D}$ with integrable distortion function. The problem of determining when a given homeomorphism between boundaries has an extension to a homeomorphism with integrable distortion is shown to be equivalent to asking a question about the inverse mapping: when does a homeomorphism of the boundaries have an extension with finite Dirichlet integral? Here we prove

Theorem 0.2. *With Ω, F and u as in Kneser's Theorem*

(i) *$u \in W^{1,p}(\mathbb{D})$ when $p < 2$*

and

(ii) *there exist $F : \mathbb{T} \rightarrow \mathbb{T}$ so that $u \notin W^{1,2}(\mathbb{D})$.*

Here $W^{1,p}(\mathbb{D})$ is the Sobolev space of $L^p(\mathbb{D})$ functions u that have distributional derivatives $\nabla u = (\frac{\partial u}{\partial x}, \frac{\partial u}{\partial y}) \in L^p(\mathbb{D})$ and norm $\|u\|_{1,p} = (\int_{\mathbb{D}} |u|^p dx dy)^{\frac{1}{p}} + (\int_{\mathbb{D}} |\nabla u|^p dx dy)^{\frac{1}{p}}$, $1 \leq p < \infty$. So (ii) implies that there exists a homeomorphism of $\partial\Omega$ onto $\mathbb{T}$ which does not admit an extension with integrable distortion [AIMO].

For $p = 2$ Friedrich Prym [Pry71] showed that the Poisson extension of a continuous function on $\mathbb{T}$ could fail to have finite Dirichlet integral in $\mathbb{D}$, i.e. fail to be

---

Received by the editors June 6, 2005 and, in revised form, September 26, 2005.

2000 *Mathematics Subject Classification.* Primary 58E20, 46E35.

The author thanks Tadeusz Iwaniec for this question.

in $W^{1,2}(\mathbb{D})$. By a recent account [MP97], p. 259 (see also [Die81], p. 38), Prym's result was ignored. Hadamard [Had06] later exhibited a lacunary series example which can be found in [Cou50], pp. 9-10 (see also [Joh82], p. 125). These examples, together with the example below, cannot arise by conformal mapping to a bounded domain since then the Jacobian would equal $\frac{1}{2}|\nabla u|^2$.

That the Dirichlet principle in its 19th century formulation (see [BB98], pp. 82-83, 85) fails for a homeomorphism of the boundary, follows here by using later technology. By Gagliardo's Theorem for $1 < p < \infty$ (see [MP97], pp. 207-210 where $\mathbb{T}$ is replaced by a Lipschitz boundary; [Ada75], pp. 204-216 for a proof), the trace $F = Tu$ on the boundary of a function $u \in W^{1,p}(\mathbb{D})$ necessarily satisfies

$$(0.1) \quad \int_0^{2\pi} \int_0^{2\pi} \left| \frac{F(e^{is}) - F(e^{it})}{s - t} \right|^p ds dt < \infty.$$

Here the trace operator is the bounded linear operator

$$T : W^{1,p}(\mathbb{D}) \rightarrow L^p(\mathbb{T}), \quad 1 \leq p < \infty,$$

such that $Tu = u$ on $\mathbb{T}$ whenever $u \in W^{1,p}(\mathbb{D}) \cap C(\overline{\mathbb{D}})$. See p. 133 of [EG92] for a proof of its existence. Consequently in order to prove part (ii) of Theorem 0.2 it suffices to exhibit a homeomorphism F for which (0.1) is infinite when $p = 2$.

In what follows, any homeomorphism F from Kneser's Theorem is considered to be of the form $F(e^{it}) = r(t)e^{if(t)}$, $-\infty < t < \infty$, where $r(t)$ is a positive 2π -periodic continuous function and $f(t)$ is a strictly increasing continuous function satisfying $f(t + 2n\pi) = f(t) + 2n\pi$. Thus the convex domain is situated (without loss of generality) to be starlike with respect to the origin.

Proof. (Part (ii) of Theorem 0.2) Consider the strictly increasing functions $f(t) = f_q(t) = \operatorname{sgn}(t)|\log|t||^{1-q}$ on the interval $-\frac{1}{2} \leq t \leq \frac{1}{2}$ when $1 < q$. Then

$$\int_{-\frac{1}{2}}^0 \int_0^{\frac{1}{2}} \left(\frac{f(s) - f(t)}{s - t} \right)^2 ds dt > \int_{-\frac{1}{2}}^0 \int_0^{\frac{1}{2}} \left(\frac{f(s)}{s - t} \right)^2 ds dt = \int_0^{\frac{1}{2}} \frac{(f(s))^2}{s(2s+1)} ds.$$

By $2s + 1 \leq 2$ and substituting f_q , the last integral is greater than

$$\frac{1}{2} \int_0^{\frac{1}{2}} \frac{|\log|s||^{2-2q}}{s} ds,$$

which diverges for $q \leq \frac{3}{2}$. For this restriction of q ,

$$\left| \frac{e^{if(s)} - e^{if(t)}}{f(s) - f(t)} \right| \geq \frac{1}{\pi} \quad \text{for all } s, t \in \left[-\frac{1}{2}, \frac{1}{2}\right].$$

Each $f(t)$ extends to a homeomorphism $[-\pi, \pi] \rightarrow [-\pi, \pi]$ and the above inequalities then show that (0.1) diverges for $p = 2$ and $F(e^{it}) = e^{if(t)}$. $\square$

Remark 0.3. The traditional Dirichlet principle has been shown to fail for an absolutely continuous homeomorphism of the boundary.

It is also a classical result (see Proposition 7' and Lemma 4' [Ste70], pp. 151-152, but change the sign on $\alpha - 1$; see [Pee76], pp. 8-10, items a. and f.) that harmonic functions in $\mathbb{D}$ will have the p -norm of their gradients bounded by (0.1). It remains to show that every homeomorphism $F = re^{if}$ satisfies the inequality (0.1) when $p < 2$.

Proof. (Part (i) of Theorem 0.2) Every bounded convex domain is also a Lipschitz domain. Because Ω is also starlike with respect to the origin it follows for any two points on the boundary that the ratio formed by a change in the distance to the origin and a change in the argument is uniformly bounded. I.e. by the setup and definition of F there is an $M < \infty$ so that $r \circ f^{-1}$ satisfies a Lipschitz condition $\left| \frac{r(s) - r(t)}{f(s) - f(t)} \right| \leq M$ for all $s \neq t$. Consequently

$$\left| \frac{r(s)e^{if(s)} - r(t)e^{if(t)}}{s - t} \right| \leq (M + \|r\|_\infty) \left(\frac{f(s) - f(t)}{s - t} \right).$$

Take $p > 1$. Assume for the moment that f is absolutely continuous and write $g(s, t) = \left(\frac{f(s) - f(t)}{s - t} \right)^p$. Integrating by parts as

$$\int g(s, t) ds = (s - t)g(s, t) - \int (s - t) \frac{\partial}{\partial s} g(s, t) ds$$

and evaluating, using the monotonicity of f and $p > 1$, yields an intermediate step in Hardy's inequality (see p. 242 of [HLP52], for example)

$$\int_0^{2\pi} \left(\frac{f(s) - f(t)}{s - t} \right)^p ds \leq \frac{p}{p-1} \int_0^{2\pi} \left(\frac{f(s) - f(t)}{s - t} \right)^{p-1} f'(s) ds.$$

The last integrand is bounded by $(2\pi)^{p-1} |t - s|^{1-p} f'(s)$. Integrating the left and right-hand sides in t from 0 to 2π , applying Fubini with $p < 2$ and then the fundamental theorem of calculus in s produces the bound

$$\int \int \left(\frac{f(s) - f(t)}{s - t} \right)^p ds dt \leq \frac{p 2^{p+1} \pi^2}{(p-1)(2-p)}$$

independent of f . When f is not absolutely continuous it can be mollified by using a smooth and symmetric mollifier so that monotonicity and periodicity are retained. By the uniformity of the a priori bounds, positivity of the difference quotients and pointwise convergence of the mollifications of f to f , Fatou's lemma yields the same bounds on the double integral for any such increasing continuous f . In general

$$\int_0^{2\pi} \int_0^{2\pi} \left| \frac{F(e^{is}) - F(e^{it})}{s - t} \right|^p ds dt \leq (M + \|r\|_\infty)^p \frac{p 2^{p+1} \pi^2}{(p-1)(2-p)}, \quad 1 < p < 2,$$

which proves (i) for $1 \leq p < 2$. $\square$

Remark 0.4. The above proof required F to be a homeomorphism and $r \circ f^{-1}$ to satisfy a Lipschitz condition. Thus the conclusion remains true if Ω is replaced by a starlike Lipschitz domain. But then u would not be a homeomorphism in general.

REFERENCES

- [Ada75] Robert A. Adams, *Sobolev spaces*, Academic Press [A subsidiary of Harcourt Brace Jovanovich, Publishers], New York-London, 1975. MR0450957 (56:9247)
- [AIMO] K. Astala, T. Iwaniec, G. J. Martin, and J. Onninen, *Extremal mappings of finite distortion*, Proc. London Math. Soc. **91** (2005), no. 3, 655–702. MR2180459
- [BB98] Haïm Brezis and Felix Browder, *Partial differential equations in the 20th century*, Adv. Math. **135** (1998), no. 1, 76–144. MR1617413 (99i:35001)
- [Cou50] R. Courant, *Dirichlet's Principle, Conformal Mapping, and Minimal Surfaces*, Interscience Publishers, Inc., New York, N.Y., 1950. MR0036317 (12:90a)
- [Die81] Jean Dieudonné, *History of functional analysis*, North-Holland Mathematics Studies, vol. 49, North-Holland Publishing Co., Amsterdam, 1981. MR0605488 (83d:46001)

- [Dur04] Peter Duren, *Harmonic mappings in the plane*, Cambridge University Press, Cambridge, 2004. MR2048384 (2005d:31001)
- [EG92] Lawrence C. Evans and Ronald F. Gariepy, *Measure theory and fine properties of functions*, Studies in Advanced Mathematics, CRC Press, Boca Raton, FL, 1992. MR1158660 (93f:28001)
- [Had06] J. Hadamard, *Sur le principe de Dirichlet*, Bull. Soc. Math. France **34** (1906), 135–138. MR1504545
- [HLP52] G. H. Hardy, J. E. Littlewood, and G. Pólya, *Inequalities*, Cambridge, at the University Press, 1952. MR0046395 (13:727e)
- [Joh82] Fritz John, *Partial differential equations*, Applied Mathematical Sciences, vol. 1, Springer-Verlag, New York, 1982. MRMR831655 (87g:35002)
- [Kne26] Hellmuth Kneser, *Lösung der Aufgabe 41*, Jahresber. Deutsch. Math.-Verein. **35** (1926), 123–124.
- [MP97] Vladimir G. Maz'ya and Sergei V. Poborchi, *Differentiable functions on bad domains*, World Scientific Publishing Co. Inc., River Edge, NJ, 1997. MR1643072 (99k:46057)
- [Pee76] Jaak Peetre, *New thoughts on Besov spaces*, Mathematics Department, Duke University, Durham, N.C., 1976. MR0461123 (57:1108)
- [Pry71] Friedrich Prym, *Zur Integration der Differentialgleichung $\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0$* , J. Reine Angew. Math. **73** (1871), 340–364.
- [Ste70] Elias M. Stein, *Singular integrals and differentiability properties of functions*, Princeton Mathematical Series, No. 30, Princeton University Press, Princeton, N.J., 1970. MR0290095 (44:7280)

DEPARTMENT OF MATHEMATICS, 215 CARNEGIE, SYRACUSE UNIVERSITY, SYRACUSE, NEW YORK 13244

E-mail address: gverchot@syr.edu