A NEW PROOF OF PROXIMALITY FOR M -IDEALS

V. INDUMATHI AND S. LALITHAMBIGAI

(Communicated by Jonathan M. Borwein)

ABSTRACT. We give a new and a simple proof of proximality for M -ideals. Unlike the known proofs, our proof derives proximality of M -ideals directly from the definition of an M -ideal, using the Bishop-Phelps theorem.

More than one proof is available for proximality of M -ideals. We refer the reader to the Notes and Remarks following Chapter II in [5], for details. The earlier proofs of Alfsen and Effros [1] (using “dominated extensions”) and that of Ando [2] (using the notion of “splittable convex sets”), are elaborate and technically involved. A simpler proof was given in Proposition 6.5 of [3] and Lemma 1.1 of [8] (see also Proposition 1.1 of [5]), using the fact that M -ideals have the 3-ball and hence the weaker $1\frac{1}{2}$ -ball intersection property. This property was used to construct a Cauchy minimizing sequence for each element to be approximated. Later, in [6], a geometric property of M -ideals was used to do the same.

In this paper we present an easy, short proof of proximality of M -ideals using just the definition of M -ideals and the Bishop-Phelps theorem.

We begin with some notation and definitions that are needed in the sequel. We assume that all subspaces are closed. Throughout X denotes a real Banach space and X^* denotes the dual of X . For any x in X and a subspace Y of X , we denote by $d(x, Y)$ the distance of x from Y . Let

$$P_Y(x) = \{y \in Y : \|x - y\| = d(x, Y)\}.$$

If $P_Y(x)$ is a non-empty set for each x in X , then the subspace Y is said to be proximal in X . For a subspace Y of X , let $Y^\perp$ denote the annihilator of Y in X , that is,

$$Y^\perp = \{f \in X^* : f \equiv 0 \text{ on } Y\}.$$

For f in X^* , the restriction of f to Y is denoted by $f|_Y$. The following definitions can be found in [5].

Definition 1 ([5]). *A subspace Y of a Banach space X is called an L -summand of X if there exists a subspace Z of X such that $X = Y \oplus_1 Z$, the l_1 -direct sum*

Received by the editors November 14, 2005.

2000 *Mathematics Subject Classification.* Primary 46B20, 41A50, 41A65.

Key words and phrases. Proximal subspace, M -ideal.

The first author was partially supported by DST-NSF project No. DST/INT/US(NSF-RPO-0141)/2003.

of Y and Z . This implies that if $x = y + z$ for x in X , y in Y and z in Z , then $\|x\| = \|y\| + \|z\|$.

Definition 2 ([5]). A subspace Y of a Banach space X is called an M -ideal in X if $Y^\perp$ is an L -summand in X^* .

If Y is an M -ideal in X , then there exists a subspace Z of X^* such that

$$(1) \quad X^* = Y^\perp \oplus_1 Z.$$

Using the canonical isometric isomorphism $Y^* \simeq X^*/Y^\perp \simeq Z$, one can conclude (see Proposition 1.12 of [5]) that

$$(2) \quad Z = \{f \in X^* : \|f\| = \|f|_Y\|\}.$$

Remark 1. If Z is a complex Banach space, Z can be considered as a real Banach space and $f \rightarrow \operatorname{Re} f$ is an isometry from $Z_{\mathbb{C}}^*$ onto $Z_{\mathbb{R}}^*$. Thus, if Y is a (complex) subspace and an M -ideal in Z , then $Y_{\mathbb{R}}$ is an M -ideal in $Z_{\mathbb{R}}$. Hence, proving proximality of M -ideals in the case of real Banach spaces would also imply the result for the complex case (see Proposition 1.23 of [5]).

We start our proof by quoting a corollary of the Hahn-Banach theorem that is needed in the sequel. This corollary characterizes the set of nearest points from a subspace.

Lemma 1 ([7]). Let X be a Banach space, Y a subspace of X and x an element of X . Then y_0 is in $P_Y(x)$ if and only if there exists an f in $Y^\perp$ such that $\|f\| = 1$ and $f(x - y_0) = \|x - y_0\|$.

We also need the following simple fact about M -ideals.

Remark 2. Let Y and W be subspaces of a Banach space X , with Y contained in W . If Y is an M -ideal in X , then Y is an M -ideal in W .

To see this, let Y be an M -ideal in X . If f_1 is in W^* , let f be a Hahn-Banach extension of f_1 to X . Using (1), get g in $Y^\perp$ and h in Z such that $f = g + h$ and $\|f\| = \|g\| + \|h\|$. Clearly, $f_1 = f|_W = g|_W + h|_W$. We have $\|f_1\| = \|f|_W\| = \|f\|$ and by (2), $\|h\| = \|h|_Y\| = \|h|_W\|$. Now

$$\|g\| = \|f|_W\| - \|h|_W\| \leq \|f|_W - h|_W\| = \|g|_W\| \leq \|g\|.$$

Hence $\|g\| = \|g|_W\|$ and $\|f|_W\| = \|g|_W\| + \|h|_W\|$. It is now clear that Y is an M -ideal in W since the annihilator of Y , in W^* , is an L -summand of W^* .

Let Y be an M -ideal in a Banach space X . Clearly Y is proximal in X if and only if it is proximal in every subspace W of X , containing Y as a hyperplane. By the above remark, Y is an M -ideal in W . Thus, it suffices to prove proximality of Y in X , when Y is a hyperplane in X .

Proposition 1. Let X be a Banach space and let Y be an M -ideal and a hyperplane in X . If there is a norm attaining functional f in X^* such that $\|f|_Y\| < \|f\|$, then Y is proximal in X .

Proof. We assume the norm of f is one and write $f = h + g$, with h in $Y^\perp$ and g in Z , where Z is given by (2). Since

$$\|g\| = \|f|_Y\| < \|f\| = 1 = \|h\| + \|g\|,$$

h is a non-zero element of $Y^\perp$.

Select any x in $X \setminus Y$. Then there is a non-zero scalar α such that f attains its norm at $\alpha x + y$, for some y in Y . We have

$$\begin{aligned} \|\alpha x + y\| &= f(\alpha x + y) = h(\alpha x + y) + g(\alpha x + y) \\ &\leq \|h\| \|\alpha x + y\| + \|g\| \|\alpha x + y\| \\ &= \|\alpha x + y\|. \end{aligned}$$

It is now clear that $\frac{h}{\|h\|}(\alpha x + y) = \|\alpha x + y\|$ and consequently $-\frac{y}{\alpha}$ is in $P_Y(x)$ by Lemma 1. For a scalar a , and y in Y , $P_Y(ax + y) = aP_Y(x) + y$, and so Y is proximal in X . $\square$

We now proceed to show that f as in Proposition 1 can be constructed for any hyperplane of a Banach space.

Theorem 1. *Let X be Banach space and let Y be a hyperplane in X . Then there is a norm attaining functional f in X^* such that $\|f|_Y\| < \|f\|$.*

Proof. Choose a norm one element x in $X \setminus Y$ and define g in X^* by $g(y + \alpha x) = \alpha$ for any scalar α and y in Y . Then $\|g\| \geq 1$ and $g|_Y = 0$. The Bishop-Phelps Theorem guarantees that there is a norm attaining functional f in X^* with $\|f - g\| < \frac{1}{3}$. Clearly $\|f|_Y\| < \frac{1}{3}$, while $\|f\| > \frac{2}{3}$. $\square$

The theorem below follows immediately from Proposition 1 and Theorem 1.

Theorem 2 (See Proposition 1.1 of [5]). *Let X be a Banach space and let Y be an M -ideal in X . Then Y is proximal in X .*

ACKNOWLEDGEMENT

The authors would like to thank the referee for comments leading to a concise presentation of the results. The second-named author's research was supported by the CSIR Research Fellowship, and she would like to thank the CSIR for their financial support.

REFERENCES

- [1] E. M. Alfsen and E.G. Effros, Structure in real Banach spaces, Part I and II, Ann. of Math. 96 (1972) 98-173. MR0352946 (50:5432)
- [2] T. Ando, Closed range theorems for convex sets and linear liftings, Pacific J. Math. 44 (1973) 393-410. MR0328546 (48:6888)
- [3] E. Behrends, M -Structure and the Banach-Stone theorem, Lecture Notes in Math. 736, Springer, Berlin, Heidelberg, New York, 1979. MR0547509 (81b:46002)
- [4] M. Fabian, P. Habala, P. Hájek, V. M. Santalucia, J. Pelant and V. Zizler, Functional Analysis and Infinite-Dimensional Geometry, Springer, Berlin, Heidelberg, New York, 2001. MR1831176 (2002f:46001)
- [5] P. Harmand, D. Werner and W. Werner, M -ideals in Banach spaces and Banach algebras, Lecture Notes in Math. 1574. Springer, Berlin, Heidelberg, New York, 1993. MR1238713 (94k:46022)
- [6] Ka-Sing Lau, On a sufficient condition for proximity, Trans. Amer. Math. Soc. 251 (1979) 343-356. MR0531983 (81b:46033)

- [7] I. Singer, Best Approximation in Normed Linear spaces by Elements of Linear Subspaces, Springer, Berlin, Heidelberg, New York, 1970. MR0270044 (42:4937)
- [8] D. Yost, Best approximation and intersections of balls in Banach spaces, Bull. Austral. Math. Soc. 20 (1979) 285-300. MR0557239 (80m:46019)

DEPARTMENT OF MATHEMATICS, PONDICHERRY UNIVERSITY, KALAPET, PONDICHERRY-605014,
INDIA

E-mail address: `pdyindumath@gmail.com`

DEPARTMENT OF MATHEMATICS, PONDICHERRY UNIVERSITY, KALAPET, PONDICHERRY-605014,
INDIA

E-mail address: `s_lalithambigai@yahoo.co.in`