POWERS AND ROOTS OF TOEPLITZ OPERATORS

ISSAM LOUHICHI

(Communicated by Joseph A. Ball)

ABSTRACT. We study the commutativity of two Toeplitz operators whose symbols are quasihomogeneous functions. We give a relationship between this commutativity and the roots (or powers) of the Toeplitz operators. We use this to characterize Toeplitz operators with symbols in $L^\infty(\mathbb{D})$ which commute with Toeplitz operators whose symbols are of the form $e^{ip\theta}r^m$.

1. INTRODUCTION

Let $\mathbb{D}$ denote the open unit disk in the complex plane $\mathbb{C}$, and let dA denote normalized Lebesgue area measure. The Bergman space, denoted by L_a^2 , is the Hilbert space of analytic functions on $\mathbb{D}$ that are square integrable with respect to dA . It is well known that L_a^2 is a closed subspace of the Hilbert space $L^2(\mathbb{D}, dA)$ and $(\sqrt{n+1}z^n)_{n \in \mathbb{N}}$ is an orthonormal basis of L_a^2 . Let P be the orthogonal projection of $L^2(\mathbb{D}, dA)$ onto L_a^2 . For a function $\phi \in L^\infty(\mathbb{D}, dA)$, the Toeplitz operator with symbol ϕ is the operator T_ϕ from L_a^2 to L_a^2 defined by $T_\phi(f) = P(\phi f)$.

If $k_z(w) = \frac{1}{(1-\bar{z}w)^2} = \sum_{j=0}^{\infty} (1+j)w^j\bar{z}^j$ is the Bergman reproducing kernel, then

$$T_\phi(f)(z) = P(\phi f)(z) = \int_{\mathbb{D}} \phi(w)f(w)\overline{k_z(w)}dA(w).$$

The question to be studied in this paper is: When do two Toeplitz operators T_ϕ and T_ψ commute? In 1964, Brown and Halmos [4] solved this problem for the analogously defined Toeplitz operators on the Hardy space. They showed that $T_\phi T_\psi = T_\psi T_\phi$ for some ϕ and $\psi \in L^\infty(\mathbb{T})$, where $\mathbb{T}$ is the unit circle of $\mathbb{C}$, if and only if either

- (a) ϕ and ψ are both analytic, or
- (b) $\bar{\phi}$ and $\bar{\psi}$ are both analytic, or
- (c) one of the two symbols is a linear function of the other.

We recall that a function in $L^\infty(\mathbb{T})$ is said to be analytic if all of its Fourier coefficients with negative indices are equal to 0.

The same question concerning Toeplitz operators on the Bergman space has a much more complicated answer. There are however some results which resemble those of [4]. In fact, Axler and Čučković proved in [2] that the condition that one of (a), (b) or (c) be true is still necessary and sufficient when the two symbols ϕ and ψ are bounded harmonic functions on $\mathbb{D}$. Moreover, with Rao [3], they proved

Received by the editors March 26, 2005 and, in revised form, December 20, 2005.

2000 *Mathematics Subject Classification.* Primary 47B35; Secondary 47L80.

Key words and phrases. Toeplitz operators, Bergman space, Mellin transform.

©2006 American Mathematical Society
 Reverts to public domain 28 years from publication

that if ϕ is a bounded analytic function and if ψ is a bounded symbol such that T_ϕ and T_ψ commute, then ψ must be analytic too. When we consider arbitrary symbols, things are different. In [5] Čučković and Rao used the Mellin transform to study the commutativity of multiplication of two Toeplitz operators T_ϕ and T_ψ on the Bergman space and describe those operators which commute with $T_{e^{ip\theta}r^m}$ for $(m, p) \in \mathbb{N} \times \mathbb{N}$. In this paper we use our results from [7] to interpret and extend the results of [5]. We give some solutions in the case where the Toeplitz operators have symbols which are “quasihomogeneous” functions and show that these solutions are related to “ p^{th} roots” and powers of the Toeplitz operators.

As in [7] we say that a bounded symbol f is quasihomogeneous of degree k if it is of the form $e^{ik\theta}\phi$ where ϕ is a radial function. In this case we say that the Toeplitz operator T_f is quasihomogeneous of degree k .

2. PRELIMINARIES

The Mellin transform of a function $\psi \in L^1([0, 1], r dr)$ is defined by

$$\widehat{\psi}(z) = \int_0^1 \psi(r) r^{z-1} dr.$$

It is easy to see that $\widehat{\psi}$ is a bounded holomorphic function on the half-plane $\Pi = \{z : \Re z > 2\}$.

We denote the Mellin convolution of two functions ϕ and ψ by $\phi *_M \psi$ and we define it by the equation

$$(\phi *_M \psi)(r) = \int_r^1 \phi\left(\frac{r}{t}\right) \psi(t) \frac{dt}{t}.$$

It is clear that the Mellin transform converts Mellin convolution into a pointwise product, i.e., that

$$(1) \quad \widehat{(\phi *_M \psi)}(r) = \widehat{\phi}(r) \widehat{\psi}(r).$$

We shall often use the following classical theorem (see [8, p. 102]).

Theorem 1. *Suppose that f is a bounded, holomorphic function on $\{z : \Re z > 0\}$ which vanishes at the pairwise distinct points $d_1, d_2, \dots$, where*

- i) $\inf\{|d_n|\} > 0$ and
- ii) $\sum_{n \geq 1} \Re\left(\frac{1}{d_n}\right) = \infty$.

Then f vanishes identically on $\{z : \Re z > 0\}$.

Remark 2. We shall often apply this theorem to show that if $\psi \in L^1([0, 1], r dr)$ and if there exist $n_0 \in \mathbb{Z}_+$, $p \in \mathbb{N}$ such that

$$\widehat{\psi}(n_0 + pk) = 0 \text{ for all } k \in \mathbb{N},$$

then $\widehat{\psi}(z) = 0$ for all $z \in \{z : \Re z > 2\}$ and so $\psi = 0$.

3. POWERS OF TOEPLITZ OPERATORS

The following lemma determines the values of powers of a bounded quasihomogeneous Toeplitz operator evaluated at any element of the orthonormal basis of L_a^2 .

Lemma 3. *Let $n \in \mathbb{N}$, $s \in \mathbb{Z}_+$ and let ψ be a bounded radial function on $\mathbb{D}$. Then, for all $k \in \mathbb{N}$ we have*

$$\begin{aligned} \left(T_{e^{is\theta}\psi}\right)^n(\xi^k)(z) &= \left[\prod_{j=0}^{n-1} 2(k+js+s+1)\widehat{\psi}(2k+2js+s+2)\right]z^{k+ns} \\ &= \frac{\prod_{j=0}^{n-1} \widehat{\psi}(2k+2js+s+2)}{\prod_{j=0}^{n-1} \widehat{\mathbb{1}}(2k+2js+2s+2)}z^{k+ns}, \end{aligned}$$

where $\mathbb{1}$ denotes the constant function with value one.

Proof. The lemma is a consequence of the following direct calculation. We write

$$T_{e^{is\theta}\psi}(\xi^k)(z) = \int_0^1 \int_0^{2\pi} \psi(r)r^k \sum_{j=0}^{\infty} (j+1)e^{i(k+s-j)\theta} r^j z^j \frac{1}{\pi} r dr d\theta$$

and interchange the integral over $[0, 2\pi]$ and the sum to see that

$$\begin{aligned} T_{e^{is\theta}\psi}(\xi^k)(z) &= 2(k+s+1)\widehat{\psi}(2k+s+2)z^{k+s} \\ &= \frac{\widehat{\psi}(2k+s+2)}{\widehat{\mathbb{1}}(2k+2s+2)}z^{k+s}. \end{aligned}$$

The lemma is proved by applying $T_{e^{is\theta}\psi}$ to ξ^k n times. □

We have the following decomposition of $L^2(\mathbb{D}, dA)$ as

$$L^2(\mathbb{D}, dA) = \bigoplus_{k \in \mathbb{Z}} e^{ik\theta} \mathcal{R},$$

where $\mathcal{R}$ is the space of functions on $[0, 1]$ that are square integrable with respect to the measure rdr . Thus every function $f \in L^2(\mathbb{D}, dA)$ has the decomposition

$$f(re^{ik\theta}) = \sum_{k=-\infty}^{+\infty} e^{ik\theta} f_k(r), \quad f_k \in \mathcal{R}.$$

Moreover, if $f \in L^\infty(\mathbb{D}, dA) \subset L^2(\mathbb{D}, dA)$, then for each $r \in [0, 1]$,

$$|f_k(r)| = \frac{1}{2\pi} \left| \int_0^{2\pi} f(re^{i\theta}) e^{-ik\theta} d\theta \right| \leq \sup_{z \in \mathbb{D}} |f(z)|, \quad \forall k \in \mathbb{Z},$$

and so the functions f_k are bounded in the disk.

In [7] we proved the following results, which we will use in the proof of our main theorem.

Proposition 4. *Let ϕ be a nonzero bounded radial function, let p be a positive integer and let $f(re^{ik\theta}) = \sum_{k=-\infty}^{+\infty} e^{ik\theta} f_k(r) \in L^\infty(\mathbb{D}, dA)$. Then*

- a) T_f commutes with $T_{e^{ip\theta}\phi}$ if and only if $T_{e^{ik\theta}f_k}$ commutes with $T_{e^{ip\theta}\phi}$ for all $k \in \mathbb{Z}$.
- b) If there exists $k \in \mathbb{Z}_-$ and a bounded radial function f_k such that

$$T_{e^{ip\theta}\phi} T_{e^{ik\theta}f_k} = T_{e^{ik\theta}f_k} T_{e^{ip\theta}\phi},$$

then f_k must be equal to zero.

- c) If there exists $k \in \mathbb{Z}_+$ and a bounded radial function f_k such that

$$T_{e^{ip\theta}\phi} T_{e^{ik\theta}f_k} = T_{e^{ik\theta}f_k} T_{e^{ip\theta}\phi},$$

then f_k is unique up to a constant factor. In particular f_0 is a constant.

Thus if $p > 0$, $f(re^{ik\theta}) = \sum_{k=-\infty}^{+\infty} e^{ik\theta} f_k(r)$ and T_f commutes with $T_{e^{ip\theta}\phi}$, then each f_k is uniquely determined up to multiplication by a constant and is equal to 0 for $k < 0$.

Next we present two technical but easy results which permit us to prove Propositions 7 and 9, the principal results of this section.

Remark 5. Let $(a_l)_{l \in \mathbb{N}}$ and $(b_l)_{l \in \mathbb{N}}$ be two nonvanishing sequences and p and s two positive integers such that

$$(2) \quad a_{l+s}b_l = b_{l+p}a_l \quad \text{for all } l \in \mathbb{N}.$$

Then if

$$A_k = \prod_{j=0}^{s-1} a_{k+jp} \quad \text{and} \quad B_k = \prod_{j=0}^{p-1} b_{k+js},$$

we have

$$A_k B_{k+p} = A_{k+p} B_k \quad \text{for all } k \in \mathbb{N}.$$

(Just multiply the p equations obtained by taking $l = k, k+s, \dots, k+(p-1)s$ in (2) together to see that, if (2) is true, then

$$\frac{B_{k+p}}{B_k} = \frac{a_{k+ps}}{a_k} = \frac{A_{k+p}}{A_k} \quad \text{for all } k \in \mathbb{N}.)$$

Notation. Let S and T be two functions (resp. two operators). We will say that $S \equiv T$ if there exists a constant $c \neq 0$ such that $S = cT$.

Lemma 6. *Let F and G be two nonzero bounded holomorphic functions on the half plane $\Pi = \{z : \Re z > 2\}$. If there exists $p \in \mathbb{N}$ such that*

$$(3) \quad F(z)G(z+p) = F(z+p)G(z) \quad \text{for all } z \in \Pi,$$

then $F \equiv G$.

Proof. Suppose that (3) is true. Then, if (as above) we multiply the k equations obtained by taking $z_n = z + np$ for $n = 0, \dots, k-1$, we have

$$(4) \quad F(z)G(z+kp) = F(z+kp)G(z) \quad \text{for all } k \in \mathbb{N}.$$

Now, let $z_0 \in \Pi$ such that $G(z_0) \neq 0$ and let $E = \{k \in \mathbb{N} : G(z_0 + kp) = 0\}$. If $\sum_{k \in E} \Re(\frac{1}{|z_0 + kp|}) = \infty$, then Theorem 1 implies that $G = 0$. This contradicts the hypothesis of the lemma. Thus $\sum_{k \in E^c} \Re(\frac{1}{|z_0 + kp|}) = \infty$, where E^c is the complement in $\mathbb{N}$ of the set E .

Now, equation (4) implies that

$$\frac{F(z_0 + kp)}{G(z_0 + kp)} = \frac{F(z_0)}{G(z_0)} \quad \text{for all } k \in E^c.$$

So, applying Theorem 1 to the function $F - cG$, where $c = \frac{F(z_0)}{G(z_0)}$, completes the proof. $\square$

Let p and s be two positive integers and ψ a bounded radial function.

If $(T_{e^{is\theta}\psi})^p$ is a Toeplitz operator, then it is the unique quasihomogeneous Toeplitz operator of degree ps (see Proposition 3 and Proposition 4 of [7]) which commutes with $T_{e^{is\theta}\psi}$. It is natural to ask whether all nonzero Toeplitz operators which are of quasihomogeneous of degree a multiple of s and which commute with $T_{e^{is\theta}\psi}$ are of this form.

Proposition 7. *Let p and s be two positive integers and ϕ and ψ be two nonzero bounded radial functions such that*

$$(5) \quad T_{e^{ip\theta}\phi} T_{e^{is\theta}\psi} = T_{e^{is\theta}\psi} T_{e^{ip\theta}\phi}.$$

Then

$$(6) \quad \left(T_{e^{ip\theta}\phi}\right)^s \equiv \left(T_{e^{is\theta}\psi}\right)^p.$$

Proof. For all $k \in \mathbb{N}$, let

$$a_k = \frac{\widehat{\phi}(2k+p+2)}{\widehat{\mathbb{1}}(2k+2p+2)} \quad \text{and} \quad b_k = \frac{\widehat{\psi}(2k+s+2)}{\widehat{\mathbb{1}}(2k+2s+2)}$$

so that

$$T_{e^{ip\theta}\phi}(\xi^k)(z) = a_k z^{k+p} \quad \text{and} \quad T_{e^{is\theta}\psi}(\xi^k)(z) = b_k z^{k+s}.$$

Then equation (5) shows that $a_{k+s}b_k = b_{k+p}a_k$ for all $k \in \mathbb{Z}_+$, and so Remark 5 implies that

$$(7) \quad \prod_{j=0}^{s-1} a_{k+jp} \prod_{j=0}^{p-1} b_{k+p+js} = \prod_{j=0}^{s-1} a_{k+p+jp} \prod_{j=0}^{p-1} b_{k+js}.$$

Let F and G be the two bounded holomorphic functions defined for all $z \in \Pi$ by

$$F(z) = \prod_{j=0}^{p-1} \widehat{\mathbb{1}}(z+2js+2s) \prod_{j=0}^{s-1} \widehat{\phi}(z+2jp+p)$$

and

$$G(z) = \prod_{j=0}^{s-1} \widehat{\mathbb{1}}(z+2jp+2p) \prod_{j=0}^{p-1} \widehat{\psi}(z+2js+s).$$

Then equation (7) is equivalent to

$$F(2k+2)G(2k+2p+2) = F(2k+2p+2)G(2k+2) \quad \text{for all } k \in \mathbb{Z}_+.$$

Now, applying Theorem 1, in the form of Remark 2, implies that

$$F(z)G(z+2p) = F(z+2p)G(z) \quad \text{for all } z \in \Pi.$$

Finally, using Lemma 6, we obtain that

$$\frac{\prod_{j=0}^{s-1} \widehat{\phi}(z+2jp+p)}{\prod_{j=0}^{s-1} \widehat{\mathbb{1}}(z+2jp+2p)} \equiv \frac{\prod_{j=0}^{p-1} \widehat{\psi}(z+2js+s)}{\prod_{j=0}^{p-1} \widehat{\mathbb{1}}(z+2js+2s)} \quad \text{for all } z \in \Pi,$$

and Lemma 3 completes the proof. $\square$

Remark 8. i) We will assume that $(T_{e^{ip\theta}\phi})^0 = I$, where I is the identity operator of L_a^2 onto L_a^2 .

ii) If p and s are both negative integers and if $T_{e^{ip\theta}\phi} T_{e^{is\theta}\psi} = T_{e^{is\theta}\psi} T_{e^{ip\theta}\phi}$, then by considering the adjoint operators we obtain

$$T_{e^{-is\theta}\psi} T_{e^{-ip\theta}\phi} = T_{e^{-ip\theta}\phi} T_{e^{-is\theta}\psi}$$

and so Proposition 7 implies that $(T_{e^{-ip\theta}\phi})^{-s} \equiv (T_{e^{-is\theta}\psi})^{-p}$.

Now, by considering once again the adjoint operators we see that

$$(T_{e^{ip\theta}\phi})^{-s} \equiv (T_{e^{is\theta}\psi})^{-p}.$$

Proposition 9. *Let ϕ and ψ be two nonzero bounded radial functions and n, p and s be positive integers. Then*

$$(T_{e^{ips\theta}\phi})^n = (T_{e^{is\theta}\psi})^{np} \implies T_{e^{ips\theta}\phi} = (T_{e^{is\theta}\psi})^p.$$

Proof. For all $k \in \mathbb{Z}_+$, let

$$a_k = 2(k + ps + 1)\widehat{\phi}(2k + ps + 2) \quad \text{and} \quad b_k = 2(k + s + 1)\widehat{\psi}(2k + s + 2),$$

so that

$$(T_{e^{ips\theta}\phi})^n = (T_{e^{is\theta}\psi})^{np} \Leftrightarrow \prod_{j=0}^{n-1} a_{k+jps} = \prod_{j=0}^{np-1} b_{k+js} \quad \text{for all } k \in \mathbb{Z}_+$$

and

$$T_{e^{ips\theta}\phi} = (T_{e^{is\theta}\psi})^p \Leftrightarrow a_k = \prod_{j=0}^{p-1} b_{k+js} \quad \text{for all } k \in \mathbb{Z}_+.$$

Suppose that

$$(8) \quad \prod_{j=0}^{n-1} a_{k+jps} = \prod_{j=0}^{np-1} b_{k+js} \quad \text{for all } k \in \mathbb{Z}_+.$$

We will prove that

$$(9) \quad a_{knps} = \prod_{j=0}^{p-1} b_{knps+js} \quad \text{for all } k \in \mathbb{Z}_+.$$

We prove (9) by induction on k . If we take $k = 0$ in equation (8), then we obtain

$$\begin{aligned} \prod_{j=0}^{n-1} a_{jps} &= \prod_{j=0}^{np-1} b_{js} \\ &= \prod_{j=0}^{p-1} b_{js} \prod_{j=p}^{np-1} b_{js} \\ &= \prod_{j=0}^{p-1} b_{js} \prod_{j=0}^{np-1} b_{ps+js}. \end{aligned}$$

Otherwise

$$\begin{aligned} \prod_{j=0}^{n-1} a_{jps} &= a_0 \prod_{j=1}^{n-1} a_{jps} \\ &= a_0 \prod_{j=0}^{n-1} a_{ps+jps}. \end{aligned}$$

But equation (8) implies that

$$\prod_{j=0}^{n-1} a_{ps+jps} = \prod_{j=0}^{np-1} b_{ps+js}.$$

Thus

$$a_0 = \prod_{j=0}^{p-1} b_{js}.$$

Now, assume (9) is true for $knps$. We show it is true for $(k+1)nps$. We set k equal to $knps$ in the left-hand side of (8) and obtain

$$\prod_{j=0}^{n-1} a_{knps+js} = a_{knps} \prod_{j=0}^{n-2} a_{knps+ps+js}.$$

Then

$$a_{(k+1)nps} \prod_{j=0}^{n-1} a_{knps+js} = a_{knps} \prod_{j=0}^{n-1} a_{knps+ps+js}.$$

But

$$\prod_{j=0}^{n-1} a_{knps+ps+js} = \prod_{j=0}^{np-1} b_{knps+ps+js}$$

and

$$\prod_{j=0}^{np-1} b_{knps+ps+js} \prod_{j=0}^{p-1} b_{knps+js} = \prod_{j=0}^{np-1} b_{knps+js} \prod_{j=0}^{p-1} b_{(k+1)nps+js}.$$

Thus (9) is proved for $(knps)_{k \in \mathbb{Z}_+}$. Hence, for all $k \in \mathbb{Z}_+$, we have

$$\widehat{\phi}(2knps+ps+2) \prod_{j=0}^{p-1} \widehat{1}(2knps+2js+2s+2) = \widehat{1}(2knp+p+2) \prod_{j=0}^{p-1} \widehat{\psi}(2knp+1+2)$$

and, using equation (1) and Remark 2, we complete the proof. $\square$

Remark 10. In [7] (Proposition 6) we prove that if $p > 0$ and ϕ is a nonzero bounded radial function and if there exists a bounded radial function ψ such that T_ψ commutes with $T_{e^{ip\theta}\phi}$, then ψ must be a constant. Here is another proof of this proposition. In fact, using Proposition 7, we have $(T_\psi)^p \equiv I$, so Proposition 9 implies that $T_\psi \equiv I$, and so, that $\psi \equiv 1$ since I is the Toeplitz operator of symbol 1 .

4. MAIN RESULT

Let p be a positive integer. We start this section with the definition of the T - p^{th} root of a quasihomogeneous Toeplitz operator of degree p or $-p$. This new notion plays an important role in the remainder of the paper.

Definition 11. Let ϕ be a nonzero bounded radial function and p be a positive integer. We say that the Toeplitz operator $T_{e^{ip\theta}\phi}$ has a T - p^{th} root $T_{e^{i\theta}\psi}$ if and only if there exists a nonzero bounded radial function ψ such that

$$T_{e^{ip\theta}\phi} = (T_{e^{i\theta}\psi})^p.$$

Remark 12. i) The T - p^{th} root of a quasihomogeneous Toeplitz operator is unique. In fact, suppose that $T_{e^{ip\theta}\phi}$ has two T - p^{th} roots $T_{e^{i\theta}\psi}$ and $T_{e^{i\theta}\tilde{\psi}}$. Then $(T_{e^{i\theta}\psi})^p = (T_{e^{i\theta}\tilde{\psi}})^p$. Then, by Proposition 9, we have that $T_{e^{i\theta}\psi} = T_{e^{i\theta}\tilde{\psi}}$, which implies that $\psi = \tilde{\psi}$.
ii) If the quasihomogeneous degree is negative we have an analogous definition of the T - p^{th} root. Let p be a positive integer and ϕ be a bounded radial function. Then, we say that $T_{e^{-ip\theta}\phi}$ has a T - p^{th} root if there exists a bounded radial function ψ such that $T_{e^{-ip\theta}\phi} = (T_{e^{-i\theta}\psi})^p$. It is easy to see,

by taking adjoints, that $T_{e^{-ip\theta}\phi}$ has a T - p^{th} root $T_{e^{-i\theta}\psi}$ if and only if $T_{e^{ip\theta}\phi}$ has a T - p^{th} root $T_{e^{i\theta}\psi}$.

Examples. i) $T_{e^{i\theta}(\frac{r+r^5}{2})}$ is the T - 2^{th} root of $T_{e^{2i\theta}r^6}$.

ii) $T_{e^{i\theta}(\frac{3r+2r^5+3r^9}{8})}$ is the T - 2^{th} root of $T_{e^{2i\theta}r^{10}}$.

Now, if $T_{e^{i\theta}\psi}$ is the T - p^{th} root of $T_{ip\theta\phi}$ and if $(T_{e^{i\theta}\psi})^k$ (for k in $\mathbb{N}$) is a Toeplitz operator, then $(T_{e^{i\theta}\psi})^k$ is the unique nonzero quasihomogeneous Toeplitz operator of degree k which can commute with $T_{e^{ip\theta}\phi}$. What we prove below is that if $T_{e^{ip\theta}\phi}$ has a T - p^{th} root $T_{e^{i\theta}\psi}$, then the *only* nonzero quasihomogeneous Toeplitz operator of degree s which commutes with $T_{e^{ip\theta}\phi}$ is an s^{th} power of $T_{e^{i\theta}\psi}$, extending the result (Propositions 7 and 9) of section 3 in this case.

Theorem 13. *Let ϕ be a nonzero bounded radial function and let p be a positive integer. Assume that $T_{e^{ip\theta}\phi}$ has a T - p^{th} root $T_{e^{i\theta}\psi}$. Suppose that*

$$f(re^{i\theta}) = \sum_{k=-\infty}^{+\infty} e^{ik\theta} f_k(r) \in L^\infty(\mathbb{D}, dA)$$

is such that

$$(10) \quad T_f T_{e^{ip\theta}\phi} = T_{e^{ip\theta}\phi} T_f.$$

Then

- i) $f_k = 0$ for $k < 0$.
- ii) If $k \geq 0$ and $(T_{e^{i\theta}\psi})^k$ is a Toeplitz operator, then either $T_{e^{ik\theta}f_k} \equiv (T_{e^{i\theta}\psi})^k$ or $f_k = 0$.
- iii) If $k \geq 0$ and $(T_{e^{i\theta}\psi})^k$ is not a Toeplitz operator, then $f_k = 0$.

Proof. Assertion a) of Proposition 4 implies that if equation (10) is true, then

$$T_{e^{ik\theta}f_k} T_{e^{ip\theta}\phi} = T_{e^{ip\theta}\phi} T_{e^{ik\theta}f_k}, \text{ for all } k \in \mathbb{Z}.$$

Thus i) is a direct consequence of assertion b) of Proposition 4.

Now, to prove ii), let k be a positive integer such that $(T_{e^{i\theta}\psi})^k$ is a Toeplitz operator. Then $(T_{e^{i\theta}\psi})^k$ is a quasihomogeneous Toeplitz operator of degree k which commutes with $T_{e^{ip\theta}\phi}$. So, if f_k is not identically equal to zero, then f_k is a bounded nonzero radial function such that $T_{e^{ik\theta}f_k}$ commutes with $T_{e^{ip\theta}\phi}$. Thus, assertion c) of Proposition 4 implies that $T_{e^{ik\theta}f_k} \equiv (T_{e^{i\theta}\psi})^k$.

Finally, let k be a positive integer such that $(T_{e^{i\theta}\psi})^k$ is not a Toeplitz operator and suppose that there exists a nonzero bounded radial function f_k such that $T_{e^{ik\theta}f_k}$ commutes with $T_{e^{ip\theta}\phi}$. Then Proposition 7 implies that

$$(T_{e^{ik\theta}f_k})^p \equiv (T_{e^{ip\theta}\phi})^k.$$

Thus $(T_{e^{ik\theta}f_k})^p \equiv (T_{e^{i\theta}\psi})^{kp}$ and Proposition 9 implies that $T_{e^{ik\theta}f_k} \equiv (T_{e^{i\theta}\psi})^k$, which contradicts our hypothesis. This proves iii). $\square$

Before starting with corollaries, we state an interesting theorem which follows from [5] and give an idea of its proof. In fact we will apply this theorem to see that if p is any positive integer and m is any nonnegative integer, then the Toeplitz operator $T_{e^{ip\theta}r^m}$ always has a T - p^{th} root.

Theorem 14. *Let $p \geq 1$ and $m \geq 0$ be two integers. For all integers s , such that $1 \leq s < p$, there exists a unique bounded radial function ψ such that*

$$(11) \quad T_{e^{is\theta}\psi} T_{e^{ip\theta}r^m} = T_{e^{ip\theta}r^m} T_{e^{is\theta}\psi}.$$

Proof. (This is a slight variation of the proof found in [5].) If $m \geq 0, p \geq 1$ and $1 \leq s < p$, we define the radial functions f and g by

$$f(r) = 2pr^{2s}(1-r^{2p})^{-\frac{s}{p}} \quad \text{and} \quad g(r) = 2pr^{m+p}(1-r^{2p})^{\frac{s}{p}-1}.$$

Let ψ be the radial function defined by

$$r^s \psi = f *_M g.$$

Čučković and Rao prove, using a long rather technical calculation, that ψ is bounded. Here, we will show that ψ satisfies (11). To do this, we need only verify that for $k \in \mathbb{Z}_+$

$$\frac{2k+2p+2}{2k+m+p+2} \widehat{r^s \psi}(2k+2p+2) = \frac{2k+2s+2}{2k+m+p+2s+2} \widehat{r^s \psi}(2k+2).$$

By (1), we have $\widehat{r^s \psi}(2k+2) = \widehat{f}(2k+2)\widehat{g}(2k+2)$. A simple substitution $t = r^{2p}$ shows that

$$\widehat{f}(2k+2) = B\left(\frac{2k+2s+2}{2p}, 1 - \frac{s}{p}\right) \quad \text{and} \quad \widehat{g}(2k+2) = B\left(\frac{2k+m+p+2}{2p}, \frac{s}{p}\right),$$

where B denotes the beta function. Using the well-known identities $B(z_1, z_2) = \frac{\Gamma(z_1)\Gamma(z_2)}{\Gamma(z_1+z_2)}$ and $\Gamma(1+z) = z\Gamma(z)$, where Γ is the gamma function, it is easy to see that

$$(12) \quad \widehat{r^s \psi}(2k+2p+2) = \frac{(2k+2s+2)(2k+m+p+2)}{(2k+2p+2)(2k+m+p+2s+2)} \widehat{r^s \psi}(2k+2),$$

which finishes the proof. $\square$

Remark 15. i) It is trivial that $T_{e^{ip\theta}r^m}$ commutes with itself. So, if $p = s$, assertion c) of Proposition 4 implies that $\psi \equiv r^m$.
ii) We wish to highlight the following case. If $m = (2n+1)p$ for $n \in \mathbb{N}$, then the function ψ exists for all $s \in \mathbb{N}$. In fact, if we substitute $m = (2n+1)p$ in (12) and use Theorem 1, we obtain for all $z \in \Pi$,

$$\frac{\widehat{r^s \psi}(z+2p)}{\widehat{r^s \psi}(z)} = \frac{F(z+2p)}{F(z)}, \quad \text{where } F(z) = \frac{\Gamma(\frac{z+2s}{2p})\Gamma(\frac{z}{2p}+n+1)}{\Gamma(\frac{z}{2p}+1)\Gamma(\frac{z+2s}{2p}+n+1)}.$$

Now, using the identity $\Gamma(1+z) = z\Gamma(z)$ repeatedly, we have

$$F(z) = 2p \frac{\prod_{j=0}^{n-1} (z+2jp+2p)}{\prod_{j=0}^n (z+2jp+2s)},$$

which is a proper fraction in z and can be written as

$$(13) \quad F(z) = \sum_{j=0}^n \frac{a_j}{z+2jp+2s}.$$

Since $\frac{1}{z+2jp+2s} = \widehat{r^{2jp+2s}}(z)$, it follows by Lemma 6 that

$$\widehat{r^s \psi}(z) \equiv \sum_{j=0}^n a_j \widehat{r^{2jp+2s}}(z),$$

where the a_j are defined by (13), and so Theorem 1 implies that

$$\psi(r) \equiv \sum_{j=0}^n a_j r^{2jp+s}.$$

Next, we give some easy but interesting consequences of Theorem 14.

Corollary 16. *For all integers $m \geq 0$, $p \geq 1$, and $s \geq 1$ there exists a bounded radial function ψ such that $(T_{e^{is\theta}\psi})^p \equiv T_{e^{ips\theta}r^m}$.*

Proof. Let $m \geq 0$, $p \geq 1$, and $s \geq 1$ be integers. Theorem 14 implies that there exists a bounded radial function ψ such that

$$T_{e^{is\theta}\psi} T_{e^{ips\theta}r^m} = T_{e^{ips\theta}r^m} T_{e^{is\theta}\psi}.$$

Using Proposition 7 we have $(T_{e^{is\theta}\psi})^{ps} \equiv (T_{e^{ips\theta}r^m})^s$ and so, an application of Proposition 9 finishes the proof. $\square$

In [4], Brown and Halmos studied multiplicativity of Toeplitz operators on the Hardy space and showed that the product of two Toeplitz operators T_f and T_g is equal to a third Toeplitz operator T_h for some f, g and h in $L^\infty(\mathbb{T})$ if and only if f is conjugate analytic or g is analytic, that is, hardly ever. The question of when the product of two Toeplitz operators on the Bergman space is equal to a third is much more complicated and still open. Most work on this question shows that it is not often true that the product of two Toeplitz operators is a Toeplitz operator (see [1] and [6]). But, below, we show that, for certain nontrivial Toeplitz operators $T_{e^{i\theta}\psi}$, not only is $(T_{e^{i\theta}\psi})^2$ equal to a Toeplitz operator, but there exists a positive integer k such that $(T_{e^{i\theta}\psi})^i$ is a Toeplitz operator for all positive integers $i \leq k$.

Corollary 17. *Let $m \geq 0$ and $p \geq 1$ be two integers. If $T_{e^{ip\theta}r^m}$ has a T - p^{th} root $T_{e^{i\theta}\psi}$ then, for all integers k with $1 \leq k \leq p$, the product $(T_{e^{i\theta}\psi})^k$ is a Toeplitz operator.*

Proof. Let k be an integer such that $1 \leq k \leq p$. By Theorem 14 we know that there exists a bounded radial function ϕ such that $T_{e^{ik\theta}\phi}$ commutes with $T_{e^{ip\theta}r^m}$. So, Proposition 7 implies that

$$(T_{e^{ik\theta}\phi})^p \equiv (T_{e^{ip\theta}r^m})^k.$$

Thus $(T_{e^{ik\theta}\phi})^p = (T_{e^{i\theta}\psi})^{kp}$ since $T_{e^{i\theta}\psi}$ is the T - p^{th} root of $T_{e^{ip\theta}r^m}$, and so Proposition 9 finishes the proof. $\square$

It is easily seen that if f is a bounded analytic function on $\mathbb{D}$, then T_f is just a multiplication operator. Thus for any integer $k \geq 1$, it is clear that $(T_f)^k$ is a Toeplitz operator of symbol f^k . By taking adjoints, we can see that the powers of a Toeplitz operator with conjugate analytic symbol are also Toeplitz operators. These are the trivial cases. The next corollary says there are nontrivial symbols f such that $(T_f)^k$ is always a Toeplitz operator for all integers $k \geq 1$.

Corollary 18. *There exist bounded radial functions ψ such that for all integers $k \geq 1$ the product $(T_{e^{i\theta}\psi})^k$ is still a Toeplitz operator.*

Proof. Let $n \geq 0$ and $p \geq 1$ be two integers. By Theorem 14 we know that the Toeplitz operator $T_{e^{ip\theta}r^{(2n+1)p}}$ has a T - p^{th} root $T_{e^{i\theta}\psi}$, where ψ is a bounded radial function. Moreover the assertion *ii*) of Remark 15 tells us that, for all integers $k \geq 1$, there exists a bounded radial function ψ_k such that $T_{e^{ik\theta}\psi_k}$ commutes with $T_{e^{ip\theta}r^{(2n+1)p}}$. Thus Proposition 7 implies that $(T_{e^{ik\theta}\psi_k})^p \equiv (T_{e^{i\theta}\psi})^{kp}$ since $T_{e^{ip\theta}r^{(2n+1)p}} = (T_{e^{i\theta}\psi})^p$ and, again, Proposition 9 finishes the proof. $\square$

ACKNOWLEDGMENTS

I am thankful to Elizabeth Strouse my thesis supervisor for helpful discussions and suggestions concerning the results of this paper.

REFERENCES

- [1] P. Ahern, Ž. Čučković, A Theorem of Brown-Halmos Type for Bergman Space Toeplitz Operators, *J. Funct. Anal.* **187** (2001), 200-210. MR1867348 (2002h:47040)
- [2] S. Axler, Ž. Čučković, Commuting Toeplitz operators with harmonic symbols, *Integral Equations and Operator Theory* **14** (1991), 1-12. MR1079815 (92f:47018)
- [3] S. Axler, Ž. Čučković, N. V. Rao, Commutants of analytic Toeplitz operators on the Bergman space, *Proc. Amer. Math. Soc.* **128** (2000), 1951-1953. MR1694299 (2000m:47035)
- [4] A. Brown, P. Halmos, Algebraic properties of Toeplitz operators, *J. Reine Angew. Math.* **213** (1964), 89-102. MR0160136 (28:3350)
- [5] Ž. Čučković, N. V. Rao, Mellin transform, monomial symbols, and commuting Toeplitz operators, *J. Funct. Anal.* **154** (1) (1998), 195-214. MR1616532 (99f:47033)
- [6] I. Louhichi, E. Strouse, L. Zakarias, Products of Toeplitz operators on the Bergman space, *Integral Equations and Operator Theory*. Published on line: 1 October 2005.
- [7] I. Louhichi, L. Zakarias, On Toeplitz operators with quasihomogeneous symbols, *Archiv der Mathematik*. Volume **85**, Number 3 (2005), 248-257. MR2172383 (2006e:47061)
- [8] R. Remmert, *Classical Topics in Complex Function Theory*, Graduate Texts in Mathematics, Springer, New York, 1998. MR1483074 (98g:30002)

UFR MATHÉMATIQUES ET INFORMATIQUE, UNIVERSITÉ BORDEAUX 1, 351 COURS DE LA LIBÉRATION, 33405 TALENCE, FRANCE

E-mail address: louhichi@math.u-bordeaux1.fr