HANKEL OPERATORS WITH UNBOUNDED SYMBOLS

P. AHERN AND E. H. YOUSSEFI

(Communicated by Joseph A. Ball)

ABSTRACT. We prove that there are holomorphic functions f in the Hardy space of the unit ball or the bidisc such that the big Hankel operator with symbol $\bar{f}$ is bounded and for any holomorphic function g the function $\bar{f} + g$ cannot be bounded.

1. INTRODUCTION

Throughout this note $\Omega = \Omega_n$ will denote the domain in $\mathbb{C}^n$ which is either the unit ball $\mathbb{B}_n$ or the unit polydisc $\mathbb{D}^n$, where $\mathbb{D}$ is the unit disc in $\mathbb{C}$. We let σ denote the normalized Lebesgue measure on the Shilov boundary $\partial_s \Omega$ of Ω . For the unit ball then $\partial_s \mathbb{B}_n := \mathbb{S}$ is the unit sphere and for the unit polydisc, $\partial_s \mathbb{D}^n = \mathbb{T}^n$ where $\mathbb{T}$ is the unit circle. Let $H^2(\Omega)$ be the Hardy space on Ω with respect to σ . The Berezin transform of a function $f \in L^2(\sigma)$ is defined by

$$\tilde{f}(z) := \int_{\partial_s \Omega} f(\xi) P(z, \xi) d\sigma(\xi), \quad z \in \Omega,$$

where $P(z, \xi)$ is the Poisson kernel of Ω with respect to σ . We follow Coburn [C] to define the space $\widetilde{BMO}(\partial\Omega)$ of functions with bounded mean oscillation with respect to σ as the subspace of functions $f \in L^2(\sigma)$ that satisfy

$$\sup_{z \in \Omega} \left[|\tilde{f}|^2(z) - |f|^2(z) \right] < \infty.$$

We point out that in the one-dimensional case this definition gives the classical BMO space. See [Z1]. The subspace of $\widetilde{BMO}(\partial\Omega)$ consisting of those analytic functions on Ω will be denoted by $\widetilde{BMOA}(\partial\Omega)$. In the case of the unit ball the space $\widetilde{BMOA}(\mathbb{B})$ was shown to be the dual of the Hardy space $H^1(\mathbb{B})$. See Theorem 5.13 of [Z2]. For the polydisc case we show in Proposition 3.5 below that $\widetilde{BMO}$ coincides with the small bmo defined by Cotlar and Sadosky [CS]. It is not the dual of the real $H^1(\mathbb{T}^2)$, and its predual is given by Theorem 1.5 of [CS].

The big Hankel operator H_f with symbol $f \in L^2(\sigma)$ is defined by

$$H_f(g) := (I - P)(fg), \quad g \in H^\infty(\Omega).$$

We recall that in the one variable setting $\Omega = \mathbb{D}$, if a function f is in $H^2(\mathbb{D})$, then the big Hankel operator $H_{\bar{f}}$ is bounded on $H^2(\mathbb{T})$ if and only if there is another $g \in H^2(\mathbb{T})$ such that $\bar{f} + g$ is bounded. See [Z1]. The goal of this note is to show this

Received by the editors November 24, 2005 and, in revised form, February 16, 2006.
 2000 *Mathematics Subject Classification*. Primary 47B35, 32A35, 32A25.

is no longer true in higher dimensions. In particular, this answers in the negative a question by E. Strouse [S] which she raised for the case of the bidisc. We shall establish the following.

Theorem A. *There are functions $f \in H^2(\Omega_2)$ such that the big Hankel operator $H_{\bar{f}}$ is bounded on $H^2(\Omega_2)$ but $\bar{f} + g$ is unbounded for any holomorphic function g on Ω_2 .*

We point out that the existence of big Hankel operators with not necessarily antianalytic unbounded symbols was established in [CS] and [BT] for the case of the bidisc.

It is not hard to see [C] that if $f \in L^2(\partial_s \Omega)$ and both operators H_f and $H_{\bar{f}}$ are bounded, then $f \in \widetilde{BMO}(\partial_s \Omega)$. The converse is true for the bidisc. Indeed,

Theorem B. *Let $f \in L^2(\mathbb{T}^2)$. Then $f \in \widetilde{BMO}(\mathbb{T}^2)$ if and only if the big Hankel operators H_f and $H_{\bar{f}}$ are bounded on $H^2(\mathbb{D}^2)$.*

The analogue of Theorem B for the unit ball in $\mathbb{C}^2$ remains open.

2. THE BALL CASE

We recall that the Bergman distance $\beta(\zeta, z)$ between two points ζ and z in the unit disc $\mathbb{D}$ in the complex plane is given by

$$\beta(\zeta, z) = \frac{1}{2} \log \frac{1 + |\varphi_z(\zeta)|}{1 - |\varphi_z(\zeta)|}, \text{ where } \varphi_z(\zeta) := \frac{\zeta - z}{1 - \bar{\zeta}z}.$$

Lemma 2.1. *There is a positive constant C independent of $n > 0$ such that*

$$I_n(z) := \int_{\mathbb{D}} (1 - |\zeta|^2)^{\frac{n}{2}} \frac{\beta(\zeta, z)}{|1 - \bar{\zeta}z|^{n+2}} dA(\zeta) \leq \frac{C}{(n+1)^{\frac{3}{2}}} (1 - |z|^2)^{-\frac{n}{2}}$$

for all $z \in \mathbb{D}$.

Proof. Setting $\zeta = \varphi_z(w)$ we see that

$$\begin{aligned} I_n(z) &= \int_{\mathbb{D}} \beta(0, w) \frac{(1 - |w|^2)^{\frac{n}{2}}}{|1 - \bar{\zeta}w|^2} dA(w) \\ &= \frac{(1 - |z|^2)^{-\frac{n}{2}}}{2} \int_0^1 r \log \frac{1+r}{1-r} (1 - r^2)^{\frac{n}{2}} \frac{1}{2\pi} \int_0^{2\pi} \frac{d\theta}{|1 - r\bar{z}e^{i\theta}|^2} dr \\ &= \frac{(1 - |z|^2)^{-\frac{n}{2}}}{2} \int_0^1 r \frac{(1 - r^2)^{\frac{n}{2}}}{1 - r^2|z|^2} \log \frac{1+r}{1-r} dr \\ &\leq \frac{(1 - |z|^2)^{-\frac{n}{2}}}{2} \int_0^1 r (1 - r^2)^{\frac{n}{2}-1} \log \frac{1+r}{1-r} dr \\ &= \frac{(1 - |z|^2)^{-\frac{n}{2}}}{2} \left(\int_0^1 r \log(1+r) (1 - r^2)^{\frac{n}{2}-1} dr \right. \\ &\quad \left. + \int_0^1 r (1 - r^2)^{\frac{n}{2}-1} \log \frac{1}{1-r} dr \right). \end{aligned}$$

Integration by parts and the change of variable $t = 1 - r$ give that the second integral in the last equality is equal to $4n^{-2}$. To estimate the first integral in terms

of n , we have

$$\begin{aligned} \int_0^1 r \log(1+r)(1-r^2)^{\frac{n}{2}-1} dr &= \frac{1}{n} \int_0^1 \frac{(1-r^2)^{\frac{n}{2}}}{1+r} dr \\ &\leq \frac{1}{2n} \int_0^1 (1-r^2)^{\frac{n}{2}} dr \\ &= \frac{1}{2n} \int_0^{\frac{\pi}{2}} (\sin t)^n dt \leq \frac{C}{n^{\frac{3}{2}}}. \end{aligned} \quad \square$$

Let $\mathcal{B}(\mathbb{D})$ be the Bloch space of $\mathbb{D}$ and if $F \in \mathcal{B}(\mathbb{D})$, we extend F to the ball $\mathbb{B}_2$ by setting $F(z, w) := F(z)$. Then we have

Lemma 2.2. *If $F \in \mathcal{B}(\mathbb{D})$, then the big Hankel operator $H_{\overline{F}}$ is bounded on the Hardy space $H^2(\mathbb{B}_2)$.*

We write each $g \in H^2(\mathbb{B})$ in the form $g(z, w) = \sum_n g_n(z)w^n$, where the functions g_n are holomorphic in $\mathbb{D}$. In addition, we have the identity

$$\int_{\mathbb{S}} g_n(z)w^n \overline{g_m(z)w^m} d\sigma(z, w) = \begin{cases} 0, & n \neq m, \\ \int_{\mathbb{D}} |g_n(z)|^2 (1-|z|^2)^n dA(z), & n = m. \end{cases}$$

Hence

$$\|g\|_{L^2(\mathbb{S})}^2 = \sum_n \int_{\mathbb{D}} |g_n(z)|^2 (1-|z|^2)^n dA(z).$$

On the other hand, for each n , we have that

$$\begin{aligned} (P(\overline{F}g_n))(z, w) &= \int_{\mathbb{S}} \frac{\overline{F}(\zeta)g_n(\zeta)\eta^n}{(1-\bar{\zeta}z-\bar{\eta}w)^2} d\sigma(\zeta, \eta) \\ &= \int_{\mathbb{D}} \frac{\overline{F}(\zeta)g_n(\zeta)}{(1-\bar{\zeta}z-\bar{\eta}w)^2} \int_0^{2\pi} \frac{(1-|\zeta|^2)^{\frac{n}{2}} e^{in\theta}}{(1-\bar{\zeta}z-\sqrt{1-|\zeta|^2}we^{i\theta})^2} \frac{d\theta}{2\pi} dA(\zeta) \\ &= \int_{\mathbb{D}} \frac{\overline{F}(\zeta)g_n(\zeta)(1-|\zeta|^2)^{\frac{n}{2}}}{(1-\bar{\zeta}z)^2} \int_0^{2\pi} \frac{e^{in\theta}}{\left(1-\frac{\sqrt{1-|\zeta|^2}w}{1-\zeta z}e^{-i\theta}\right)^2} \frac{d\theta}{2\pi} dA(\zeta) \\ &= (n+1)w^n \int_{\mathbb{D}} \frac{\overline{F}(\zeta)g_n(\zeta)(1-|\zeta|^2)^n}{(1-\bar{\zeta}z)^{n+2}} dA(\zeta)w^n. \end{aligned}$$

Since $(n+1)\frac{(1-|\zeta|^2)^n}{(1-\bar{\zeta}z)^{n+2}}$ reproduces holomorphic functions on the unit disc, it follows that

$$\begin{aligned} (H_{\overline{F}}g)(z, w) &= \sum_n (n+1)w^n \int_{\mathbb{D}} \frac{\overline{F}(z)-\overline{F}(\zeta)}{(1-\bar{\zeta}z)^{n+2}} g_n(\zeta)(1-|\zeta|^2)^n dA(\zeta) \\ &= \sum_n H_n(z)w^n, \end{aligned}$$

where

$$H_n(z) := (n+1) \int_{\mathbb{D}} \frac{\overline{F}(z)-\overline{F}(\zeta)}{(1-\bar{\zeta}z)^{n+2}} g_n(\zeta)(1-|\zeta|^2)^n dA(\zeta).$$

Since $F \in \mathcal{B}(\mathbb{D})$, we see that $|F(z)-F(\zeta)| \leq \|F\|_{\mathcal{B}(\mathbb{D})}\beta(z, \zeta)$. Hence

$$|H_n(z)| \leq (n+1)\|F\|_{\mathcal{B}(\mathbb{D})} \int_{\mathbb{D}} \frac{\beta(z, \zeta)|g_n(\zeta)|}{|1-\bar{\zeta}z|^{n+2}} (1-|\zeta|^2)^n dA(\zeta)$$

so that by Hölder's inequality and Lemma 2.1 we see that

$$\begin{aligned} |H_n(z)|^2 &\leq (n+1)^2 \|F\|_{\mathcal{B}(\mathbb{D})}^2 (I_n(z))^2 \int_{\mathbb{D}} \frac{\beta(z, \zeta) |g_n(\zeta)|^2 (1 - |\zeta|^2)^{\frac{3n}{2}}}{|1 - \bar{\zeta}z|^{n+2}} dA(\zeta) \\ &\leq C\sqrt{n} \|F\|_{\mathcal{B}(\mathbb{D})}^2 (1 - |z|^2)^{-\frac{n}{2}} \int_{\mathbb{D}} \frac{\beta(z, \zeta) |g_n(\zeta)|^2 (1 - |\zeta|^2)^{\frac{n}{2}}}{|1 - \bar{\zeta}z|^{n+2}} (1 - |\zeta|^2)^n dA(\zeta). \end{aligned}$$

Therefore, again applying Lemma 2.1, we have

$$\begin{aligned} &\int_{\mathbb{D}} |H_n(z)|^2 (1 - |z|^2)^n dA(\zeta) \\ &\leq 2\sqrt{n} \|F\|_{\mathcal{B}(\mathbb{D})}^2 \int_{\mathbb{D}} |g_n(\zeta)|^2 (1 - |\zeta|^2)^{\frac{3n}{2}} \int_{\mathbb{D}} \frac{\beta(z, \zeta) (1 - |z|^2)^{\frac{n}{2}}}{|1 - \bar{\zeta}z|^{n+2}} dA(z) dA(\zeta) \\ &\leq 2 \frac{\|F\|_{\mathcal{B}(\mathbb{D})}^2}{n} \int_{\mathbb{D}} |g_n(\zeta)|^2 (1 - |\zeta|^2)^n dA(\zeta). \end{aligned}$$

Thus

$$\begin{aligned} \|(H_{\bar{F}}g)\|_{H^2(\mathbb{B})}^2 &= \sum_n \int_{\mathbb{D}} |H_n(z)|^2 (1 - |z|^2)^n dA(\zeta) \\ &\leq \sum_n \frac{\|F\|_{\mathcal{B}(\mathbb{D})}^2}{n} \int_{\mathbb{D}} |g_n(\zeta)|^2 (1 - |\zeta|^2)^n dA(\zeta) \\ &\leq \|F\|_{\mathcal{B}(\mathbb{D})}^2 \|g\|_{H^2(\mathbb{B})}^2. \end{aligned}$$

Proof of Theorem A in the ball case $\Omega_2 = \mathbb{B}_2$. Let $F \in \mathcal{B}(\mathbb{D}) \setminus BMOA(\mathbb{T})$ be extended to $\mathbb{B}_2$ as the function $(z, w) \mapsto F(z)$. By Lemma 2.2 the big Hankel operator $H_{\bar{F}}$ is bounded on the Hardy space $H^2(\mathbb{B}_2)$. Assume that there is a holomorphic function g on $\mathbb{B}_2$ such that the function $G := \bar{F} + g$ is bounded. Then the function $z \mapsto G(z, 0)$ is bounded on the unit disc and hence in $BMO(\mathbb{T})$. This shows that both functions F and $z \mapsto g(z, 0)$ are in $BMOA(\mathbb{T})$. This contradicts the fact that $F \notin BMOA(\mathbb{T})$. $\square$

3. THE BIDISC CASE

Let $bmo(\mathbb{T}^2)$ be the subspace of functions $f \in L^1(\mathbb{T}^2)$ such that f is in $BMO(\mathbb{T})$ in each variable and

$$\sup_{z \in \mathbb{T}} \|f(z, \cdot)\|_{BMO} + \sup_{w \in \mathbb{T}} \|f(\cdot, w)\|_{BMO} < +\infty.$$

In particular, if $f \in L^1(\mathbb{T}^2)$, then f is in $bmo(\mathbb{T}^2)$ if and only if

$$\sup_{z \in \mathbb{T}} \|Mf(z, \cdot)\|_{\infty} + \sup_{w \in \mathbb{T}} \|Mf(\cdot, w)\|_{\infty} < +\infty,$$

where

$$(M\varphi)(\zeta) := \sqrt{|\varphi|^2(\zeta) - |\tilde{\varphi}|^2(\zeta)}, \quad \zeta \in \mathbb{D}, \varphi \in L^2(\mathbb{T}).$$

The subspace of $bmo(\mathbb{T}^2)$ consisting of holomorphic functions on $\mathbb{D}^2$ will be denoted by $bmoa(\mathbb{T}^2)$.

Theorem 3.1. *If $\varphi \in L^\infty(\mathbb{D})$, then*

$$(T\varphi)(z, w) := \int_{\mathbb{D}} \frac{\varphi(\zeta)}{(1 - z\bar{\zeta})(1 - w\bar{\zeta})} dA(\zeta), \quad (z, w) \in \mathbb{T}^2,$$

is in $bmoa(\mathbb{T}^2)$. In addition, the operator T is bounded from $L^\infty(\mathbb{D})$ into $bmoa(\mathbb{T}^2)$.

To prove this theorem we need some auxiliary results. The following lemma can be found in [J].

Lemma 3.2. *There is a positive constant C such that*

$$\int_{\mathbb{D}} \frac{dA(\zeta)d}{|1 - z\bar{\zeta}|^\alpha |1 - w\bar{\zeta}|^\beta} \leq \begin{cases} C, & \alpha + \beta < 2, \\ C \log \frac{1}{|1 - z\bar{w}|}, & \alpha + \beta = 2, \\ \frac{C}{|1 - z\bar{w}|^{\alpha + \beta - 2}}, & \alpha + \beta > 2, \end{cases}$$

for all $z, w \in \mathbb{D}$.

Lemma 3.3. *There is a positive constant C such that*

$$I(z, w) := \int_{\mathbb{D} \times \mathbb{D}} \frac{dA(\zeta)dA(\eta)}{|1 - z\bar{\zeta}||1 - z\bar{\eta}||1 - w\bar{\zeta}||1 - w\bar{\eta}||1 - \zeta\bar{\eta}|} \leq \frac{C}{|1 - z\bar{w}|}$$

for all $z, w \in \mathbb{D}$.

Proof. We observe that $I(z, w) \leq I_1(z, w) + I_2(z, w) + I_3(z, w)$, where

$$I_j(z, w) := \int_{\mathbb{E}_j} \frac{dA(\zeta)dA(\eta)}{|1 - z\bar{\zeta}||1 - z\bar{\eta}||1 - w\bar{\zeta}||1 - w\bar{\eta}||1 - \zeta\bar{\eta}|}, \quad j = 1, 2, 3,$$

$$\mathbb{E}_1 := \{(\zeta, \eta) \in \mathbb{D} \times \mathbb{D} : |1 - \zeta\bar{\eta}| \leq \frac{1}{2}|1 - z\bar{\zeta}|\},$$

$$\mathbb{E}_2 := \{(\zeta, \eta) \in \mathbb{D} \times \mathbb{D} : |1 - \zeta\bar{\eta}| \leq \frac{1}{2}|1 - w\bar{\eta}|\}$$

and $\mathbb{E}_3$ is the complement of $\mathbb{E}_1 \cup \mathbb{E}_2$ in $\mathbb{D} \times \mathbb{D}$. By symmetry the integrals $I_1(z, w)$ and $I_2(z, w)$ can be estimated similarly. Since by Lemma 3.2

$$\begin{aligned} I_3(z, w) &\leq 2 \int_{\mathbb{D} \times \mathbb{D}} \frac{dA(\zeta)dA(\eta)}{|1 - z\bar{\zeta}|^{\frac{3}{2}}|1 - z\bar{\eta}||1 - w\bar{\zeta}||1 - w\bar{\eta}|^{\frac{3}{2}}} \\ &\leq \frac{C}{|1 - z\bar{w}|}, \end{aligned}$$

we only need to estimate $I_1(z, w)$. To do so, we write $I_1(z, w) = I_{11}(z, w) + I_{12}(z, w)$, where

$$I_{1j}(z, w) := \int_{\mathbb{E}_{1j}} \frac{dA(\zeta)dA(\eta)}{|1 - z\bar{\zeta}||1 - z\bar{\eta}||1 - w\bar{\zeta}||1 - w\bar{\eta}||1 - \zeta\bar{\eta}|}, \quad j = 1, 2,$$

$\mathbb{E}_{11} := \{(\zeta, \eta) \in \mathbb{E}_1 : |1 - \zeta\bar{\eta}| \leq \frac{1}{2}|1 - w\bar{\zeta}|\}$ and $\mathbb{E}_{12}$ is the complement of $\mathbb{E}_{11}$ in $\mathbb{E}_1$. Set $R(z, w, \zeta) := \min(|1 - z\bar{\zeta}|, |1 - w\bar{\zeta}|)$. Since on $\mathbb{E}_{11}$ we have $|1 - z\bar{\zeta}| \approx |1 - z\bar{\eta}|$ and $|1 - w\bar{\zeta}| \approx |1 - w\bar{\eta}|$, it follows that

$$\begin{aligned} I_{11}(z, w) &\approx \int_{\mathbb{E}_{11}} \frac{dA(\zeta)dA(\eta)}{|1 - z\bar{\zeta}|^2|1 - w\bar{\zeta}|^2|1 - \zeta\bar{\eta}|} \\ &\leq \int_{\mathbb{D}} \int_{\{\eta \in \mathbb{D} : |1 - \zeta\bar{\eta}| \leq \frac{1}{2}R(z, w, \zeta)\}} \frac{dA(\eta)}{|1 - \zeta\bar{\eta}|} \frac{dA(\zeta)}{|1 - z\bar{\zeta}|^2|1 - w\bar{\zeta}|^2} \\ &\leq \int_{\mathbb{D}} \frac{R(z, w, \zeta)}{|1 - z\bar{\zeta}|^2|1 - w\bar{\zeta}|^2} dA(\zeta) \\ &\leq \frac{C}{|1 - z\bar{w}|}, \end{aligned}$$

where the latter inequality holds in view of Lemma 3.2. It now remains to establish the estimate of $I_{12}(z, w)$. We write $I_{12}(z, w) = \sum_{j=1}^3 I_{12j}(z, w)$, where

$$\begin{aligned} I_{12j}(z, w) &:= \int_{\mathbb{E}_{12j}} \frac{dA(\zeta)dA(\eta)}{|1 - z\bar{\zeta}||1 - z\bar{\eta}||1 - w\bar{\zeta}||1 - w\bar{\eta}||1 - \zeta\bar{\eta}|}, \quad j = 1, 2, 3, \\ \mathbb{E}_{121} &:= \{(\zeta, \eta) \in \mathbb{E}_{12} : |1 - w\bar{\eta}| \leq \frac{1}{2}|1 - \zeta\bar{\eta}|\}, \\ \mathbb{E}_{122} &:= \{(\zeta, \eta) \in \mathbb{E}_{12} : |1 - \zeta\bar{\eta}| \leq \frac{1}{2}|1 - w\bar{\eta}|\}, \end{aligned}$$

and $\mathbb{E}_{123}$ is the complement of $\mathbb{E}_{121} \cup \mathbb{E}_{122}$ in $\mathbb{E}_{12}$. On $\mathbb{E}_{121}$ we have $|1 - w\bar{\zeta}| \approx |1 - \zeta\bar{\eta}|$ and $|1 - w\bar{\eta}| \leq CR(z, w, \zeta)$. Hence, by Lemma 3.2,

$$\begin{aligned} I_{121}(z, w) &\approx \int_{\mathbb{D}} \int_{\{\eta \in \mathbb{D} : |1 - w\bar{\eta}| \leq CR(z, w, \zeta)\}} \frac{A(\eta)}{|1 - w\bar{\eta}|} \frac{dA(\zeta)}{|1 - z\bar{\zeta}|^2 |1 - w\bar{\zeta}|^2} \\ &\leq C \int_{\mathbb{D}} \frac{R(z, w, \zeta)}{|1 - z\bar{\zeta}|^2 |1 - w\bar{\zeta}|^2} dA(\zeta) \\ &\leq \frac{C}{|1 - z\bar{w}|}. \end{aligned}$$

On $\mathbb{E}_{122}$ we have $|1 - w\bar{\zeta}| \approx |1 - w\bar{\eta}|$ and $|1 - \zeta\bar{\eta}| \leq CR(z, w, \zeta)$. Hence as in the previous case we have

$$\begin{aligned} I_{122}(z, w) &\approx \int_{\mathbb{D}} \int_{\{\eta \in \mathbb{D} : |1 - w\bar{\eta}| \leq CR(z, w, \zeta)\}} \frac{A(\eta)}{|1 - w\bar{\eta}|} \frac{dA(\zeta)}{|1 - z\bar{\zeta}|^2 |1 - w\bar{\zeta}|^2} \\ &\leq \frac{C}{|1 - z\bar{w}|}. \end{aligned}$$

Finally, on $\mathbb{E}_{123}$ we have $|1 - w\bar{\eta}| \approx |1 - \zeta\bar{\eta}|$ and $|1 - w\bar{\zeta}| \leq 2|1 - \zeta\bar{\eta}|$. Thus

$$\begin{aligned} I_{123}(z, w) &\approx \int_{\mathbb{D}} \int_{\{\eta \in \mathbb{D} : |1 - \zeta\bar{\eta}| \leq \frac{1}{2}|1 - z\bar{\zeta}|\}} \frac{A(\eta)}{|1 - \zeta\bar{\eta}|^{\frac{3}{2}}} \frac{dA(\zeta)}{|1 - z\bar{\zeta}|^2 |1 - w\bar{\zeta}|^{\frac{3}{2}}} \\ &\leq C \int_{\mathbb{D}} \frac{dA(\zeta)}{|1 - z\bar{\zeta}|^{\frac{3}{2}} |1 - w\bar{\zeta}|^{\frac{3}{2}}} \\ &\leq \frac{C}{|1 - z\bar{w}|}. \end{aligned}$$

□

Proof of Theorem 3.1. Let

$$f(z, w) = (T\varphi)(z, w) = \int_{\mathbb{D}} \frac{\varphi(\zeta)}{(1 - z\bar{\zeta})(1 - w\bar{\zeta})} dA(\zeta), \quad (z, w) \in \mathbb{D}^2.$$

Then f is in the Hardy space $H^2(\mathbb{D}^2)$ and satisfies $\widetilde{f(\cdot, w)}(z) = f(z, w)$ and

$$\begin{aligned} |\widetilde{f(\cdot, w)}|^2(z) &= \frac{1}{2i\pi} \int_{\mathbb{T}} \int_{\mathbb{D} \times \mathbb{D}} \frac{\varphi(\zeta)\overline{\varphi(\eta)}(1 - |z|^2)dA(\zeta)dA(\eta)}{(1 - \tau\bar{\zeta})(1 - w\bar{\zeta})(1 - \bar{\tau}\eta)(1 - \bar{w}\eta)|1 - z\bar{\tau}|^2} \frac{d\tau}{\tau} \\ &= \int_{\mathbb{D} \times \mathbb{D}} \frac{\varphi(\zeta)\overline{\varphi(\eta)}(1 - \eta\bar{\zeta}|z|^2)dA(\zeta)dA(\eta)}{(1 - z\bar{\zeta})(1 - w\bar{\zeta})(1 - \bar{z}\eta)(1 - \bar{w}\eta)(1 - \zeta\bar{\eta})} \end{aligned}$$

by the Cauchy formula. Therefore, by Lemma 3.3 we see that

$$\begin{aligned} (Mf(\cdot, w))^2(z) &= |\widetilde{f(\cdot, w)}|^2(z) - |\widetilde{f}(z, w)|^2 \\ &= \left| \int_{\mathbb{D} \times \mathbb{D}} \frac{\zeta \bar{\eta} \varphi(\zeta) \overline{\varphi(\eta)} (1 - |z|^2) dA(\zeta) dA(\eta)}{(1 - z\bar{\zeta})(1 - w\bar{\zeta})(1 - \bar{z}\eta)(1 - \bar{w}\eta)(1 - \zeta\bar{\eta})} \right| \\ &\leq \|\varphi\|_\infty^2 (1 - |z|^2) \int_{\mathbb{D} \times \mathbb{D}} \frac{dA(\zeta) dA(\eta)}{|(1 - z\bar{\zeta})(1 - w\bar{\zeta})(1 - \bar{z}\eta)(1 - \bar{w}\eta)(1 - \zeta\bar{\eta})|} \\ &\leq C \|\varphi\|_\infty^2. \end{aligned}$$

Similarly we have $\|Mf(z, \cdot)\|_\infty \leq C \|\varphi\|_\infty$, showing that $T\varphi \in bmoa(\mathbb{T}^2)$ and

$$\|T\varphi\|_{bmoa} \leq C \|\varphi\|_\infty.$$

□

Theorem 3.4. Suppose that $h \in \mathcal{B}(\mathbb{D})$ and $h \notin BMOA(\mathbb{T})$ and let

$$f(z, w) := \int_{\mathbb{D}} \frac{h(\zeta)}{(1 - z\bar{\zeta})(1 - w\bar{\zeta})} dA(\zeta), \quad (z, w) \in \mathbb{T}^2.$$

Then the big Hankel operator $H_{\bar{f}}$ is bounded on $H^2(\mathbb{D}^2)$, and for any holomorphic function g on $\mathbb{D} \times \mathbb{D}$, the function $\bar{f} + g$ cannot be bounded.

Proof. Since $h \in \mathcal{B}(\mathbb{D})$, there is $\varphi \in L^\infty(\mathbb{D})$ such that

$$h(\lambda) = \int_{\mathbb{D}} \frac{\varphi(\zeta)}{(1 - \lambda\bar{\zeta})^2} dA(\zeta), \quad \lambda \in \mathbb{D}.$$

By the reproducing formula of the Bergman kernel and Theorem 3.1 we see that

$$f(z, w) := \int_{\mathbb{D}} \frac{\varphi(\zeta)}{(1 - z\bar{\zeta})(1 - w\bar{\zeta})} dA(\zeta) = (T\varphi)(z, w) \in bmoa(\mathbb{T}^2).$$

By [FS] we see that $H_{\bar{f}}$ is bounded on $H^2(\mathbb{D}^2)$. Now suppose that there is a holomorphic function g on $\mathbb{D}^2$ such that the function $F := \bar{f} + g$ is bounded. Then the restriction of F to the diagonal is also bounded and hence in $BMO(\mathbb{T})$. Since if $\lambda \in \mathbb{D}$, we have $F(\lambda, \lambda) = h(\lambda) + g(\lambda, \lambda)$, it follows that both functions h and $g(\lambda, \lambda)$ are in $BMOA(\mathbb{T})$. This contradicts the fact that $h \notin BMOA(\mathbb{T})$. □

Now, Theorem A in the bidisc case follows from Theorem 3.4.

Proposition 3.5. $bmo(\mathbb{T}^2) = \widetilde{BMO}(\mathbb{T}^2)$.

Proof. For $z, w \in \mathbb{D}$ and $\theta \in [0, 2\pi]$, set $P_z(\theta) := \frac{1 - |z|^2}{1 - ze^{-i\theta}\bar{z}}$ and define the measure $\mu_{z,w}$ on $[0, 2\pi] \times [0, 2\pi] \times [0, 2\pi] \times [0, 2\pi]$ by

$$d\mu_{z,w}(\theta, s, \psi, t) := \frac{1}{(2\pi)^4} P_z(\theta) P_z(\psi) P_w(s) P_w(t) d\theta ds d\psi dt.$$

A little computing shows that if $f \in L^2(\mathbb{T}^2)$, then we have the following key equality:

$$\begin{aligned} 2 \left(|\widetilde{f}|^2 - |\widetilde{f}|^2 \right) (z, w) \\ = \int_0^{2\pi} \int_0^{2\pi} \int_0^{2\pi} \int_0^{2\pi} |f(e^{i\theta}, e^{is}) - f(e^{i\psi}, e^{it})|^2 d\mu_{z,w}(\theta, s, \psi, t), \end{aligned}$$

which will be used to prove both inclusions. On the one hand, suppose that $f \in bmo(\mathbb{T}^2)$. Then the key inequality and the triangle inequality imply that

$$\sqrt{2 \left(\widetilde{|f|^2} - |\tilde{f}|^2 \right)}(z, w) \leq \sqrt{A(z, w)} + \sqrt{B(z, w)},$$

where

$$\begin{aligned} A(z, w) &:= \int_0^{2\pi} \int_0^{2\pi} \int_0^{2\pi} \int_0^{2\pi} |f(e^{i\theta}, e^{is}) - f(e^{i\theta}, e^{it})|^2 d\mu_{z,w}(\theta, s, \psi, t) \\ &= \frac{1}{(2\pi)^3} \int_0^{2\pi} \int_0^{2\pi} \int_0^{2\pi} |f(e^{i\theta}, e^{is}) - f(e^{i\theta}, e^{it})|^2 P_z(\theta) P_w(s) P_w(t) d\theta ds dt \\ &= \frac{1}{\pi} \int_0^{2\pi} (M[f(e^{i\theta}, \cdot)](w))^2 P_z(\theta) d\theta \end{aligned}$$

and similarly

$$\begin{aligned} B(z, w) &:= \int_0^{2\pi} \int_0^{2\pi} \int_0^{2\pi} \int_0^{2\pi} |f(e^{i\theta}, e^{it}) - f(e^{i\psi}, e^{it})|^2 d\mu_{z,w}(\theta, s, \psi, t) \\ &= \frac{1}{\pi} \int_0^{2\pi} (M[f(\cdot, e^{it})](z))^2 P_w(t) dt \end{aligned}$$

showing that $bmo(\mathbb{T}^2) \subseteq \widetilde{BMO}(\mathbb{T}^2)$.

On the other hand, suppose that $f \in \widetilde{BMO}(\mathbb{T}^2)$. By Proposition 1.3 of [CS], in order to show that $f \in bmo(\mathbb{T}^2)$, we only need prove that f is of bounded mean oscillation on rectangles; that is, for some constant $C > 0$ we have

$$\frac{1}{|I|} \frac{1}{|J|} \int_{I \times J} |f(e^{i\theta}, e^{is}) - f_{I \times J}|^2 d\theta ds \leq C$$

for all intervals $I, J \subset [0, 2\pi]$, where $|I|$ is the length of an interval I and

$$f_{I \times J} := \frac{1}{|I||J|} \int_I \int_J f(e^{i\theta}, e^{is}) d\theta ds.$$

To do so, let $z = \sqrt{1 - \varepsilon} e^{i\theta_0}$ and $w = \sqrt{1 - \varepsilon} e^{is_0}$ where $\theta_0, s_0 \in [0, 2\pi]$ and $\varepsilon \in]0, 1]$, then for some constant $C > 0$ independent of all parameters we have

$$P_z(\theta) \geq \frac{C}{1 - |z|^2} \quad \text{and} \quad P_w(\theta) \geq \frac{C}{1 - |w|^2}.$$

Thus by the key equality above we have

$$\begin{aligned} &2 \left(\widetilde{|f|^2} - |\tilde{f}|^2 \right)(z, w) \\ &\geq \int_{\theta_0 - \varepsilon}^{\theta_0 + \varepsilon} \int_{s_0 - \varepsilon}^{s_0 + \varepsilon} \int_{\theta_0 - \varepsilon}^{\theta_0 + \varepsilon} \int_{s_0 - \varepsilon}^{s_0 + \varepsilon} |f(e^{i\theta}, e^{is}) - f(e^{i\psi}, e^{it})|^2 d\mu_{z,w}(\theta, s, \psi, t) \\ &\geq \frac{C^4}{(2\varepsilon\pi)^4} \int_{\theta_0 - \varepsilon}^{\theta_0 + \varepsilon} \int_{s_0 - \varepsilon}^{s_0 + \varepsilon} \int_{\theta_0 - \varepsilon}^{\theta_0 + \varepsilon} \int_{s_0 - \varepsilon}^{s_0 + \varepsilon} |f(e^{i\theta}, e^{is}) - f(e^{i\psi}, e^{it})|^2 dt d\psi ds d\theta \\ &= \frac{2C^4}{(2\pi)^4} \frac{1}{(2\varepsilon)^2} \int_{\theta_0 - \varepsilon}^{\theta_0 + \varepsilon} \int_{s_0 - \varepsilon}^{s_0 + \varepsilon} |f(e^{i\theta}, e^{is}) - f_{] \theta_0 - \varepsilon, \theta_0 + \varepsilon[\times] s_0 - \varepsilon, s_0 + \varepsilon[}|^2 ds d\theta. \end{aligned}$$

This shows that

$$\frac{1}{|I|} \frac{1}{|J|} \int_{I \times J} |f(e^{i\theta}, e^{is}) - f_{I \times J}|^2 d\theta ds \leq \frac{(2\pi)^4}{C^4} \sup_{z, w \in \mathbb{D}} \left(\widetilde{|f|^2} - |\tilde{f}|^2 \right) (z, w)$$

for all intervals I, J with length less than or equal to 2. Similar argument shows that

$$\sup_{|I| \leq 2, |J| > 2} \frac{1}{|I|} \frac{1}{|J|} \int_{I \times J} |f(e^{i\theta}, e^{is}) - f_{I \times J}|^2 d\theta ds \leq \frac{(2\pi)^2}{C^2} \sup_{z \in \mathbb{D}} \left(\widetilde{|f|^2} - |\tilde{f}|^2 \right) (z, 0),$$

and by symmetry we have

$$\sup_{|J| \leq 2, |I| > 2} \frac{1}{|I|} \frac{1}{|J|} \int_{I \times J} |f(e^{i\theta}, e^{is}) - f_{I \times J}|^2 d\theta ds \leq \frac{(2\pi)^2}{C^2} \sup_{w \in \mathbb{D}} \left(\widetilde{|f|^2} - |\tilde{f}|^2 \right) (0, w).$$

The case where both intervals have length greater than 2 can be handled easily. $\square$

Proposition 3.5, combined with Theorem 2.1 of [FS], proves Theorem B and answers in the affirmative a conjecture due to Berger, Coburn and Zhu [C] for the bidisc case.

REFERENCES

- [BT] M. Bakonyi and D. Timotin, *On a conjecture of Coltar and Sadosky on multidimensional Hankel operators*, C. R. Acad. Sci. Paris **t. 325, Série I** (1997), 1071-1075. MR1614007 (99b:47032)
- [C] L. Coburn, *Toeplitz operators, quantum mechanics and mean oscillation in the Bergman metric*, Proceedings of Symposia in Pure Mathematics **51** (1990), 97-104. MR1077380 (92j:47045)
- [CS] M. Cotlar and C. Sadosky, *Two distinguished subspaces of product BMO and Nehari-AAK theory of Hankel operators on the torus*, Integral Equations and Operator Theory **26** (1996), 273-304. MR1415032 (98e:47043)
- [FS] S. Ferguson and C. Sadosky, *Characterizations of bounded mean oscillation on the polydisc in terms of Hankel operators and Carleson measures*, J. Anal. Math. **81** (2000), 239-267. MR1785283 (2001h:47040)
- [J] S. Jacobson, *The harmonic Bergman kernel and the Friedrichs operator*, Ark. Mat. **40** (2002), 89-104. MR1948888 (2003j:30019)
- [S] E. Strouse, *Conference in Holomorphic Function Spaces and their Operators*, CIRM Marseille, 2002.
- [Z1] K. Zhu, *Operator Theory in Function Spaces*, Marcel Dekker, New York, 1990. MR1074007 (92c:47031)
- [Z2] K. Zhu, *Spaces of holomorphic functions in the unit ball*, Graduate Texts in Math., 226, Springer, New York, 2005. MR2115155 (2006d:46035)

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WISCONSIN, 480 LINCOLN DRIVE, MADISON, WISCONSIN 53705

E-mail address: `ahern@math.wisc.edu`

LATP, U.M.R. C.N.R.S. 6632, CMI, UNIVERSITÉ DE PROVENCE, 39 RUE F-JOLIOT-CURIE, 13453 MARSEILLE CEDEX 13, FRANCE

E-mail address: `youssfi@gyptis.univ-mrs.fr`