RANK OF THE FUNDAMENTAL GROUP OF ANY COMPONENT OF A FUNCTION SPACE

GREGORY LUPTON AND SAMUEL BRUCE SMITH

(Communicated by Paul Goerss)

ABSTRACT. We compute the rank of the fundamental group of any connected component of the space $\text{map}(X, Y)$ for X and Y connected, nilpotent CW complexes of finite type with X finite. For the component corresponding to a general homotopy class $f: X \rightarrow Y$, we give a formula directly computable from the Sullivan model for f . For the component of the constant map, our formula retrieves a known expression for the rank in terms of classical invariants of X and Y . When both X and Y are rationally elliptic spaces with positive Euler characteristic, we use our formula to determine the rank of the fundamental group of any component of $\text{map}(X, Y)$ explicitly in terms of the homomorphism induced by f on rational cohomology.

1. INTRODUCTION

The *rank* of a finitely generated abelian group G is the cardinality of a basis of the free abelian group G/T , where T is the torsion subgroup of G . This notion can be extended to finitely generated nilpotent groups Γ by means of the central series $\Gamma = \Gamma_0 \supset \Gamma_1 \supset \cdots \supset \Gamma_n = \{1\}$, where $\Gamma_n = [\Gamma_{n-1}, \Gamma]$. The *rank* of Γ is the sum of the ranks of the finitely generated abelian quotients Γ_{k-1}/Γ_k for $k = 1, \dots, n$.

Given connected spaces X and Y , let $\text{map}(X, Y)$ denote the function space of all (not-necessarily based) maps from X to Y with the compact-open topology. Given a map $f: X \rightarrow Y$, write $\text{map}(X, Y; f)$ for the connected component of $\text{map}(X, Y)$ containing f , that is, the space of maps freely homotopic to f . Under reasonable hypotheses on X and Y , the fundamental groups of the connected components of $\text{map}(X, Y)$ are finitely generated nilpotent groups. To be precise, say a space X is *nilpotent* if $\pi_1(X)$ is a nilpotent group and the action of $\pi_1(X)$ on the higher homotopy groups of X is a nilpotent action. If X is a finite CW complex and Y is a nilpotent CW complex of finite type, then the components of $\text{map}(X, Y)$ are themselves nilpotent CW complexes of finite type ([11] and [9, Th.II.2.5]). Thus with these hypotheses $\pi_1(\text{map}(X, Y; f))$ is a finitely generated nilpotent group. Our purpose in this paper is to give a formula for the rank of this group expressed in terms of accessible invariants of the map $f: X \rightarrow Y$.

In [10], we have described the higher rational homotopy groups of $\text{map}(X, Y; f)$. Our description there is in terms of the homology of chain complexes of derivations

Received by the editors February 1, 2006 and, in revised form, April 13, 2006.

2000 *Mathematics Subject Classification.* Primary 55Q52, 55P62.

Key words and phrases. Function space, fundamental group, nilpotent space, nilpotent group, rank, rational homotopy, minimal models, derivations.

©2007 American Mathematical Society
 Reverts to public domain 28 years from publication

of Sullivan minimal models. Specifically, for each $n \geq 2$ we constructed a vector space isomorphism

$$(1) \quad \pi_n(\text{map}(X, Y; f)) \otimes \mathbb{Q} \cong H_n(\text{Der}(\mathcal{M}_Y, \mathcal{M}_X; \mathcal{M}_f)),$$

where $\text{Der}_*(\mathcal{M}_Y, \mathcal{M}_X; \mathcal{M}_f)$ is a graded vector space of (generalized) algebra derivations constructed directly from a Sullivan model $\mathcal{M}_f: \mathcal{M}_Y \rightarrow \mathcal{M}_X$ of the map $f: X \rightarrow Y$ (see Section 2 below for precise definitions). Now within the framework established there, we may also compute $H_1(\text{Der}(\mathcal{M}_Y, \mathcal{M}_X; \mathcal{M}_f))$, which is a rational homotopy invariant of the map f . But the fundamental group $\pi_1(\text{map}(X, Y; f))$ is generally non-abelian, even after rationalization—see Example 1.1 below. So it is evidently not possible to identify the rationalized fundamental group, *as a group*, with the (abelian) vector space $H_1(\text{Der}(\mathcal{M}_Y, \mathcal{M}_X; \mathcal{M}_f))$. In other words, the isomorphism (1) does not extend to the case $n = 1$. Nonetheless, one intuitively expects this latter vector space to be related to the rationalization of the fundamental group. Our main result of this paper makes their relationship explicit:

Theorem 1. *Let X and Y be connected, nilpotent CW complexes of finite type with X finite. Let $f: X \rightarrow Y$ be any given map. Then*

$$\text{rank}(\pi_1(\text{map}(X, Y; f))) = \dim_{\mathbb{Q}}(H_1(\text{Der}(\mathcal{M}_Y, \mathcal{M}_X; \mathcal{M}_f))).$$

By $\dim_{\mathbb{Q}}$ on the right-hand side of the equation we mean the dimension as a rational vector space.

We prove this result in Section 4. As an illustration of the possible applications, in Theorem 4.1 we express $\text{rank}(\pi_1(\text{map}(X, Y; f)))$ for a map $f: X \rightarrow Y$ between certain formal spaces directly in terms of the map induced by f on rational cohomology. A particular consequence is Example 4.2: let G be a compact, simple Lie group of rank $n \geq 2$, $T^n \subseteq G$ a maximal torus and $f: S^2 \rightarrow G/T^n$ any map. Then $\pi_1(\text{map}(S^2, G/T^n; f))$ is finite—that is, has rank 0—if and only if f is essential.

Specializing Theorem 1 to the case in which $f = 0$, we obtain the following corollary. (This result may also be deduced from Corollary 1.6 of [3].) Let $b_n(X)$ denote the n th Betti number of X . If $n \geq 2$ let $\rho_n(Y)$ denote the integer $\dim_{\mathbb{Q}}(\pi_n(Y) \otimes \mathbb{Q})$ —sometimes called the n th *Hurewicz number* of Y .

Corollary 1. *Let X and Y be connected, nilpotent CW complexes of finite type with X a finite complex of dimension N . Then*

$$\text{rank}(\pi_1(\text{map}(X, Y; 0))) = \text{rank}(\pi_1(Y)) + \sum_{n=2}^{N+1} \rho_n(Y) \cdot b_{n-1}(X).$$

Our results here and in [10] complement and augment those of others. In addition to [7] and [3], we cite [14] in which the author obtains results for the null-component case. Recent results of Buijs and Murillo [4] overlap with, indeed extend some aspects of, our work. We will discuss our methods after a brief, motivating example.

Example 1.1. Let $X = \mathbb{C}P^2$ and let Y be the total space of the principal fibration $K(\mathbb{Z}, 5) \rightarrow Y \rightarrow K(\mathbb{Z} \times \mathbb{Z}, 3)$ whose k -invariant is $k: K(\mathbb{Z} \times \mathbb{Z}, 3) \rightarrow K(\mathbb{Z}, 6)$, corresponding to the cup-product $x \cup y \in H^6(K(\mathbb{Z} \times \mathbb{Z}, 3); \mathbb{Z})$. Here, x and y denote the generators in $H^3(K(\mathbb{Z} \times \mathbb{Z}, 3); \mathbb{Z}) \cong \mathbb{Z} \times \mathbb{Z}$. We now consider the null-component $\text{map}(X, Y; 0)$. The Sullivan minimal model of Y is $\mathcal{M}_Y = \Lambda(x_3, y_3, z_5; d)$ with the single non-trivial differential $d(z_5) = x_3 y_3$. Here a subscript on a generator denotes its degree. The rational homology coalgebra of X is the vector space $\langle 1, a_2, b_4 \rangle$,

with a_2 primitive and the coproduct on b_4 given by $\Delta(b_4) = b_4 \otimes 1 + 1 \otimes b_4 + a_2 \otimes a_2$. Because we are focussing on the null-component, and because X is a formal space, we may readily assemble the Sullivan minimal model of $\text{map}(X, Y; 0)$ from Theorems 1.9 and 6.1 of [3]. We may write it as $\Lambda(\alpha_1, \beta_1, \gamma_1, x_3, y_3, \eta_3, z_5; \delta)$ with non-zero differentials $\delta(\gamma_1) = \alpha_1\beta_1$, $\delta(\eta_3) = \alpha_1x_3 + \beta_1y_3$ and $\delta(z_5) = x_3y_3$. (In the notation of [3], our generators α_1 , β_1 and γ_1 , for instance, would be written as $x_3 \otimes a_2$, $y_3 \otimes a_2$ and $z_5 \otimes b_4$, respectively, and the formulas for the differentials were obtained via the comments preceding Theorem 1.9 of that paper.) Then the fact that there are 3 degree-one generators implies that $\text{rank}(\pi_1(\text{map}(X, Y; 0))) = 3$ and the non-zero differential $\delta(\gamma_1) = \alpha_1\beta_1$ implies that *the rationalized fundamental group is non-abelian*.

This example indicates that, even for relatively simple choices of X , Y and f , the rationalization of $\pi_1(\text{map}(X, Y; f))$ can display quite rich structure. Example 1.1 also illustrates the effectiveness of the models of [7] and [3] for concrete computations. As regards our methods here, we must emphasize an important point about these models, since our proof of Theorem 1 does not use them. These models are often quite effective tools for gleaning information about a null-component $\text{map}(X, Y; 0)$ (see, for example, [14]). But for $f: X \rightarrow Y$ an essential map, these models are generally quite complicated. By focussing purely on the rank of $\pi_1(\text{map}(X, Y; f))$, we are able to give a “closed form” description that proceeds directly from the ordinary Sullivan model of the map $f: X \rightarrow Y$ and is relatively easy to handle in practice.

2. DERIVATIONS AND FUNCTION SPACE COMPONENTS

We assume basic familiarity with localization of nilpotent groups and spaces, as discussed in [9]. Recall that a nilpotent space Y (respectively, group G) admits a *rationalization*, which we denote $e_Y: Y \rightarrow Y_{\mathbb{Q}}$ (respectively, $e_G: G \rightarrow G_{\mathbb{Q}}$). If G is abelian, then $G_{\mathbb{Q}} \cong G \otimes \mathbb{Q}$. If $f: X \rightarrow Y$ is a map, we write $f_{\mathbb{Q}}: X \rightarrow Y_{\mathbb{Q}}$ for the composition $e_Y \circ f$ and likewise for a group homomorphism. Post-composition with a map of spaces $g: Y \rightarrow Z$ induces a map $g_*: \text{map}(X, Y; f) \rightarrow \text{map}(X, Z; g \circ f)$. By [9, Th.II.3.1], if X and Y satisfy the hypotheses of Theorem 1, then $\text{map}(X, Y; f)$ is a nilpotent space and the induced map $(e_Y)_*: \text{map}(X, Y; f) \rightarrow \text{map}(X, Y_{\mathbb{Q}}; f_{\mathbb{Q}})$ is a rationalization. Under these hypotheses we have, for all $n \geq 1$,

$$(2) \quad \pi_n(\text{map}(X, Y; f))_{\mathbb{Q}} \cong \pi_n(\text{map}(X, Y_{\mathbb{Q}}; f_{\mathbb{Q}})).$$

We also assume familiarity with rational homotopy theory using the DG (differential graded) algebra minimal models introduced by Sullivan. The basic facts that we rely on are as follows: Each nilpotent space of finite-type Y has a Sullivan minimal model $(\mathcal{M}_Y, d_Y)$ in the category of nilpotent DG algebras over $\mathbb{Q}$. This DG algebra is unique up to isomorphism and is of the form $\mathcal{M}_Y = \Lambda W$, a free graded commutative algebra generated by a positively graded vector space W of finite type. A map $f: X \rightarrow Y$ of nilpotent, finite-type spaces also has a Sullivan minimal model, which is a DG algebra map $\mathcal{M}_f: \mathcal{M}_Y \rightarrow \mathcal{M}_X$. The Sullivan minimal model is a complete rational homotopy invariant for a space or a map. Our main reference for this material is [5], although that book restricts to the simply connected case. See [2] for a treatment of the non-simply-connected case.

Fix two DG algebras (A, d_A) and (B, d_B) and a DG algebra map $\phi: A \rightarrow B$. Define a ϕ -*derivation* to be a linear map $\theta: A \rightarrow B$ that *reduces degree by n* and

satisfies the derivation law $\theta(xy) = \theta(x)\phi(y) + (-1)^{|x|}\phi(x)\theta(y)$. Let $\text{Der}_n(A, B; \phi)$ denote the vector space of ϕ -derivations of degree $n \geq 0$. Define a linear map $D: \text{Der}_n(A, B; \phi) \rightarrow \text{Der}_{n-1}(A, B; \phi)$ by $D(\theta) = d_B \circ \theta - (-1)^{|\theta|}\theta \circ d_A$. A standard check now shows that $D^2 = 0$ and thus $(\text{Der}_*(A, B; \phi), D)$ is a chain complex. To ease notation, we will suppress the differential and only write the outermost degree in the homology of a DG space. In particular, $H_n(\text{Der}(A, B; \phi))$ will denote the homology in degree n of the chain complex $(\text{Der}_*(A, B; \phi), D)$.

In [10] we showed that the homology theory of these chain complexes may be used to model the rational homotopy theory of function spaces at the level of (higher) homotopy groups. The link between the two is provided by a map

$$(3) \quad \Phi_f: \pi_n(\text{map}(X, Y; f)) \rightarrow H_n(\text{Der}(\mathcal{M}_Y, \mathcal{M}_X; \mathcal{M}_f)),$$

where $\mathcal{M}_f: \mathcal{M}_Y \rightarrow \mathcal{M}_X$ is any choice of a Sullivan minimal model of f . The construction of this map originates with Thom [13] and is as follows. Suppose $\beta \in \pi_n(\text{map}(X, Y; f))$ has adjoint $B: S^n \times X \rightarrow Y$. Then B induces a DG algebra map $\mathcal{A}_B: \mathcal{M}_Y \rightarrow H^*(S^n; \mathbb{Q}) \otimes \mathcal{M}_X$ that is of the form

$$\mathcal{A}_B(\chi) = 1 \otimes \mathcal{M}_f(\chi) + u \otimes \theta_B(\chi),$$

where $u \in H^n(S^n; \mathbb{Q})$ denotes a generator and $\chi \in \mathcal{M}_Y$ is of positive degree. Here $H^*(S^n; \mathbb{Q})$ is viewed as a DG algebra with zero differential. This expression defines a linear map $\theta_B: \mathcal{M}_Y \rightarrow \mathcal{M}_X$ that reduces the degree by n . A standard check shows θ_B is an $\mathcal{M}_f$ -derivation and a cycle. Set $\Phi_f(\beta) = [\theta_B] \in H_n(\text{Der}(\mathcal{M}_Y, \mathcal{M}_X; \mathcal{M}_f))$.

Theorem 2.1. *Let $f: X \rightarrow Y$ be a map between connected, nilpotent CW complexes of finite type with X finite. Let $\mathcal{M}_f: \mathcal{M}_Y \rightarrow \mathcal{M}_X$ be any choice of a Sullivan minimal model of f . Then for $n \geq 1$ the map*

$$\Phi_f: \pi_n(\text{map}(X, Y; f)) \rightarrow H_n(\text{Der}(\mathcal{M}_Y, \mathcal{M}_X; \mathcal{M}_f))$$

is a well-defined map (of sets). Furthermore, we have:

- (a) *For $n \geq 2$, Φ_f is a group homomorphism whose rationalization*

$$\Phi_{f_{\mathbb{Q}}}: \pi_n(\text{map}(X, Y_{\mathbb{Q}}; f_{\mathbb{Q}})) \rightarrow H_n(\text{Der}(\mathcal{M}_Y, \mathcal{M}_X; \mathcal{M}_f))$$

is an isomorphism.

- (b) *Let Y be an H -space and suppose f is null-homotopic. For $n \geq 1$, Φ_f is a group homomorphism whose rationalization is an isomorphism.*

- (c) *For nilpotent Y of finite type, we have a well-defined surjection of sets*

$$\Phi_{f_{\mathbb{Q}}}: \pi_1(\text{map}(X, Y_{\mathbb{Q}}; f_{\mathbb{Q}})) \rightarrow H_1(\text{Der}(\mathcal{M}_Y, \mathcal{M}_X; \mathcal{M}_f)).$$

Proof. For X and Y simply connected, part (a) is proven in Theorem 2.1 of [10]. We observe that all parts of that proof remain valid if X and Y are only assumed nilpotent since the same ingredients are available in the nilpotent setting. The argument given in [10] to show that Φ_f is well-defined also remains valid for X and Y nilpotent and $n = 1$. We point out, however, that our argument given in [10] to show that Φ_f is a homomorphism breaks down for $n = 1$, since at one point it would use the auxiliary space $S^1 \vee S^1$ which is non-nilpotent. For this reason, we give more details for parts (b) and (c).

Part (b) is an easy generalization of the result of Thom [13, Th.2]. We argue that Φ_f is a homomorphism. That the rationalization of Φ_f is both 1-1 and onto then follows from the argument of [10, Th.2.1]. Suppose the multiplication on Y is $m: Y \times Y \rightarrow Y$. We then have an induced multiplication $M: \text{map}(X, Y; 0) \times$

$\text{map}(X, Y; 0) \rightarrow \text{map}(X, Y; 0)$ defined by $M(g, h)(x) = m(g(x), h(x))$. Suppose that $\alpha, \beta \in \pi_n(\text{map}(X, Y; f))$ have adjoints $A, B: S^n \times X \rightarrow Y$. Then $\alpha + \beta$ may be written as a composition $M \circ (\alpha \times \beta) \circ \Delta: S^n \rightarrow \text{map}(X, Y; 0)$. The adjoint of $\alpha + \beta$, which we write as C , is given by $C = m \circ (A \times B) \circ (1 \times T \times 1) \circ \Delta: S^n \times X \rightarrow Y$. Here, $T: S^n \times X \rightarrow X \times S^n$ denotes the switching map. It follows immediately that $\mathcal{A}_C(\chi) = u \otimes \theta_A(\chi) + u \otimes \theta_B(\chi)$ and hence that Φ_f is a homomorphism.

Finally, for part (c), we identify $\pi_1(\text{map}(X, Y; f))_{\mathbb{Q}}$ with $\pi_1(\text{map}(X, Y_{\mathbb{Q}}; f_{\mathbb{Q}}))$ as in (2). Now suppose given $[\theta] \in H_1(\text{Der}(\mathcal{M}_Y, \mathcal{M}_X; \mathcal{M}_f))$. Define $\phi(\chi) = 1 \otimes \mathcal{M}_f(\chi) + s_n \otimes \theta(\chi)$ for $\chi \in \mathcal{M}_Y$. Since θ is an $\mathcal{M}_f$ -derivation that is a cycle, this defines a DG algebra map $\phi: \mathcal{M}_Y \rightarrow H^*(S^n; \mathbb{Q}) \otimes \mathcal{M}_X \cong \mathcal{M}_{S^1} \otimes \mathcal{M}_X$. By the standard correspondence between (DG homotopy classes of) maps of minimal models and (homotopy classes of) maps of rational spaces, this gives a map $F: S^1 \times X \rightarrow Y_{\mathbb{Q}}$ that satisfies $F \circ i_2 \sim f_{\mathbb{Q}}: X \rightarrow Y_{\mathbb{Q}}$. We may adjust F into a homotopic map $F': S^1 \times X \rightarrow Y_{\mathbb{Q}}$ that satisfies $F' \circ i_2 = f_{\mathbb{Q}}: X \rightarrow Y_{\mathbb{Q}}$, so that F' has as adjoint a *based map* $S^1 \rightarrow \text{map}(X, Y_{\mathbb{Q}}; f_{\mathbb{Q}})$ that represents a homotopy class $\alpha \in \pi_1(\text{map}(X, Y_{\mathbb{Q}}; f_{\mathbb{Q}}))$. Evidently, we have $\Phi_{f_{\mathbb{Q}}}(\alpha) = [\theta]$. $\square$

For our proof of Theorem 1, we require certain naturality properties of Φ_f . First, suppose given $k: Y \rightarrow Z$, with X, Y and Z connected, nilpotent CW complexes of finite type with X finite. Let $\mathcal{M}_k: \mathcal{M}_Z \rightarrow \mathcal{M}_Y$ be a minimal model. By pre-composing with $\mathcal{M}_k$, we obtain a map of chain complexes

$$(\mathcal{M}_k)^*: \text{Der}_*(\mathcal{M}_Y, \mathcal{M}_X; \mathcal{M}_f) \rightarrow \text{Der}_*(\mathcal{M}_Z, \mathcal{M}_X; \mathcal{M}_f \circ \mathcal{M}_k).$$

Then for $n \geq 2$, we have a commutative square

$$(4) \quad \begin{array}{ccc} \pi_n(\text{map}(X, Y_{\mathbb{Q}}; f_{\mathbb{Q}})) & \xrightarrow{((k_{\mathbb{Q}})_*)_{\sharp}} & \pi_n(\text{map}(X, Z_{\mathbb{Q}}; k_{\mathbb{Q}} \circ f_{\mathbb{Q}})) \\ \Phi_{f_{\mathbb{Q}}} \downarrow \cong & & \cong \downarrow \Phi_{k_{\mathbb{Q}} \circ f_{\mathbb{Q}}} \\ H_n(\text{Der}(\mathcal{M}_Y, \mathcal{M}_X; \mathcal{M}_f)) & \xrightarrow{H((\mathcal{M}_k)^*)} & H_n(\text{Der}(\mathcal{M}_Z, \mathcal{M}_X; \mathcal{M}_f \circ \mathcal{M}_k)) \end{array}$$

of group homomorphisms, with isomorphisms where indicated by part (a) of Theorem 2.1. Commutativity of the diagram follows easily from the definitions. Our proof of Theorem 1 uses this commutative diagram of group homomorphisms in the case in which $n = 2$.

We also need a naturality property in the case in which $n = 1$. Suppose given $f: X \rightarrow Y$ with X and Y connected, nilpotent CW complexes of finite type with X finite, and $k: Y \rightarrow K$ with K a connected H -space of finite type. Suppose further that $k \circ f$ is null-homotopic. Then we also have a commutative square

$$(5) \quad \begin{array}{ccc} \pi_1(\text{map}(X, Y_{\mathbb{Q}}; f_{\mathbb{Q}})) & \xrightarrow{((k_{\mathbb{Q}})_*)_{\sharp}} & \pi_1(\text{map}(X, K_{\mathbb{Q}}; 0)) \\ \Phi_{f_{\mathbb{Q}}} \downarrow & & \cong \downarrow \Phi_0 \\ H_1(\text{Der}(\mathcal{M}_Y, \mathcal{M}_X; \mathcal{M}_f)) & \xrightarrow{H((\mathcal{M}_k)^*)} & H_1(\text{Der}(\mathcal{M}_K, \mathcal{M}_X; \mathcal{M}_f \circ \mathcal{M}_k)). \end{array}$$

The top horizontal map is a group homomorphism, the right-hand vertical map is a group isomorphism by part (b) of Theorem 2.1, the bottom horizontal map is a group homomorphism and the left-hand vertical map is a *surjection of sets*.

3. PRINCIPAL FIBRATIONS AND FUNCTION SPACES

Our strategy for proving Theorem 1 is to identify the long exact homology sequence corresponding to a certain principal fibration on function spaces within the framework of chain complexes described above. Recall that a fibration $p: E \rightarrow B$ is *principal* if p is obtained as a pullback as in the left-hand diagram below:

$$(6) \quad \begin{array}{ccc} E & \xrightarrow{\bar{k}} & PK \\ p \downarrow & & \downarrow q \\ B & \xrightarrow{k} & K, \end{array} \quad \begin{array}{ccc} \mathrm{map}(X, E; \tilde{g}) & \xrightarrow{(\bar{k})_*} & \mathrm{map}(X, PK; \bar{k} \circ \tilde{g}) \\ p_* \downarrow & & \downarrow q_* \\ \mathrm{map}(X, B; g) & \xrightarrow{k_*} & \mathrm{map}(X, K; k \circ g). \end{array}$$

Here, $q: PK \rightarrow K$ is the usual path fibration over some space K and $k: B \rightarrow K$ is some map—usually referred to as the k -invariant of the (principal) fibration. The contractibility of PK implies that $E \xrightarrow{p} B \xrightarrow{k} K$ is a fibre sequence.

Lemma 3.1. *Suppose given a principal fibration $p: E \rightarrow B$ induced from a k -invariant $k: B \rightarrow K$. Let $g: X \rightarrow B$ be a map and $\tilde{g}: X \rightarrow E$ be a lift of g through p . Then the induced map $p_*: \mathrm{map}(X, E; \tilde{g}) \rightarrow \mathrm{map}(X, B; g)$ is a principal fibration induced by the k -invariant $k_*: \mathrm{map}(X, B; g) \rightarrow \mathrm{map}(X, K; 0)$. In particular,*

$$(7) \quad \mathrm{map}(X, E; \tilde{g}) \xrightarrow{p_*} \mathrm{map}(X, B; g) \xrightarrow{k_*} \mathrm{map}(X, K; 0)$$

is a fibration sequence.

Proof. Applying $\mathrm{map}(X, -)$ to the left-hand pullback diagram of (6) and restricting to appropriate components, we obtain the right-hand diagram of (6), which again is a pullback. Notice that $k \circ p$, and hence $k \circ g$, is null-homotopic. Since PK is contractible, $\mathrm{map}(X, PK)$ is connected and we have $P\mathrm{map}(X, K; k \circ g) \approx \mathrm{map}(X, PK)$. $\square$

We focus on the case $K = K(G, r+1)$ for $r \geq 1$ and describe the rationalization of the long exact homotopy sequence of (7). In this case, we may write the Sullivan minimal model of E as $\mathcal{M}_E = \mathcal{M}_B \otimes \Lambda \overline{V}$, where $\overline{V}$ is concentrated in degree r and is isomorphic to $G \otimes \mathbb{Q}$. The differential d_E restricts to d_B on $\mathcal{M}_B$ while $d_E(\overline{V}) \subseteq \mathcal{M}_B$. Furthermore, we may take the minimal model of p to be the DG algebra inclusion $\mathcal{M}_p: \mathcal{M}_B \rightarrow \mathcal{M}_E$. Now suppose given choices of minimal models $\mathcal{M}_k: \Lambda V \rightarrow \mathcal{M}_B$ and $\mathcal{M}_g: \mathcal{M}_B \rightarrow \mathcal{M}_X$ for k and g . Here, V is isomorphic to $G \otimes \mathbb{Q}$ but is concentrated in degree $r+1$ and ΛV has trivial differential. We may assume that $\mathcal{M}_k(v) = d_E(\overline{v})$ for each $v \in V$, where $v \mapsto \overline{v}$ is the obvious degree $+1$ identification of V with $\overline{V}$. Finally, let $\mathcal{M}_{\tilde{g}}: \mathcal{M}_E \rightarrow \mathcal{M}_X$ be a minimal model for $\tilde{g}$ that restricts to $\mathcal{M}_g$ on the sub-DG algebra $\mathcal{M}_B$.

By pre-composing with $\mathcal{M}_k$, as before, we obtain a map of chain complexes

$$(\mathcal{M}_k)^*: \mathrm{Der}_*(\mathcal{M}_B, \mathcal{M}_X; \mathcal{M}_g) \rightarrow \mathrm{Der}_*(\Lambda V, \mathcal{M}_X; \mathcal{M}_g \circ \mathcal{M}_k).$$

Using the mapping cone construction for chain maps (cf. [12, p.166]), we may place the homomorphism induced by $(\mathcal{M}_k)^*$ on homology into a long exact sequence of

$$\begin{array}{c}
 \cdots \longrightarrow H_{n+1}(\mathrm{Der}(\Lambda V, \mathcal{M}_X; \mathcal{M}_{\bar{g}})) \xrightarrow{J_{n+1}} H_{n+1}(\mathrm{Rel}((\mathcal{M}_k)^*)) \\
 \downarrow P_n \searrow \hspace{10em} \nearrow \\
 {}_B, \mathcal{M}_X; \mathcal{M}_{\bar{q}}) \xrightarrow{H((\mathcal{M}_k)^*)} H_n(\mathrm{Der}(\Lambda V, \mathcal{M}_X; \mathcal{M}_{\bar{q}})) \xrightarrow{J_n} \dots
 \end{array}$$
$$\mathrm{Rel}_n((\mathcal{M}_k)^*) = \mathrm{Der}_{n-1}(\mathcal{M}_B, \mathcal{M}_X; \mathcal{M}_q) \oplus \mathrm{Der}_n(\Lambda V, \mathcal{M}_X; \mathcal{M}_{\bar{a}})$$
$$\delta(\theta, \varphi) = (-D(\theta), (\mathcal{M}_k)^*(\theta) + D'(\varphi)).$$

Lemma 3.2. *Given the above notation, there is a chain isomorphism*

$$\Psi: \mathrm{Rel}_*((\mathcal{M}_k)^*) \rightarrow \mathrm{Der}_{*-1}(\mathcal{M}_E, \mathcal{M}_X; \mathcal{M}_{\tilde{q}})$$

$$H_2(\mathrm{Rel}_*((\mathcal{M}_k)^*)) \xrightarrow{\cong} H_1(\mathrm{Der}(\mathcal{M}_E, \mathcal{M}_X; \mathcal{M}_{\tilde{q}})) .$$

We will use the following result as the induction step in our proof of Theorem 1. Once again, the hypotheses are indicated in diagram (6).

$$\mathrm{rank}(\pi_1(\mathrm{map}(X, B; g))) = \dim_{\mathbb{Q}}(H_1(\mathrm{Der}(\mathcal{M}_B, \mathcal{M}_X; \mathcal{M}_q))),$$
$$\mathrm{rank}(\pi_1(\mathrm{map}(X, E; \widetilde{g}))) = \dim_{\mathbb{Q}}(H_1(\mathrm{Der}(\mathcal{M}_E, \mathcal{M}_X; \mathcal{M}_{\widetilde{g}}))).$$
$$(k_*)_{\#}: \pi_1(\mathrm{map}(X, B; g)) \rightarrow \pi_1(\mathrm{map}(X, K(G, r+1); 0))$$
$$(k_*)_{\#}: \pi_2(\mathrm{map}(X, B; g)) \rightarrow \pi_2(\mathrm{map}(X, K(G, r+1); 0)).$$
$$I = \text{im}(k_*)_{\#}: \pi_1(\text{map}(X, B; q)) \rightarrow \pi_1(\text{map}(X, K(G, r+1); 0)).$$

Then $\text{rank}(\ker(k_*)_{\#}) = \text{rank}(\pi_1(\text{map}(X, B; g))) - \text{rank}(I)$, and we have

$$(8) \quad \text{rank}(\pi_1(\text{map}(X, E; \tilde{g}))) = \text{rank}(\pi_1(\text{map}(X, B; g))) - \text{rank}(I) + \text{rank}(C).$$

Now we establish a comparable expression in the framework of chain complexes of derivations of minimal models. Choose a minimal model $\mathcal{M}_k: (\Lambda V, d=0) \rightarrow \mathcal{M}_B$ of the k -invariant $k: B \rightarrow K(G, r+1)$. Here, V is a vector space isomorphic to $G \otimes \mathbb{Q}$ concentrated in degree $r+1$. Pre-composition with $\mathcal{M}_k$ gives the map of chain complexes $(\mathcal{M}_k)^*: \text{Der}_*(\mathcal{M}_B, \mathcal{M}_X; \mathcal{M}_g) \rightarrow \text{Der}_*(\Lambda V, \mathcal{M}_X; \mathcal{M}_g \circ \mathcal{M}_k)$ that are featured in Lemma 3.2. The long exact sequence in homology obtained by applying the mapping cone construction to $(\mathcal{M}_k)^*$ displays the vector space $H_2(\text{Rel}((\mathcal{M}_k)^*))$ as a vector space isomorphic to the direct sum of the cokernel of

$$H((\mathcal{M}_k)^*): H_2(\text{Der}(\mathcal{M}_B, \mathcal{M}_X; \mathcal{M}_g)) \rightarrow H_2(\text{Der}(\Lambda V, \mathcal{M}_X; \mathcal{M}_g \circ \mathcal{M}_k))$$

and the kernel of

$$H((\mathcal{M}_k)^*): H_1(\text{Der}(\mathcal{M}_B, \mathcal{M}_X; \mathcal{M}_g)) \rightarrow H_1(\text{Der}(\Lambda V, \mathcal{M}_X; \mathcal{M}_g \circ \mathcal{M}_k)).$$

Write this cokernel as $\mathcal{C}$ and set

$$\mathcal{I} = \text{im } H((\mathcal{M}_k)^*): H_1(\text{Der}(\mathcal{M}_B, \mathcal{M}_X; \mathcal{M}_g)) \rightarrow H_1(\text{Der}(\Lambda V, \mathcal{M}_X; \mathcal{M}_g \circ \mathcal{M}_k))$$

so that we obtain

$$(9) \quad \begin{aligned} \dim_{\mathbb{Q}}(H_2(\text{Rel}((\mathcal{M}_k)^*))) &= \dim_{\mathbb{Q}}(H_1(\text{Der}(\mathcal{M}_B, \mathcal{M}_X; \mathcal{M}_g))) \\ &\quad - \dim_{\mathbb{Q}}(\mathcal{I}) + \dim_{\mathbb{Q}}(\mathcal{C}). \end{aligned}$$

Now, part (a) of Theorem 2.1 and (4) imply that $\text{rank}(C) = \dim_{\mathbb{Q}}(\mathcal{C})$, and parts (b) and (c) of Theorem 2.1, and (5) imply that $\text{rank}(I) = \dim_{\mathbb{Q}}(\mathcal{I})$. Therefore, by (8) and (9), and our hypothesis, we have that

$$\text{rank}(\pi_1(\text{map}(X, E; \tilde{g}))) = \dim_{\mathbb{Q}}(H_2(\text{Rel}((\mathcal{M}_k)^*))).$$

The result now follows from Lemma 3.2. $\square$

4. CONCLUSION

In this section, we prove our main result and its corollary and give a sample application.

Proof of Theorem 1. We obtain by induction the desired formula for an arbitrary Postnikov section Y^r of Y . That is, we show that

$$\text{rank}(\pi_1(\text{map}(X, Y^r; f^r))) = \dim_{\mathbb{Q}}(H_1(\text{Der}(\mathcal{M}_{Y^r}, \mathcal{M}_X; \mathcal{M}_{f^r})))$$

for any given $r \geq 0$, where $f^r: X \rightarrow Y^r$ denotes the map induced by $f: X \rightarrow Y$. Since $Y^0 = \{*\}$, the induction starts trivially.

For the induction step, recall that, since Y is a nilpotent space, by [9, Th.II.2.9] the Postnikov decomposition of Y admits a *principal refinement* at each stage. That is, $p_r: Y^r \rightarrow Y^{r-1}$ —the r th stage of the Postnikov decomposition of Y —factors into a finite sequence of principal fibrations (c_r of them, say)

$$Y^r = Y_{c_r}^r \rightarrow Y_{c_{r-1}}^r \rightarrow \cdots \rightarrow Y_1^r \rightarrow Y_0^r = Y^{r-1},$$

each induced from a path-loop fibration by a k -invariant of the form $k_j^r: Y_{j-1}^r \rightarrow K(G_j^r, r+1)$. Using Proposition 3.3 for each stage of the principal refinement completes the induction.

To complete the proof of Theorem 1 we observe that, by obstruction theory, the r -equivalence $p_r: Y \rightarrow Y^r$ induces an $(r-N)$ -equivalence $(p_r)_*: \text{map}(X, Y; f) \rightarrow$

Here $(\rho_Y)^*$ and $(d_Y)^*$ are the maps induced by pre-composition. The map $\mathcal{H}$ is defined as follows. Given $\theta \in \text{Der}_1(\Lambda V_1, H^*(X; \mathbb{Q}); 0)$, first define a linear map $H(\theta): V \rightarrow \mathcal{M}_X$ by setting $H(\theta)(V_0) = \{0\}$ and, on a basis element of $y \in V_1$, $H(\theta(y)) = P \in \mathcal{M}_X$, where P is representative of the class $\theta(y) \in H(\mathcal{M}_X) \cong H^*(X; \mathbb{Q})$. Extend $H(\theta)$ to a linear map of degree -1 , then to an element of $\text{Der}_1(\mathcal{M}_Y, \mathcal{M}_X; \mathcal{M}_f)$, by the $\mathcal{M}_f$ -derivation law. The result is a cycle derivation $H(\theta)$. Finally, let $\mathcal{H}(\theta)$ be the homology class of $H(\theta)$. Grivel's proof of [6, Th.4.4] is directly adapted to show $\mathcal{H}$ is well-defined and the sequence is exact. $\square$

As a consequence of Theorem 4.1, we deduce, for instance, the following example.

Example 4.2. Let G be a connected, compact, simple Lie group of rank $n > 1$ and $T^n \subseteq G$ a maximal torus. Let $f: S^2 \rightarrow G/T^n$ be any given map. We show

$$\pi_1(\text{map}(S^2, G/T^n; f)) \text{ is a finite group if and only if } f \text{ is essential.}$$

The space G/T^n is an F_0 -space with rational cohomology generated in degree 2 and we have $b_2(G/T^n) = \rho_2(G/T^n) = n$. It is well-known that $\rho_3(G) = 1$, and hence $\rho_3(G/T^n) = 1$ (see, for instance, p.426 of [1]). Thus by Theorem 4.1, we have $\text{rank}(\pi_1(\text{map}(S^2, G/T^n; f))) = D_2(f) + 1 - n$. If f is rationally null-homotopic, then $\text{rank}(\pi_1(\text{map}(S^2, G/T^n; 0))) = 1$ by Corollary 1. Suppose f is rationally essential. Fix an additive basis $\{t_1, \dots, t_n\}$ for $H^2(Y; \mathbb{Q})$ and suppose, say, $H^*(f)(t_i) \neq 0$. Let $\theta_i \in \text{Hom}_2(H^*(Y; \mathbb{Q}), H^*(X; \mathbb{Q}))$ be defined on this basis by $\theta_i(t_i) = 1$ while $\theta_i(t_j) = 0$ for $j \neq i$. Suppose θ_i extends to an $H^*(f)$ -derivation. Since the Weyl group of G is a finite reflection group, the cohomology class $t_1^2 + \dots + t_n^2$ vanishes in $H^4(G/T^n; \mathbb{Q})$. However, $\theta_i(t_1^2 + \dots + t_n^2) = 2H^*(f)(t_i)\theta_i(t_i) = 2H^*(f)(t_i) \neq 0$. This contradiction implies $D_2(f) \leq n - 1$ and so $\pi_1(\text{map}(S^2, G/T^n; f))$ must be a finite group in this case. Finally, by the result of E. Cartan, $\pi_2(G) = 0$ and so $\pi_2(G/T^n)$ is free abelian. Thus $f: S^2 \rightarrow G/T^n$ is essential if and only if it is rationally essential.

REFERENCES

- [1] A. Borel, *Topology of Lie groups and characteristic classes*, Bull. Amer. Math. Soc. **61** (1955), 397–432. MR0072426 (17:282b)
- [2] A. K. Bousfield and V. K. A. M. Gugenheim, *On PL de Rham theory and rational homotopy type*, Mem. Amer. Math. Soc. **8** (1976), no. 179, ix+94. MR0425956 (54:13906)
- [3] E. H. Brown, Jr. and R. H. Szczarba, *On the rational homotopy type of function spaces*, Trans. Amer. Math. Soc. **349** (1997), no. 12, 4931–4951. MR1407482 (98c:55015)
- [4] U. Buijs and A. Murillo, *The rational homotopy Lie algebra of function spaces*, preprint.
- [5] Y. Félix, S. Halperin, and J.-C. Thomas, *Rational homotopy theory*, Graduate Texts in Mathematics, vol. 205, Springer-Verlag, New York, 2001. MR1802847 (2002d:55014)
- [6] P.-P. Grivel, *Algèbres de Lie de dérivations de certaines algèbres pures*, J. Pure Appl. Algebra **91** (1994), no. 1-3, 121–135. MR1255925 (95a:55024)
- [7] A. Haefliger, *Rational homotopy of the space of sections of a nilpotent bundle*, Trans. Amer. Math. Soc. **273** (1982), no. 2, 609–620. MR0667163 (84a:55010)
- [8] S. Halperin, *Finiteness in the minimal models of Sullivan*, Trans. Amer. Math. Soc. **230** (1977), 173–199. MR0461508 (57:1493)
- [9] P. Hilton, G. Mislin, and J. Roitberg, *Localization of nilpotent groups and spaces*, North-Holland Publishing Co., Amsterdam, 1975, North-Holland Mathematics Studies, No. 15. MR0478146 (57:17635)
- [10] G. Lupton and S. B. Smith, *Rationalized evaluation subgroups of a map I: Sullivan models, derivations, and G-sequences*, J. Pure Appl. Algebra, **209** (2007) 159–171.
- [11] J. Milnor, *On spaces having the homotopy type of CW-complex*, Trans. Amer. Math. Soc. **90** (1959), 272–280. MR0100267 (20:6700)
- [12] E. H. Spanier, *Algebraic topology*, Springer-Verlag, New York, 1989. MR0666554 (83i:55001)
- [13] R. Thom, *L'homologie des espaces fonctionnels*, Colloque de topologie algébrique, Louvain, 1956, Georges Thone, Liège, 1957, pp. 29–39. MR0089408 (19:669h)
- [14] M. Vigué-Poirrier, *Sur l'homotopie rationnelle des espaces fonctionnels*, Manuscripta Math. **56** (1986), 177–191. MR0850369 (87h:55009)

DEPARTMENT OF MATHEMATICS, CLEVELAND STATE UNIVERSITY, CLEVELAND, OHIO 44115

E-mail address: `G.Lupton@csuohio.edu`

DEPARTMENT OF MATHEMATICS, SAINT JOSEPH'S UNIVERSITY, PHILADELPHIA, PENNSYLVANIA
19131

E-mail address: `smith@sju.edu`