BANACH-STONE THEOREM FOR BANACH LATTICE VALUED CONTINUOUS FUNCTIONS

Z. ERCAN AND S. ÖNAL

(Communicated by Joseph A. Ball)

ABSTRACT. Let X and Y be compact Hausdorff spaces, E be a Banach lattice and F be an AM space with unit. Let $\pi : C(X, E) \rightarrow C(Y, F)$ be a Riesz isomorphism such that $0 \notin f(X)$ if and only if $0 \notin \pi(f)(Y)$ for each $f \in C(X, E)$. We prove that X is homeomorphic to Y and E is Riesz isomorphic to F . This generalizes some known results.

INTRODUCTION

In this paper we use the standard terminology and notation of the Riesz spaces theory. For unexplained definitions and notation we refer to [1]. Throughout this paper X and Y stand for compact Hausdorff spaces. For a compact Hausdorff space Z and a Banach lattice E , the Banach lattice (under pointwise algebraic operations and order) of continuous functions from Z into E is denoted by $C(Z, E)$. If $E = \mathbb{R}$ we write $C(Z)$ instead of $C(Z, E)$. $\mathbf{1}_Z \in C(Z)$ is defined by $\mathbf{1}_Z(x) = 1$.

One of the versions of the Banach-Stone Theorem states that X and Y are homeomorphic if and only if $C(X)$ and $C(Y)$ are Riesz isomorphic. More precisely, if $\pi : C(X) \rightarrow C(Y)$ is a Riesz homomorphism, then there exists a homeomorphism $\sigma : Y \rightarrow X$ such that $\pi(f) = \pi(\mathbf{1}_X)(f\sigma)$. A simple and direct proof of this theorem can be found in [5]. This theorem is generalized in [2] as follows:

Theorem 1. *Let E be a Banach lattice and $\pi : C(X, E) \rightarrow C(Y)$ be a Riesz isomorphism such that $\pi(f)(y) \neq 0$ for each $y \in Y$ whenever $f(x) \neq 0$ for each $x \in X$. Then X is homeomorphic to Y and E is Riesz isomorphic to $E = \mathbb{R}$.*

A simple and direct proof of the above theorem is given in [3], also providing a positive answer to a conjecture in [2]. By using Theorem 2 of [4] it can also be proved that the above theorem is still true for d -isomorphism π ; that is, $x \perp y$ if and only if $\pi(x) \perp \pi(y)$.

THE MAIN RESULT

The aim of this paper is to generalize the above theorem as follows:

Received by the editors June 16, 2005 and, in revised form, May 21, 2006.

2000 *Mathematics Subject Classification.* Primary 46E40; Secondary 46B42.

Key words and phrases. Riesz isomorphism, Banach lattices, Banach-Stone Theorem.

Theorem 2. Let E be a Banach lattice and F be an AM space with unit. Let $\pi : C(X, E) \longrightarrow C(Y, F)$ be a Riesz isomorphism. If

$$f(x) \neq 0 \text{ for each } x \in X \iff \pi f(y) \neq 0 \text{ for each } y \in Y,$$

then X is homeomorphic to Y and E is Riesz isomorphic to F .

To prove this theorem we will need the following lemmas. The proofs of the first two lemmas are clear.

Lemma 3. Let M and N be compact Hausdorff spaces and $\sigma : X \times M \longrightarrow Y \times N$ be a homeomorphism. If for each $x \in X$ there exists a unique $y \in Y$ such that $\sigma(\{x\} \times M) = \{y\} \times N$, then X and M are homeomorphic to Y and N , respectively.

Lemma 4. Let $f : X \longrightarrow Y$ be a continuous open surjective map. If K is a compact subset of Y and O is an open subset of X with $K \subset f(O)$, then there is a closed subset F of X such that $F \subset O$ and $K \subset f(F)$.

Lemma 5. Let M and N be compact Hausdorff spaces and $\pi : C(X \times M) \longrightarrow C(Y \times N)$ be a Riesz isomorphism with $\pi(\mathbf{1}_{X \times M}) = \mathbf{1}_{Y \times N}$. If

$$f(\{x\} \times M) \neq 0 \text{ for each } x \in X \iff \pi f(\{y\} \times N) \neq 0 \text{ for each } y \in Y,$$

then X and M are homeomorphic to Y and N , respectively.

Proof. From the Banach-Stone theorem there exists a homeomorphism $\sigma : Y \times N \longrightarrow X \times M$ such that

$$\pi(f) = f\sigma \text{ and } \pi^{-1}(g) = g\sigma^{-1}.$$

Let P_X be the projection of $X \times M$ onto X ; that is, $P_X : X \times M \longrightarrow X$ is defined by $P_X(x, m) = x$.

Claim a: For each $y_0 \in Y$, $P_X\sigma((Y - \{y_0\}) \times N) \neq X$. Indeed, suppose it does not hold. Note that $P_X\sigma : Y \times N \longrightarrow X$ is a surjective open map and $(Y - \{y_0\}) \times N$ is open in $Y \times N$. Since $X \subset P_X\sigma((Y - \{y_0\}) \times N)$ there exists a closed set F in $Y \times N$ such that

$$F \subset Y - \{y_0\} \times N \text{ and } X \subset P_X\sigma(F).$$

Let $A = \sigma(F)$ and $B = \sigma(\{y_0\} \times N)$. Clearly A and B are closed in $Y \times N$ and $A \cap B = \emptyset$. Then there exists $f \in C(Y \times N)$ such that

$$f(A) = \{1\} \text{ and } f(B) = \{0\}.$$

Let $x \in X$ be given. Since $X = P_X\sigma(F)$, there exists $m \in M$, $(y, n) \in F$ such that

$$x = P_X(x, m) = P_X\sigma(y, n).$$

Then $\sigma(y, n) \in A$ and so $1 = f(\sigma(y, n)) = f(x, m)$. From the hypothesis, $\pi f(\{y\} \times N) \neq \{0\}$ for each $y \in Y$, in particular,

$$\{0\} = f(B) = \pi f(\{y_0\} \times N) \neq \{0\}.$$

This is a contradiction.

Claim b: For each $n \in N$, $P_X\sigma(Y \times \{n\}) = X$. To see this, suppose that $x_0 \notin P_X\sigma(Y \times \{n\})$ for some $x_0 \in X$ and $n \in N$. Since $\{x_0\} \times M$ and $\sigma(Y \times \{n\})$ are nonempty disjoint closed sets, there exists $f \in C(X \times M)$ such that

$$f\sigma(Y \times \{n\}) = \{1\} \text{ and } f(\{x_0\} \times N) = \{0\}.$$

That implies that $\pi f(Y \times \{n_0\}) = \{1\}$, so $f(\{x_0\} \times N) \neq \{0\}$. This contradiction shows that Claim b is also true.

Claim c: For each $y \in Y$ there exists a unique $x \in X$ such that $\sigma^{-1}(\{y\} \times N) = \{x\} \times M$. Let $y \in Y$ be given. Since for each $n \in N$,

$$P_X \sigma((Y - \{y\}) \times N) \neq P_X \sigma(Y \times \{n\}) = X,$$

there exists $x \in X$ such that $(x, m) \notin \sigma(Y - \{y\} \times N)$ for each $m \in M$, so $\{x\} \times M \subset \sigma(\{y\} \times N)$. Hence $\sigma^{-1}(\{x\} \times M) \subset \{y\} \times N$. Similarly there exists $y_1 \in Y$ such that $\sigma(\{y_1\} \times N) \subset \{x\} \times M$. From

$$\{y_1\} \times N \subset \sigma^{-1}(\{x\} \times M) \subset \{y\} \times N$$

we have $y_1 = y$. Now, it is clear that $\sigma^{-1}(\{x\} \times M) = \{y\} \times N$ and y must be unique. Now Lemma 3 is applied to complete the proof. $\square$

Now we are ready to give the proof of the theorem.

Proof of Theorem 2. From the Kakutani Representation Theorem there exists a compact Hausdorff space N such that F is isometrically Riesz isomorphic to $C(N)$. Since for each $x \in X$, $\pi^{-1}(\mathbf{1}_Y \otimes e)(x)$ is a strong order unit for E ($\mathbf{1}_Y \otimes e$ is defined by $\mathbf{1}_Y \otimes e(y) = e$), E is (norm) isomorphic and Riesz isomorphic to $C(M)$ for some compact Hausdorff space. Since for compact Hausdorff spaces Y and Z , the Banach lattices $C(Y, C(Z))$ and $C(Y \times Z)$ are Riesz isomorphic spaces, we have that $C(X \times M)$ and $C(Y \times N)$ are Riesz isomorphic spaces. Moreover, it is clear that from the hypotheses of the theorem there exists a Riesz isomorphism $\pi_1 : C(X \times M) \rightarrow C(Y \times N)$ such that $\pi_1(\mathbf{1}_{X \times M}) = \mathbf{1}_{Y \times N}$ and

$$f(\{x\} \times M) \neq 0 \quad \text{for each } x \in X \iff \pi_1 f(\{y\} \times N) \neq 0 \quad \text{for each } y \in Y.$$

Hence from Lemma 5, X and M are homeomorphic to Y and N respectively, so E is (norm) isomorphic and Riesz isomorphic to F . $\square$

Let X and Y be compact Hausdorff spaces, and E and F be Banach lattices. Let $\pi : C(X, E) \rightarrow C(Y, F)$ be a Riesz isomorphism such that $0 \notin f(X)$ if and only if $0 \notin \pi(f)(Y)$. We conjecture that X and Y are homeomorphic and E and F are Riesz isomorphic.

REFERENCES

- [1] C. D. Aliprantis and O. Burkinshaw, *Positive Operators*, Academic Press (1985). MR0809372 (87h:47086)
- [2] J. Cao, I. Reilly and H. Xiong, *A lattice-valued Banach-Stone Theorem*, Acta Math. Hungar., **98** (1-2), (2003), 103-110. MR1958470 (2003m:46028)
- [3] Z. Ercan and S. Onal, *An answer to a conjecture of Cao, Reilly and Xiong*, to appear in Czechoslovak Math. J.
- [4] J.J. Font and S. Hernandez, *On separating maps between locally compact spaces*, Arch. Math., Vol. **63**, (1994), 158-165. MR1289298 (95k:46083)
- [5] E. de Jonge and A.C.M. van Rooij, *Introduction to Riesz Spaces* (Mathematical Center Tracts **78**), Mathematisch Centrum, Amsterdam, 1977. MR0473777 (57:13439)

DEPARTMENT OF MATHEMATICS, MIDDLE EAST TECHNICAL UNIVERSITY, 06531 ANKARA, TURKEY

E-mail address: zercan@metu.edu.tr

DEPARTMENT OF MATHEMATICS, MIDDLE EAST TECHNICAL UNIVERSITY, 06531 ANKARA, TURKEY

E-mail address: osul@metu.edu.tr