ON MOD 3 GALOIS REPRESENTATIONS WITH CONDUCTOR 4

HYUNSUK MOON

(Communicated by Ken Ono)

ABSTRACT. The non-existence is proved of 2-dimensional mod 3 irreducible representations of $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ of Artin conductor dividing 4.

Let $G_{\mathbb{Q}}$ be the absolute Galois group $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ of $\mathbb{Q}$. For any prime number ℓ , let $\overline{\mathbb{F}}_{\ell}$ denote an algebraic closure of the finite field $\mathbb{F}_{\ell}$ of ℓ elements. In this paper, we prove the non-existence of certain mod 3 Galois representations:

Theorem 1. *There exist no irreducible representations $\rho : G_{\mathbb{Q}} \rightarrow \text{GL}_2(\overline{\mathbb{F}}_3)$ with $N(\rho)$ dividing 4.*

Here, $N(\rho) = \prod_{p \nmid 3} p^{n_p(\rho)}$ is the Artin conductor of ρ outside 3 ([7], §1.2; the definition of the exponent $n_p(\rho)$ will be recalled below). This proves a special case of Serre’s conjecture ([7]). Indeed, the conjecture predicts that if such a representation is odd, then up to twist by a power of the mod 3 cyclotomic character, it comes from a cuspidal eigenform of level 4 and weight ≤ 4 , but there are no such forms. Khare ([3]) proved the level 1 case of Serre’s conjecture by induction on the primes. Our result may serve as a first step of such an inductive proof of the level 4 case of Serre’s conjecture if Khare’s proof can be extended to this case.*

Serre’s conjecture is known to be true if ρ is odd and its image $\text{Im}(\rho)$ of ρ is solvable ([5], Thm. 4). So, it remains for us to prove the theorem in the following two cases: (i) $\text{Im}(\rho)$ is non-solvable, (ii) ρ is even and $\text{Im}(\rho)$ is solvable.

Our strategy in case (i) is basically the same as in [10]; that is to deduce a contradiction by comparing two kinds of inequalities of the opposite direction for the discriminant of the field corresponding to the kernel of ρ — one from above (the refined Tate bound ([5], Thm. 3) and the other from below (the Odlyzko bound [6]). A new ingredient in this paper is the estimate of the prime-to-3 part of the discriminant (Lemma 4). The case (ii) is settled by class field theory with the help of known results ([9]) on solvable subgroup of $\text{GL}_2(\overline{\mathbb{F}}_{\ell})$ and Jones’ table ([2]) of number fields.

In Section 1, we analyze the ramification of ρ at 2. The proof of the theorem is given in Section 2.

Received by the editors March 2, 2006 and, in revised form, July 24, 2006.

2000 *Mathematics Subject Classification.* Primary 11F80, 11S15.

*Note added in proof: Serre’s conjecture has recently been proved completely by Khare and Wintenberger independently of our work.

1. RAMIFICATION AT 2

Let $D_p (\subset G_{\mathbb{Q}})$ be the decomposition subgroup for a choice of an extension of the prime ideal (p) to $\overline{\mathbb{Q}}$, and I_p its inertia subgroup. For a continuous representation $\rho : D_p \rightarrow \mathrm{GL}(V)$, where V is a finite-dimensional $\overline{\mathbb{F}}_{\ell}$ -vector space with $\ell \neq p$, we define the *exponent of Artin conductor* of ρ by

$$n_p(\rho) := \sum_{i=0}^{\infty} \frac{1}{(G_0 : G_i)} \dim_{\overline{\mathbb{F}}_{\ell}}(V/V^{G_i}).$$

Here, G_i is the i th ramification subgroup of $G := \mathrm{Im}(\rho)$.

Lemma 2. *Let p and ℓ be two distinct prime numbers, and let $\rho : D_p \rightarrow \mathrm{GL}_2(\overline{\mathbb{F}}_{\ell})$ be a continuous representation with $n_p(\rho) = 2$. If ρ is wildly ramified at p , then it is a direct sum of two characters, one of which is unramified and the other has exponent of Artin conductor 2. In particular, if ρ is irreducible, then it is tamely ramified at p .*

Remark. This lemma holds true if D_p is the absolute Galois group of any complete discrete valuation field with finite residue field of characteristic p .

Proof. Suppose ρ is wildly ramified at p , so that $\dim(V/V^{G_1}) < 2$. By assumption, we have

$$(*) \quad n_p(\rho) = \dim(V/V^{G_0}) + \frac{1}{(G_0 : G_1)} \dim(V/V^{G_1}) + \cdots = 2.$$

This implies that $\dim(V/V^{G_0}) = 1$. Then V is reducible as a representation of D_p , because G_0 is normal in G and hence G stabilizes V^{G_0} . We may assume that ρ is of the form

$$\rho = \begin{pmatrix} \psi_1 & * \\ 0 & \psi_2 \end{pmatrix},$$

where $\psi_i : D_p \rightarrow \overline{\mathbb{F}}_{\ell}^{\times}$ are characters of D_p and ψ_1 is unramified. Let $\rho^{\mathrm{ss}} = \begin{pmatrix} \psi_1 & 0 \\ 0 & \psi_2 \end{pmatrix}$ be the semisimplification of ρ and put $G^{\mathrm{ss}} := \mathrm{Im}(\rho^{\mathrm{ss}})$. Then G sits in a short exact sequence

$$1 \rightarrow H \rightarrow G \rightarrow G^{\mathrm{ss}} \rightarrow 1,$$

where $H = G \cap \left\{ \begin{pmatrix} 1 & * \\ 0 & 1 \end{pmatrix} \right\}$ is the kernel of the natural homomorphism $G \rightarrow G^{\mathrm{ss}}$. Note that G^{ss} is abelian of order prime to ℓ , and H is an elementary abelian ℓ -group of rank at most 2. Let $H_0 := H \cap G_0$. If $H_0 \neq 1$, then it is mapped by the projection $G_0 \rightarrow G_0/G_1$ to the unique ℓ -Sylow subgroup of the tame inertia group G_0/G_1 . Let G_0^b be the inverse image in G_0 of the maximal prime-to- ℓ subgroup of the cyclic group G_0/G_1 . Then H_0 and G_0^b are both normal in G_0 , $H_0 G_0^b = G_0$, and $H_0 \cap G_0^b = 1$. Hence we have $G_0 = H_0 \times G_0^b$. But this is impossible, because any two non-trivial elements of the form $\begin{pmatrix} * & * \\ 0 & * \end{pmatrix}$, one of order ℓ and the other of order prime to ℓ , do not commute. (Note that $G_0^b \neq 1$, as $G_1 \neq 1$.) Hence $H_0 = 1$ and G_0 has order prime to ℓ . Next we argue in the same way with G/G_0 in place of G_0/G_1 . If $H \neq 1$, then it is mapped by the projection $G \rightarrow G/G_0$ to the unique ℓ -Sylow subgroup of G/G_0 . Let G^b be the inverse image in G of the maximal prime-to- ℓ subgroup of the cyclic group G/G_0 . Then H and G^b are both normal in G , $H G^b = G$, and $H \cap G^b = 1$. Hence $G = H \times G^b$. But this is again impossible by the same reason as above. Hence $H = 1$ and $G = G^{\mathrm{ss}}$ is semisimple. $\square$

The next lemma in group theory is proved in §§19–21 of [9], and it is used in the proof of Lemma 4 and in Section 2.

Lemma 3. *Let G be an irreducible solvable subgroup of $\mathrm{GL}_2(\overline{\mathbb{F}}_\ell)$ and $\overline{G}$ the image of G in $\mathrm{PGL}_2(\overline{\mathbb{F}}_\ell)$. Then G is either of the following two types.*

- (a) *Imprimitive case: G is isomorphic to a subgroup of the wreath product $\overline{\mathbb{F}}_\ell^\times \wr (\mathbb{Z}/2\mathbb{Z})$.*
- (b) *Primitive case: $\overline{G}$ sits in an exact sequence*

$$1 \rightarrow \overline{A} \rightarrow \overline{G} \rightarrow \overline{H} \rightarrow 1,$$

where $\overline{A}$ is isomorphic to a subgroup of $(\mathbb{Z}/2\mathbb{Z})^{\oplus 2}$ and the conjugate action of $\overline{H}$ on $\overline{A}$ is faithful, so that $\overline{H}$ is identified with a subgroup of $\mathrm{Aut}(\overline{A})$. In particular, $\overline{G}$ is either a 2-group or isomorphic to the symmetric group S_4 or the alternating group A_4 .

In the special case $p = 2$ and $\ell = 3$, Lemma 2 can be strengthened as follows.

Lemma 4. *Let $\rho : D_2 \rightarrow \mathrm{GL}_2(\overline{\mathbb{F}}_3)$ be a continuous representation with $n_2(\rho) = 2$. Then it is a direct sum of two characters, one of which is unramified and the other has exponent of Artin conductor 2. If G_i denotes the i th ramification subgroup of $G := \mathrm{Im}(\rho)$, then one has $G_0 = G_1 \simeq \mathbb{Z}/2\mathbb{Z}$ and $G_2 = 1$. The different of the extension $K/\mathbb{Q}_2$ corresponding to $\mathrm{Ker}(\rho)$ is (2).*

Remark. This lemma holds true if D_2 is the absolute Galois group of any complete discrete valuation field with residue field $\mathbb{F}_2$.

Proof. We first show that ρ cannot be irreducible. Suppose ρ is irreducible. Then by Lemma 2, it is tamely ramified. In particular, G is meta-abelian. By Lemma 3, G is an extension of an elementary abelian 2-group $\overline{G}$ of rank at most 2 by an abelian group H of order prime to 3. Since ρ is tamely ramified, the extension $F/\mathbb{Q}_2$ corresponding to $\overline{G}$ is unramified and $\overline{G} \simeq \mathbb{Z}/2\mathbb{Z}$. Now H is the Galois group of a tamely ramified abelian extension of F . Since the residue field of F is $\mathbb{F}_4$, the inertia subgroup H_0 of H is a quotient of $\mathbb{F}_4^\times \simeq \mathbb{Z}/3\mathbb{Z}$. Since H has order prime to 3, we must have $H_0 = 1$. This contradicts the assumption that $n_2(\rho) = 2$.

Thus ρ is reducible, and we may assume that ρ is of the form

$$\rho = \begin{pmatrix} \psi_1 & * \\ 0 & \psi_2 \end{pmatrix},$$

where $\psi_i : D_2 \rightarrow \overline{\mathbb{F}}_3^\times$ are characters of D_2 . They factor through the abelianization D_2^{ab} of D_2 . Since the inertia subgroup of D_2^{ab} is isomorphic to the pro-2 group $\mathbb{Z}_2^\times$, these characters are either unramified or wildly ramified. Since $n_2(\rho) = 2$, the only possible case is that ψ_1 is unramified and ψ_2 is wildly ramified (if $* = 0$, then the role of ψ_1 and ψ_2 may be exchanged). By Lemma 2, we have $* = 0$ and $\rho \simeq \psi_1 \oplus \psi_2$. Then since $n_2(\rho) = n_2(\psi_2) = 2$, it follows that $G_0 = G_1 \simeq \mathbb{Z}_2^\times / (1 + 2^2\mathbb{Z}_2) \simeq \mathbb{Z}/2\mathbb{Z}$ and $G_2 = 1$. The statement on the different follows by the Führerdiskriminantenproduktformel [8, Ch. VI, §3, Cor. 2 to Prop. 6] $\square$

2. PROOF OF THE THEOREM

Now we can prove the theorem. Suppose there were a ρ as in the theorem. Since the case $N(\rho)|2$ was done in [4], we may assume $N(\rho) = 4$.

(i) Assume $G := \text{Im}(\rho)$ is non-solvable. Let K be the field corresponding to the kernel of ρ . Let $n := [K : \mathbb{Q}]$, and $|d_K|^{1/n}$ denote the root discriminant of K . By Theorem 3 in [5], the 3-part of $|d_K|^{1/n}$ is bounded by $3^{2+\frac{1}{6}-\frac{1}{3m}}$, where m is the wild ramification index of $K/\mathbb{Q}$ at 3. By Lemma 4, the 2-part of $|d_K|^{1/n}$ is 2. All together, we have

$$|d_K|^{1/n} \leq 3^{2+\frac{1}{6}-\frac{1}{3m}} \cdot 2 \\ \leq \begin{cases} 19.133 & \text{if } m \leq 2, \\ 21.617 & \text{for any } m. \end{cases}$$

If 3^μ is the exact power of 3 dividing the order of G , then by §251–253 of [1], the projective image $\overline{G}$ of G in $\text{PGL}_2(\overline{\mathbb{F}}_3)$ is isomorphic to either $\text{PGL}_2(\mathbb{F}_{3^\mu})$ or $\text{PSL}_2(\mathbb{F}_{3^\mu})$. Thus we have $n = |G| \geq |\text{PSL}_2(\mathbb{F}_{3^\mu})|$. Note that we have $\mu \geq 2$ because $\overline{G}$ is solvable if $\mu = 1$. By [6], we have

$$|d_K|^{1/n} > \begin{cases} 19.567 & \text{if } n \geq 360 = |\text{PSL}_2(\mathbb{F}_{3^2})|, \\ 22.021 & \text{if } n \geq 9828 = |\text{PSL}_2(\mathbb{F}_{3^3})|. \end{cases}$$

Comparing these two sets of inequalities, we obtain contradictions.

(ii) Assume ρ is even and $G = \text{Im}(\rho)$ is solvable. Note that, since ρ is even, it maps the complex conjugation to $\pm \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$, so that the field K cut out by ρ is totally real or CM. We shall show that there exists no such extension $K/\mathbb{Q}$.

If either G is of type (a) in Lemma 3 or G is of type (b) in Lemma 3 and $\overline{G}$ is a 2-group, then K contains a non-trivial abelian extension of degree prime to 3 over a real quadratic field F . Since K is unramified outside $\{2, 3\}$ and its conductor (or, exactly speaking, the conductor of ρ) at 2 is 2^2 , F is the field $\mathbb{Q}(\sqrt{3})$. Then K/F is unramified at 2 since $K/\mathbb{Q}$ has ramification index 2 at the prime 2 (Lemma 4). Since any ray class group of F of 3-power conductor has 3-power order, there are no non-trivial abelian extensions of F which are unramified outside 3 and of degree prime to 3.

Suppose now that G is of type (b) in Lemma 3 and $\overline{G}$ is isomorphic to S_4 or A_4 . According to [2], there are three S_4 -extensions (resp. one A_4 -extension) of $\mathbb{Q}$ which are unramified outside $\{2, 3\}$ and whose ramification index at 2 divides 2. However, each of these fields has 2-component of the root discriminant greater than 2, which contradicts Lemma 4. $\square$

REFERENCES

- [1] L. E. Dickson, *Linear Groups*, Teubner, 1901, Leipzig.
- [2] J. Jones, *Tables of number fields with prescribed ramification*, <http://math.la.asu.edu/~jj/>.
- [3] C. Khare, *On Serre's modularity conjecture: the level one case*, *Duke Math. J.* **134** (2006), 557–589. MR2254626
- [4] H. Moon, *The non-existence of certain mod p Galois representations*, *Bull. Korean Math. Soc.* **40** (2003), 537–544. MR1996862 (2004d:11044)
- [5] H. Moon and Y. Taguchi, *Refinement of Tate's discriminant bound and non-existence theorems for mod p Galois representations*, *Documenta Math. Extra Volume: Kazuya Kato's Fiftieth Birthday* (2003), 641–654. MR2046611 (2005a:11069)
- [6] A. M. Odlyzko, *Discriminant bounds*, unpublished manuscript (1976), available at: <http://www.dtc.umn.edu/~odlyzko/unpublished/discr.bound.table2>.
- [7] J.-P. Serre, *Sur les représentations modulaires de degré 2 de $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$* , *Duke Math. J.* **54** (1987), 179–230. MR885783 (88g:11022)
- [8] J.-P. Serre, *Local fields*, Springer-Verlag, New York, 1979.

- [9] D.A. Suprunenko, *Matrix Groups*, A.M.S., Providence, 1976. MR0390025 (52:10852)
- [10] J. Tate, *The non-existence of certain Galois extensions of $\mathbb{Q}$ unramified outside 2*, Contemp. Math. **174** (1994), 153–156. MR1299740 (95i:11132)

DEPARTMENT OF MATHEMATICS, COLLEGE OF NATURAL SCIENCES, KYUNGPOOK NATIONAL UNIVERSITY, DAEGU 702-701, KOREA

E-mail address: `hsmoon@knu.ac.kr`