L-FUNCTIONS OF TWISTED DIAGONAL EXPONENTIAL SUMS OVER FINITE FIELDS

SHAOFANG HONG

(Communicated by Wen-Ching Winnie Li)

ABSTRACT. Let $\mathbf{F}_q$ be the finite field of q elements with characteristic p and $\mathbf{F}_{q^m}$ its extension of degree m . Fix a nontrivial additive character Ψ and let $\chi_1, \dots, \chi_n$ be multiplicative characters of $\mathbf{F}_p$. For

$$f(x_1, \dots, x_n) \in \mathbf{F}_q[x_1, x_1^{-1}, \dots, x_n, x_n^{-1}],$$

one can form the twisted exponential sum $S_m^*(\chi_1, \dots, \chi_n, f)$. The corresponding L -function is defined by

$$L^*(\chi_1, \dots, \chi_n, f; t) = \exp\left(\sum_{m=0}^{\infty} S_m^*(\chi_1, \dots, \chi_n, f) \frac{t^m}{m}\right).$$

In this paper, by using the p -adic gamma function and the Gross–Koblitz formula on Gauss sums, we give an explicit formula for the L -function $L^*(\chi_1, \dots, \chi_n, f; t)$ if f is a Laurent diagonal polynomial. We also determine its p -adic Newton polygon.

1. INTRODUCTION

Let $\mathbf{F}_q$ be the finite field of q elements with characteristic p and $\mathbf{F}_{q^m}$ its extension of degree m . Fix a nontrivial additive character Ψ and let $\chi_1, \dots, \chi_n$ be multiplicative characters of $\mathbf{F}_q$. If $f(x_1, \dots, x_n) \in \mathbf{F}_q[x_1, x_1^{-1}, \dots, x_n, x_n^{-1}]$, then one can form the twisted exponential sum

$$(1.1) \quad \begin{aligned} & S_m^*(\chi_1, \dots, \chi_n, f) \\ &= \sum_{(x_1, \dots, x_n) \in (\mathbf{F}_{q^m}^*)^n} \chi_1(N_m(x_1)) \dots \chi_n(N_m(x_n)) \Psi(\text{Tr}_m(f(x_1, \dots, x_n))), \end{aligned}$$

where $\text{Tr}_m : \mathbf{F}_{q^m} \rightarrow \mathbf{F}_q$ is the trace map and $N_m : \mathbf{F}_{q^m} \rightarrow \mathbf{F}_q$ is the norm map. The corresponding L -function is defined as follows:

$$L^*(\chi_1, \dots, \chi_n, f; t) = \exp\left(\sum_{m=0}^{\infty} S_m^*(\chi_1, \dots, \chi_n, f) \frac{t^m}{m}\right).$$

Received by the editors May 1, 2006 and, in revised form, July 20, 2006.

2000 *Mathematics Subject Classification*. Primary 11L03, 11T23, 14G10.

Key words and phrases. Twisted exponential sum, L -function, p -adic Newton polygon, Γ -function, Gross–Koblitz formula, Hasse–Davenport relation.

The research of this author was supported by New Century Excellent Talents in University Grant # NCET-060785, by SRF for ROCS, SEM and by NNSF of China Grant # 10101015.

©2007 American Mathematical Society
 Reverts to public domain 28 years from publication

By the renowned Dwork–Grothendieck theorem, one knows that the L -function $L^*(\chi_1, \dots, \chi_n, f; t)$ is a rational function.

Finding sharp p -adic estimates of the exponential sums $S_m^*(\chi_1, \dots, \chi_n, f)$ is a fundamental question in number theory. This is well known to be reduced to determining the p -adic absolute values of the reciprocal roots and reciprocal poles. The best way to describe this problem is in terms of p -adic Newton polygons (see [11]). Of course one may consider the question of finding the p -adic Newton polygon of Dwork’s unit root L -function attached to a family of algebraic varieties over $\mathbf{F}_q$, which is proved by Wan [18, 19, 20] to be p -adic meromorphic as conjectured by Dwork [4]. Such question should be very deep and hard.

On the other hand, Dwork [5] determined the Newton polygon for the L -functions of the one-dimensional Kloosterman sum, and then Sperber [13] generalized Dwork’s result to the n -dimensional case while Sperber [14] considered the general hyper-Kloosterman sums. Wan [16, 17, 21] solved a conjecture of Adolphson and Sperber [1]. Note also that Wan [17] presented a p -adic proof of the Dwork–Mazur conjecture [3, 11]. When all the χ_i are trivial, then the p -adic Newton polygon of the L -function of the one-variable polynomial $f(x)$ received attention by Sperber [15], Hong [7, 8, 9] and Yang [22]. A general interesting question is to determine the p -adic Newton polygon of the L -function $L^*(\chi, f; t)$ if f is of one variable, even for small degree, and χ is a nontrivial multiplicative character.

In this paper, we are interested in the case that $f(x_1, \dots, x_n)$ is a Laurent diagonal polynomial. We obtain an explicit formula for the L -function $L^*(\chi_1, \dots, \chi_n, f; t)$ if f is a Laurent diagonal polynomial by using the p -adic gamma function. We also determine the p -adic Newton polygon for such an L -function $L^*(\chi_1, \dots, \chi_n, f; t)$. In the forthcoming works, we will use the results presented here to investigate a conjecture of Adolphson and Sperber [2] on the p -adic Newton polygons of twisted exponential sums.

2. PRELIMINARIES ON p -ADIC GAMMA FUNCTIONS

In this section, we present some properties of the p -adic gamma function Γ_p . For $s \in \mathbf{Z}^+$ one sets

$$(2.1) \quad \Gamma(s) = (-1)^s \cdot \prod_{i=1}^{s-1} i.$$

It is known (see, for example, [12]) that this extends to a continuous function on $\mathbf{Z}_p$, and we define Γ_p to be this extension. Therefore it follows from continuity that the following functional equation holds:

$$\frac{\Gamma_p(x+1)}{\Gamma_p(x)} = \begin{cases} -1, & \text{if } x \in p\mathbf{Z}_p, \\ -x, & \text{if } x \notin p\mathbf{Z}_p, \end{cases}$$

and also we have

$$\Gamma_p(x)\Gamma_p(1-x) = (-1)^{\langle x \rangle_p},$$

where $\langle x \rangle_p$ denotes the least nonnegative residue of x modulo p .

Denote by ζ_p a fixed primitive p -th root of unity in $\mathbf{C}_p$, the completion of the algebraic closure of the p -adic field $\mathbf{Q}_p$. Then there is a generator π of the maximal ideal of the ring of integers of $\mathbf{Q}_p(\zeta_p)$ uniquely characterized by $\pi^{p-1} = -p$ and $\pi \equiv \zeta_p - 1 \pmod{(\zeta_p - 1)^2}$. Let ϖ be any fixed Teichmüller character of $\mathbf{F}_q$. Let $\sigma_p(a)$ denote the sum of the digits of the p -adic representation of an integer $a \geq 1$.

Lemma 2.1 (Gross–Koblitz formula, see [6] or [12]). *Let $1 \leq a < p^e := q$. Then the value of the Gauss sum $g_q(a)$ is explicitly given by*

$$g_q(a) := - \sum_{0 \neq x \in \mathbf{F}_q} \omega(x)^{-a} \Psi(\text{Tr}(x)) = \pi^{\sigma_p(a)} \prod_{j=0}^{e-1} \Gamma_p \left(\frac{\langle p^j a \rangle_{q-1}}{q-1} \right).$$

3. TWISTED DIAGONAL EXPONENTIAL SUMS

For brevity, we define $x^\alpha := x_1^{\alpha_1} \dots x_n^{\alpha_n}$ for any vector $x = (x_1, \dots, x_n) \in (\mathbf{F}_{q^m}^*)^n$ and any n -tuple $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbf{Z}^n$. Since ζ_p is a given primitive p th root of unity in $\mathbf{C}_p$, we know that for any given nontrivial additive character Ψ of $\mathbf{F}_q$, we have $\Psi(c) = \zeta_p^{\text{Tr}_{\mathbf{F}_q/\mathbf{F}_p}(c)}$, $c \in \mathbf{F}_q$. Let ϖ_l be any fixed Teichmüller character of $\mathbf{F}_{q^l}$ for any given integer $l \geq 1$. Then $\varpi_1 = \varpi$ and for multiplicative characters $\chi_1, \dots, \chi_n$, there must be nonnegative integers $b_1, \dots, b_n$, such that $\chi_i = \varpi^{b_i}$, $1 \leq i \leq n$. Thus we can rewrite (1.1) as follows:

$$(3.1) \quad S_m^*(\chi_1, \dots, \chi_n, f) = \sum_{x \in (\mathbf{F}_{q^m}^*)^n} \zeta_p^{\text{Tr}_{\mathbf{F}_{q^m}/\mathbf{F}_p}(f(x))} \cdot \prod_{i=1}^n (\varpi^{b_i} \circ \mathbf{N}_m)(x_i).$$

Lemma 3.1. *For any $c \in \mathbf{F}_q$ and any integer $l \geq 1$, we have $\varpi_l(c) = \varpi(c)$.*

Proof. For any positive integer c , let $\mathbf{Z}_c^{\text{unram}}$ denote the ring of integers of the unramified extension of $\mathbf{Q}_p$ of degree c . Since the diagram

$$\begin{array}{ccc} \mathbf{F}_q & \hookrightarrow & \mathbf{F}_{q^l} \\ \downarrow & & \downarrow \\ \mathbf{Z}_{da}^{\text{unram}} & \hookrightarrow & \mathbf{Z}_{dla}^{\text{unram}} \end{array}$$

commutes, the result follows immediately. The proof is complete. $\square$

Now for $x_i \in \mathbf{F}_{q^m}^*$, we have

$$(\varpi^{b_i} \circ \mathbf{N}_m)(x_i) = \varpi^{b_i} \left(x_i^{\frac{q^m-1}{q-1}} \right) = \varpi \left(x_i^{\frac{q^m-1}{q-1} \cdot b_i} \right).$$

Since $x_i \in \mathbf{F}_{q^m}^*$, we have $x_i^{\frac{q^m-1}{q-1} \cdot b_i} \in \mathbf{F}_q$. By Lemma 3.1 we obtain

$$(\varpi^{b_i} \circ \mathbf{N}_m)(x_i) = \varpi_m \left(x_i^{\frac{q^m-1}{q-1} \cdot b_i} \right).$$

So by (3.1) our sum (1.1) may be rewritten as

$$(3.2) \quad S_m^*(\chi_1, \dots, \chi_n, f) = \sum_{x \in (\mathbf{F}_{q^m}^*)^n} \zeta_p^{\text{Tr}_{\mathbf{F}_{q^m}/\mathbf{F}_p}(f(x))} \cdot \varpi_m \left(x^{\frac{q^m-1}{q-1} \cdot b} \right),$$

where $b = (b_1, \dots, b_n)$. From the definition of Gauss sums over $\mathbf{F}_{q^m}$, one can deduce that for each $c \in \mathbf{F}_{q^m}$, the Gauss sum satisfies the following interpolation relation:

$$(3.3) \quad \zeta_p^{\text{Tr}_{\mathbf{F}_{q^m}/\mathbf{F}_p}(c)} = \frac{1}{1-q^m} \sum_{k=0}^{q^m-2} g_{q^m}(k) \cdot \varpi_m(c)^k.$$

Now let $f(x) = f(x_1, \dots, x_n) = \sum_{i=1}^n a_i x^{V_i} \in \mathbf{F}_q[x_1, x_1^{-1}, \dots, x_n, x_n^{-1}]$ be a diagonal polynomial. Then by (3.2) and (3.3) we derive that

$$\begin{aligned}
S_m^*(\chi_1, \dots, \chi_n, f) &= \sum_{x \in (\mathbf{F}_{q^m}^*)^n} \zeta_p^{\text{Tr}_{\mathbf{F}_{q^m}/\mathbf{F}_p}(\sum_{i=1}^n a_i x^{V_i})} \cdot \varpi_m \left(x^{\frac{q^m-1}{q-1} \cdot b} \right) \\
&= \sum_{x \in (\mathbf{F}_{q^m}^*)^n} \varpi_m \left(x^{\frac{q^m-1}{q-1} \cdot b} \right) \cdot \prod_{i=1}^n \zeta_p^{\text{Tr}_{\mathbf{F}_{q^m}/\mathbf{F}_p}(a_i x^{V_i})} \\
&= \sum_{x \in (\mathbf{F}_{q^m}^*)^n} \varpi_m \left(x^{\frac{q^m-1}{q-1} \cdot b} \right) \cdot \prod_{i=1}^n \frac{1}{1-q^m} \sum_{k_i=0}^{q^m-2} g_{q^m}(k_i) \cdot \varpi_m(a_i x^{V_i})^{k_i} \\
&= \sum_{x \in (\mathbf{F}_{q^m}^*)^n} \varpi_m \left(x^{\frac{q^m-1}{q-1} \cdot b} \right) \cdot \sum_{k_1=0}^{q^m-2} \dots \sum_{k_n=0}^{q^m-2} \prod_{i=1}^n \frac{g_{q^m}(k_i)}{1-q^m} \cdot \varpi_m(a_i x^{V_i})^{k_i} \\
&= \sum_{k_1=0}^{q^m-2} \dots \sum_{k_n=0}^{q^m-2} \sum_{x \in (\mathbf{F}_{q^m}^*)^n} \varpi_m \left(x^{\frac{q^m-1}{q-1} \cdot b} \right) \prod_{i=1}^n \frac{g_{q^m}(k_i)}{1-q^m} \cdot \varpi_m(a_i x^{V_i})^{k_i} \\
&= \sum_{k_1=0}^{q^m-2} \dots \sum_{k_n=0}^{q^m-2} \left(\prod_{i=1}^n \frac{g_{q^m}(k_i)}{1-q^m} \cdot \varpi_m(a_i)^{k_i} \right) \cdot \sum_{x \in (\mathbf{F}_{q^m}^*)^n} \varpi_m \left(x^{\sum_{i=1}^n k_i V_i + \frac{q^m-1}{q-1} \cdot b} \right).
\end{aligned}$$

Noting that

$$\sum_{x \in (\mathbf{F}_{q^m}^*)^n} \varpi_m \left(x^{\sum_{i=1}^n k_i V_i + \frac{q^m-1}{q-1} \cdot b} \right)$$

is equal to $(q^m - 1)^n$ if $\sum_{i=1}^n k_i V_i + \frac{q^m-1}{q-1} \cdot b \equiv 0 \pmod{q^m - 1}$, and 0 otherwise, one then deduces that

$$\begin{aligned}
S_m^*(\chi_1, \dots, \chi_n, f) &= \sum_{\substack{\sum_{i=1}^n k_i V_i + \frac{q^m-1}{q-1} \cdot b \equiv 0 \pmod{q^m-1} \\ 0 \leq k_i \leq q^m-2 \text{ for } 1 \leq i \leq n}} (q^m - 1)^n \prod_{i=1}^n \frac{g_{q^m}(k_i)}{1-q^m} \cdot \varpi_m(a_i)^{k_i} \\
(3.4) \quad &= (-1)^n \cdot \sum_{\substack{\sum_{i=1}^n k_i V_i + \frac{q^m-1}{q-1} \cdot b \equiv 0 \pmod{q^m-1} \\ 0 \leq k_i \leq q^m-2 \text{ for } 1 \leq i \leq n}} \prod_{i=1}^n g_{q^m}(k_i) \cdot \varpi_m(a_i)^{k_i}.
\end{aligned}$$

Let each V_i be written as a column vector, and let $M = (V_1, \dots, V_n)$ be the $n \times n$ matrix having V_i as its i -th column for $1 \leq i \leq n$. Write $u_i = \frac{k_i}{q^m-1}$ and $b'_i = \frac{b_i}{q-1}$ for $1 \leq i \leq n$. Then for all $1 \leq i \leq n$, we have $0 \leq u_i, b'_i < 1$. It is clear that

$$\sum_{i=1}^n k_i V_i + \frac{q^m-1}{q-1} \cdot b \equiv 0 \pmod{q^m-1}$$

if and only if

$$Mu + b' \equiv 0 \pmod{1},$$

where $u = (k'_1, \dots, k'_n)^T$ and $b' = (b'_1, \dots, b'_n)^T$ are written as column vectors. Then from (3.4) we finish the proof of the following result.

Theorem 3.2. *For any integer $m \geq 1$, we have*

$$S_m^*(\chi_1, \dots, \chi_n, f) = (-1)^n \cdot \sum_{\substack{Mu+b' \equiv 0 \pmod{1} \\ (q^m-1)u \in \mathbf{Z}^n}} \prod_{i=1}^n g_{q^m}(u_i(q^m-1)) \cdot \varpi_m(a_i)^{u_i(q^m-1)}.$$

4. EXPLICIT FORMULAS OF L -FUNCTIONS OF TWISTED DIAGONAL SUMS
AND NEWTON POLYGONS

In this section, we will give an explicit formula for the L -function of the twisted sum of the diagonal Laurent polynomial over $\mathbf{F}_q$. Let $f(x) = \sum_{i=1}^n a_i x^{V_i}$ be a diagonal Laurent polynomial and Δ the Newton polyhedron generated by the n indices $V_i, 1 \leq i \leq n$. Let $S(\Delta, b')$ be the set of solutions u of the following linear system of congruences:

$$Mu + b' \equiv 0 \pmod{1}, \quad u_i \text{ rational numbers, } 0 \leq u_i < 1.$$

Now suppose that $(\det M, p) = 1$. Then for each $u \in S(\Delta, b')$, there exists a smallest positive integer, denoted by $d(q, b', u)$, such that the multiplication by $q^{d(q, b', u)}$ acts trivially on $u + b'$. We claim that this is equivalent to $(q^{d(q, b', u)} - 1)u \in \mathbf{Z}^n$. In fact, since $(q - 1)b' = b \in \mathbf{Z}^n$, we have $q^{d(q, b', u)}(u + b') \equiv u + b' \pmod{1} \Leftrightarrow (q^{d(q, b', u)} - 1)u \equiv (1 - q^{d(q, b', u)})b' \pmod{1} \Leftrightarrow (q^{d(q, b', u)} - 1)u \equiv 0 \pmod{1}$. So the assertion is proved. Therefore $d(q, b', u)$ is the smallest positive integer d' such that $(q^{d'} - 1)u \in \mathbf{Z}^n$.

For any given integer $d \in \mathbf{Z}^+$, let $S(q, b', d) = \{u \in S(\Delta, b') \mid d(q, b', u) = d\}$. Define a set I of indices by $I := \{d \in \mathbf{Z}^+ \mid \exists u \in S(\Delta, b'), \text{ such that } d(q, b', u) = d\}$. Since $S(\Delta, b')$ is finite, so is I .

To give the main result, one first needs two lemmas.

Lemma 4.1. *For any integer l and $0 \leq k \leq q^d - 2$, we have*

$$g_{q^{dl}} \left(k \cdot \frac{q^{dl} - 1}{q^d - 1} \right) = g_{q^d}(k)^l.$$

Proof. First we have

$$g_{q^{dl}} \left(k \cdot \frac{q^{dl} - 1}{q^d - 1} \right) = - \sum_{c \in \mathbf{F}_{q^{dl}}^*} \varpi_{dl}(N_{dl}(c))^{-k} \cdot \zeta_p^{\text{Tr}_{\mathbf{F}_{q^{dl}}/\mathbf{F}_p}(c)}.$$

Since $N_{dl}(c) \in \mathbf{F}_q$, by Lemma 3.1 we have $\varpi_{dl}(N_{dl}(c)) = \varpi_d(N_{dl}(c))$. Thus

$$g_{q^{dl}} \left(k \cdot \frac{q^{dl} - 1}{q^d - 1} \right) = - \sum_{c \in \mathbf{F}_{q^{dl}}^*} \varpi_d^{-k}(N_{dl}(c)) \cdot \zeta_p^{\text{Tr}_{\mathbf{F}_{q^{dl}}/\mathbf{F}_p}(c)}.$$

Then it follows from the well-known Hasse–Davenport relation (see, for example, [10]) that

$$g_{q^{dl}} \left(k \cdot \frac{q^{dl} - 1}{q^d - 1} \right) = \left(- \sum_{c \in \mathbf{F}_{q^d}^*} \varpi_d^{-k}(N_d(c)) \cdot \zeta_p^{\text{Tr}_{\mathbf{F}_{q^d}/\mathbf{F}_p}(c)} \right)^l = (g_{q^d}(k))^l = g_{q^d}(k)^l,$$

as desired. The proof is complete. $\square$

Lemma 4.2. *The Gauss sum is invariant under p -action; i.e., we have $g_q(pk) = g_q(k)$.*

Proof. We have

$$g_q(pk) = - \sum_{c \in \mathbf{F}_q^*} \varpi(c)^{-pk} \cdot \zeta_p^{\text{Tr}_{\mathbf{F}_q/\mathbf{F}_p}(c)} = - \sum_{c \in \mathbf{F}_q^*} \varpi(c^p)^{-k} \cdot \zeta_p^{\text{Tr}_{\mathbf{F}_q/\mathbf{F}_p}(c)}.$$

Noting that $\text{Tr}_{\mathbf{F}_q/\mathbf{F}_p}(c) = \text{Tr}_{\mathbf{F}_q/\mathbf{F}_p}(c^p)$, we then have

$$g_q(pk) = - \sum_{c \in \mathbf{F}_q^*} \varpi(c^p)^{-k} \cdot \zeta_p^{\text{Tr}_{\mathbf{F}_q/\mathbf{F}_p}(c^p)}.$$

Let $c' = c^p$. It is easy to see that if c runs over $\mathbf{F}_q^*$, then c' runs also over $\mathbf{F}_q^*$. Therefore

$$g_q(pk) = - \sum_{c \in \mathbf{F}_q^*} \varpi(c')^{-k} \cdot \zeta_p^{\text{Tr}_{\mathbf{F}_q/\mathbf{F}_p}(c')} = g_q(k).$$

This completes the proof of Lemma 4.2. $\square$

As usual, let $\{r\}$ denote the fractional part of the real number r , namely $\{r\} = r - [r]$. For any $u \in S(q, b', d)$, one can easily check that $\{u\} \in S(q, b', d)$. So we can define the q -action on u by $q \cdot u := \{qu\}$, where $\{qu\} = (\{qu_1\}, \dots, \{qu_n\})$. Evidently for any $0 \leq i \neq j \leq d-1$, we have $\{q^i u\} \neq \{q^j u\}$ and $\{q^d u\} = u$. Namely $\{\{q^l u\} | l \in \mathbf{Z}_{\geq 0}\} = \{u, \{qu\}, \dots, \{q^{d-1}u\}\}$. Call it an orbit (closed point) of u under q -action. Hence there are d distinct points in the orbit of $u \in S(q, b', d)$ under q -action. Then $S(q, b', d)$ can be partitioned into distinct orbits. We take an element from each orbit of $S(q, b', d)$ as the representation of this orbit and put these representations together to form a set $O(S(q, b', d))$.

We are now ready to give the main result in this paper.

Theorem 4.3. *Let $f(x) = \sum_{i=1}^n a_i x^{V_i} \in \mathbf{F}_q[x_1, x_1^{-1}, \dots, x_n, x_n^{-1}]$ be a diagonal Laurent polynomial, and let $M = (V_1, \dots, V_n)$ be the $n \times n$ matrix having V_i as its i -th column for $1 \leq i \leq n$. Let $(p, \det M) = 1$ and $q = p^e$. Then we have*

$$\begin{aligned} & L^*(\chi_1, \dots, \chi_n, f; t) \\ &= \prod_{d \in I} \prod_{u \in O(S(q, b', d))} \left(1 - t^d \cdot \pi^{\sum_{i=1}^n \sigma_p(u_i(q^d-1))} \right. \\ & \quad \cdot \prod_{i=1}^n \varpi(a_i)^{u_i(q^d-1)} \prod_{j=0}^{de-1} \Gamma_p \left(\frac{\langle p^j u_i(q^d-1) \rangle_{q^d-1}}{q^d-1} \right) \Big). \end{aligned} \quad (4.1)$$

Furthermore, if $u \in O(S(q, b', d))$ and ω_u is any one of the d reciprocal roots of the corresponding factor in (4.1), then we have

$$\text{ord}_q \omega_u = \frac{1}{d} \sum_{j=0}^{de-1} \sum_{i=1}^n \{p^j u_i\}.$$

Proof. From Theorem 3.2 we deduce that

$$\begin{aligned} & L^*(\chi_1, \dots, \chi_n, f; t) \\ &= \exp \left(\sum_{m=1}^{\infty} (-1)^n \cdot \sum_{\substack{u \in S(\Delta, b') \\ (q^m-1)u \in \mathbf{Z}^n}} \prod_{i=1}^n \varpi_m(a_i)^{u_i(q^m-1)} \cdot g_{q^m}(u_i(q^m-1)) \cdot \frac{t^m}{m} \right). \end{aligned} \quad (4.2)$$

Clearly, for any given u satisfying $Mu + b' \equiv 0 \pmod{1}$, there exists a unique integer d such that $u \in S(q, b', d)$, i.e., such that $(q^d-1)u + b' \in \mathbf{Z}^n$, but $(q^{d-1}-1)u + b' \notin \mathbf{Z}^n$.

For such a u , we have $(q^{dk} - 1)u + b' \in \mathbf{Z}^n$ for any integer $k \geq 1$. So by (4.2) we have

$$(4.3) \quad L^*(\chi_1, \dots, \chi_n, f; t) = \exp \left((-1)^n \cdot \sum_{d=1}^{\infty} \sum_{u \in S(q, d, b')} \prod_{k=1}^{\infty} \prod_{i=1}^n \varpi(a_i)^{u_i(q^{dk}-1)} \cdot g_{q^{dk}}(u_i(q^{dk}-1)) \cdot \frac{t^{dk}}{dk} \right).$$

By Lemma 4.1, one knows that for any $u \in S(q, b', d)$,

$$(4.4) \quad g_{q^{dk}}(u_i(q^{dk}-1)) = g_{q^d}(u_i(q^d-1))^k.$$

Note also that for any $u \in S(q, b', d)$, we have $\varpi(a_i)^{u_i(q^{dk}-1)} = \varpi(a_i)^{u_i(q^d-1)k}$. Then by (4.3) and (4.4) we have

$$(4.5) \quad \begin{aligned} & L^*(\chi_1, \dots, \chi_n, f; t) \\ &= \exp \left((-1)^n \cdot \sum_{d=1}^{\infty} \sum_{u \in S(q, b', d)} \sum_{k=1}^{\infty} \frac{t^{dk}}{dk} \cdot \prod_{i=1}^n \varpi(a_i)^{u_i(q^d-1)k} \cdot g_{q^d}(u_i(q^d-1))^k \right) \\ &= \exp \left(\sum_{d=1}^{\infty} \frac{(-1)^{n-1}}{d} \sum_{u \in S(q, b', d)} \sum_{k=1}^{\infty} - \frac{\prod_{i=1}^n \varpi(a_i)^{u_i(q^d-1)k} \cdot g_{q^d}(u_i(q^d-1))^k \cdot t^{dk}}{k} \right) \\ &= \exp \left(\sum_{d=1}^{\infty} \frac{(-1)^{n-1}}{d} \sum_{u \in S(q, b', d)} \log \left(1 - t^d \prod_{i=1}^n \varpi(a_i)^{u_i(q^d-1)} \cdot g_{q^d}(u_i(q^d-1)) \right) \right) \\ &= \prod_{d=1}^{\infty} \prod_{u \in S(q, b', d)} \exp \left(\frac{(-1)^{n-1}}{d} \log \left(1 - t^d \prod_{i=1}^n \varpi(a_i)^{u_i(q^d-1)} \cdot g_{q^d}(u_i(q^d-1)) \right) \right) \\ &= \prod_{d=1}^{\infty} \prod_{u \in S(q, b', d)} \left(1 - t^d \prod_{i=1}^n \varpi(a_i)^{u_i(q^d-1)} \cdot g_{q^d}(u_i(q^d-1)) \right)^{\frac{(-1)^{n-1}}{d}}. \end{aligned}$$

It follows immediately from (4.5) that

$$(4.6) \quad \begin{aligned} & L^*(\chi_1, \dots, \chi_n, f; t)^{(-1)^{n-1}} \\ &= \prod_{d=1}^{\infty} \prod_{u \in S(q, b', d)} \left(1 - t^d \prod_{i=1}^n \varpi(a_i)^{u_i(q^d-1)} \cdot g_{q^d}(u_i(q^d-1)) \right)^{\frac{1}{d}} \\ &= \prod_{d \in I} \prod_{u \in S(q, b', d)} \left(1 - t^d \prod_{i=1}^n \varpi(a_i)^{u_i(q^d-1)} \cdot g_{q^d}(u_i(q^d-1)) \right)^{\frac{1}{d}}. \end{aligned}$$

Now let $u \in O(S(q, b', d))$. Then note that $\omega(a_i)^q = \omega(a_i)$ and by Lemma 4.2 we know that for each of the d points in the orbit of $u \in S(q, b', d)$ under q -action, the corresponding factor in (4.6) is the same. So we can remove the power $\frac{1}{d}$ in (4.6) by restricting u to run over the set $O(S(q, b', d))$. Therefore we get

$$(4.7) \quad \begin{aligned} & L^*(\chi_1, \dots, \chi_n, f; t)^{(-1)^{n-1}} \\ &= \prod_{d \in I} \prod_{u \in O(S(q, b', d))} \left(1 - t^d \prod_{i=1}^n \varpi(a_i)^{u_i(q^d-1)} \cdot g_{q^d}(u_i(q^d-1)) \right). \end{aligned}$$

But Lemma 2.1 applied to the finite field $\mathbf{F}_{q^d}$ gives

$$(4.8) \quad g_{q^d}(u_i(q^d - 1)) = \pi^{\sum_{i=1}^n \sigma_p(u_i(q^d - 1))} \prod_{j=0}^{de-1} \Gamma_p \left(\frac{\langle p^j u_i(q^d - 1) \rangle_{q^d - 1}}{q^d - 1} \right).$$

So by (4.7) and (4.8) we then get the desired formula (4.1).

Now from (4.1) we deduce that

$$(4.9) \quad \begin{aligned} & \text{ord}_q \omega_u \\ &= \frac{1}{d} \cdot \text{ord}_q \left(\pi^{\sum_{i=1}^n \sigma_p(u_i(q^d - 1))} \cdot \prod_{i=1}^n \varpi(a_i)^{u_i(q^d - 1)} \prod_{j=0}^{de-1} \Gamma_p \left(\frac{\langle p^j u_i(q^d - 1) \rangle_{q^d - 1}}{q^d - 1} \right) \right) \\ &= \frac{1}{d} \cdot \text{ord}_q \left(\pi^{\sum_{i=1}^n \sigma_p(u_i(q^d - 1))} \right) \\ &= \frac{1}{d(p-1)} \sum_{i=1}^n \sigma_p(u_i(q^d - 1)) \\ &= \frac{1}{d(p-1)} \sum_{i=1}^n \frac{p-1}{q^d - 1} \sum_{j=0}^{de-1} \langle p^j u_i(q^d - 1) \rangle_{q^d - 1} \\ &= \frac{1}{d} \sum_{i=1}^n \frac{1}{q^d - 1} \sum_{j=0}^{de-1} \langle p^j u_i(q^d - 1) \rangle_{q^d - 1}. \end{aligned}$$

Since

$$\langle p^j u_i(q^d - 1) \rangle_{q^d - 1} - \{p^j u_i\}(q^d - 1) \equiv [p^j u_i](q^d - 1) \equiv 0 \pmod{q^d - 1}$$

and note that $0 \leq \{p^j u_i\}(q^d - 1) \leq q^d - 1$, we have

$$(4.10) \quad \langle p^j u_i(q^d - 1) \rangle_{q^d - 1} = \{p^j u_i\}(q^d - 1).$$

So we get by (4.9) as well (4.10),

$$\text{ord}_q \omega_u = \frac{1}{d} \sum_{i=1}^n \frac{1}{q^d - 1} \sum_{j=0}^{de-1} \{p^j u_i\}(q^d - 1) = \frac{1}{d} \sum_{j=0}^{de-1} \sum_{i=1}^n \{p^j u_i\},$$

as required. The proof of Theorem 4.3 is complete. $\square$

For any rational number $0 \leq r \leq n$, we define a number h_r by

$$h_r := \# \left\{ u \in S(\Delta, b') \mid \sum_{i=1}^n \sum_{j=0}^{de-1} \{p^j u_i\} = dr \right\}.$$

Therefore by Theorem 4.3 we derive the following result.

Theorem 4.4. *Let $f(x) = \sum_{i=1}^n a_i x^{V_i} \in \mathbf{F}_q[x_1, x_1^{-1}, \dots, x_n, x_n^{-1}]$ be a diagonal Laurent polynomial, and let $M = (V_1, \dots, V_n)$ be the $n \times n$ matrix having V_i as its i -th column for $1 \leq i \leq n$ satisfying $(p, \det M) = 1$. Then for each rational number $0 \leq r \leq n$, the p -adic Newton polygon of the L -function $L^*(\chi_1, \dots, \chi_n, f; t)$ has a side of slope r with horizontal length h_r .*

Corollary 4.5. *Let $f(x) = \sum_{i=1}^n a_i x^{V_i} \in \mathbf{F}_q[x_1, x_1^{-1}, \dots, x_n, x_n^{-1}]$ be a diagonal Laurent polynomial, and let $M = (V_1, \dots, V_n)$ be the $n \times n$ matrix having V_i as its i -th column for $1 \leq i \leq n$. Let $(p, \det M) = 1$. Then each of the following is true:*

(i) We have

$$(4.11) \quad L^*(f; t) = \prod_{d \in I_0} \prod_{u \in O(S(q, d))} \left(1 - t^d \cdot \pi^{\sum_{i=1}^n \sigma_p(u_i(q^d - 1))} \cdot \prod_{i=1}^n \varpi(a_i)^{u_i(q^d - 1)} \prod_{j=0}^{de-1} \Gamma_p \left(\frac{\langle p^j u_i(q^d - 1) \rangle_{q^d - 1}}{q^d - 1} \right) \right),$$

where I_0 is defined by $I_0 := \{d \in \mathbf{Z}^+ | \exists u \in S(\Delta, 0), \text{ such that } d(q, 0, u) = d\}$ and $S(q, d) := S(q, 0, d)$.

(ii) If $u \in O(S(q, d))$ and ω_u is any one of the d reciprocal roots of the corresponding factor in (4.11), then we have

$$\text{ord}_q \omega_u = \frac{1}{d} \sum_{j=0}^{de-1} \sum_{i=1}^n \{p^j u_i\}.$$

(iii) For each rational number $0 \leq r \leq n$, the p -adic Newton polygon of the L -function $L^*(f; t)$ has a side of slope r with horizontal length h_r .

Finally we point out that if f is degenerate, which means that $p | \det M$, then we can change it to the nondegenerate case by using some suitable transformations. For instance, let $f(x_1, x_2) = x_1 x_2^{-1} + x_1^{-1} x_2^{1-p}$. Then $|\det(M)| = p$ and so f is degenerate. First let $y_1 = x_1 x_2^{-1}, y_2 = x_2$. Then it is an invertible transformation. Second let $y'_1 = y_1, y'_2 = y_2^p$. Finally let $z_1 = y'_1, z_2 = y'_2 y'_1{}^{-1}$. This is also invertible. Hence we get $f(x_1, x_2) = z_1 + z_2 := h(z_1, z_2)$ and

$$(4.12) \quad L^*(\chi_1, \chi_2, f; t) = L^*(\chi'_1, \chi'_2, h; t)$$

for some multiplicative characters χ'_1, χ'_2 of $\mathbf{F}_q$. But $h = z_1 + z_2$ is nondegenerate. Therefore we can use Theorems 4.3 and 4.4 to compute the L -function $L^*(\chi'_1, \chi'_2, h; t)$ (hence $L^*(\chi_1, \chi_2, f; t)$ by (4.12)) and determine its p -adic Newton polygon.

REFERENCES

1. A. Adolphson and S. Sperber, *Exponential sums and Newton polyhedra: Cohomology and estimates*, Ann. of Math.(2) 130 (1989), 367-406. MR1014928 (91e:11094)
2. A. Adolphson and S. Sperber, *p-Adic estimates for exponential sums*, Lectures Notes in Math. 1454 (1990), 367-406. MR1094845 (92d:11086)
3. B. Dwork, *On the zeta function of a hypersurface*, Publ. Math. I.H.E.S. 12 (1962), 5-68. MR0159823 (28:3039)
4. B. Dwork, *Normalized period matrices II*, Ann. of Math.(2) 98 (1973), 1-57. MR0396580 (53:442b)
5. B. Dwork, *Bessel functions as p-adic functions of the argument*, Duke Math. J. 41 (1974), 711-738. MR0387281 (52:8124)
6. B.H. Gross and N. Koblitz, *Gauss sums and the p-adic Γ -function*, Ann. of Math.(2) 109 (1979), 569-581. MR534763 (80g:12015)
7. S. Hong, *Newton polygons of L-functions associated with exponential sums of polynomials of degree four over finite fields*, Finite Fields Appl. 7 (2001), 205-237. MR1803945 (2001j:11120)
8. S. Hong, *Newton polygons for L-functions of exponential sums of polynomials of degree six over finite fields*, J. Number Theory 97 (2002), 368-396. MR1942966 (2003i:11116)
9. S. Hong, *Newton polygons for L-functions of exponential sums of polynomials of degree five over finite fields*, preprint.

10. K. Ireland and M. Rosen, *A classical introduction to modern number theory*, GTM 84, Springer-Verlag, New York, 2nd edition, 1990. MR1070716 (92e:11001)
11. B. Mazur, *Frobenius and the Hodge filtration*, Bull. A.M.S. 78 (1972), 635-667. MR0330169 (48:8507)
12. A.M. Robert, *A course in p -adic analysis*, GTM 198, Springer-Verlag, New York, 2000. MR1760253 (2001g:11182)
13. S. Sperber, *Congruence properties of the hyper-Kloosterman sum*, Compositio Math. 40 (1980), 3-33. MR558257 (81j:10059)
14. S. Sperber, *Newton polygons for general hyper-Kloosterman sums*, Astérisque 119-120 (1984), 267-330. MR773095 (86i:11044)
15. S. Sperber, *On the p -adic theory of exponential sums*, Amer. J. Math. 108 (1986), 255-296. MR833359 (87j:11055)
16. D. Wan, *Newton polygons and congruence decompositions of L -functions over finite fields*, Contemp. Math. 133 (1992), 221-241. MR1183982 (93i:11099)
17. D. Wan, *Newton polygons of zeta functions and L -functions*, Ann. of Math.(2) 137(1993), 249-293. MR1207208 (94f:11074)
18. D. Wan, *Dwork's conjecture on unit root zeta functions*, Ann. of Math.(2) 150 (1999), 867-927. MR1740990 (2001a:11108)
19. D. Wan, *Higher rank case of Dwork's conjecture*, J. Amer. Math. Soc. 13 (2000), 807-852. MR1775738 (2001f:11101a)
20. D. Wan, *Rank one case of Dwork's conjecture*, J. Amer. Math. Soc. 13 (2000), 853-908. MR1775761 (2001f:11101b)
21. D. Wan, *Variation of p -adic Newton polygons for L -functions of exponential sums*, Asian J. Math. 8 (2004), 427-472. MR2129244 (2006b:11095)
22. R. Yang, *Newton polygons of L -functions of polynomials of the form $x^d + \lambda x$* , Finite Fields Appl. 9 (2003), 59-88. MR1954784 (2003i:11180)

MATHEMATICAL COLLEGE, SICHUAN UNIVERSITY, CHENGDU 610064, PEOPLE'S REPUBLIC OF CHINA

E-mail address: s-f.hong@tom.com

E-mail address: hongsf02@yahoo.com