ON THE NUMBER OF CERTAIN GALOIS EXTENSIONS OF LOCAL FIELDS

DA-SHENG WEI AND CHUN-GANG JI

(Communicated by Ken Ono)

ABSTRACT. In this paper, we will calculate the number of Galois extensions of local fields with Galois group A_n or S_n .

1. INTRODUCTION

Let p be a prime, F a finite extension of p -adic field $\mathbb{Q}_p$ with $[F : \mathbb{Q}_p] = m$. Let k be its residue field with $[k : \mathbb{Z}/p\mathbb{Z}] = f$. It's easy to see that $f|m$. Let π be a uniformizer of F and e the absolute ramification index of $F/\mathbb{Q}_p$. Then $m = ef$. In this paper let μ_l denote the set of l -th roots of unity. All notations are standard if not explained.

Since the number of the extensions of local fields with a given degree inside the fixed algebraic closure is finite, see [2], one can ask for a formula that gives the number of extensions of a given degree. Krasner [1] gave such a formula, and Serre [6] also computed the number of extensions using a different method. Pauli and Roblot [3] gave the third proof for that formula. Similarly one can also ask for a formula that gives the number of Galois extensions of a given degree. In particular, it is possible to ask for a formula that gives the number of the Galois extensions with the prescribed finite Galois group G . We denote this number by $\nu(F, G)$. If G is a p -group with $\mu_p \not\subset F$, Šafarevič [4] gave an explicit formula for the number of the G -extensions over F :

$$\nu(F, G) = \frac{1}{|Aut(G)|} \left(\frac{|G|}{p^d} \right)^{m+1} \prod_{i=1}^{d-1} (p^{m+1} - p^i),$$

where d is the minimal number of generators of G . If G is a p -group, and $\mu_p \subset F$, Yamagishi [7] obtained a formula for $\nu(F, G)$.

In this paper, we will calculate the number of S_n -extensions and A_n -extensions over F , where S_n is the n -th symmetric group and A_n is the n -th alternating group.

The cases for $n \geq 5$ that are quickly dismissed as S_n and A_n are not solvable in these cases, and the Galois groups of extensions of local fields are always solvable. So we only need to handle the remaining cases, $n \leq 4$.

Received by the editors June 13, 2006.

2000 *Mathematics Subject Classification*. Primary 11S15, 11S20.

Key words and phrases. Local fields, Galois closure, ramified extensions.

This work was partially supported by grants #10171046 and #10201013 from NNSF of China and Jiangsu planned projects for postdoctoral research funds.

Theorem 1.1. *Let F be a finite extension over $\mathbb{Q}_p$ with $[F : \mathbb{Q}_p] = m$, $\nu(F, G)$ the number of the Galois extensions K/F with $\text{Gal}(K/F) = G$.*

(1) *Suppose the prime $p \neq 3$; then*

$$\nu(F, S_3) = \begin{cases} 0 & \text{if } \mu_3 \subset F, \\ 3 & \text{if } \mu_3 \not\subset F. \end{cases}$$

(2) *Suppose $p = 3$; then*

$$\nu(F, S_3) = \begin{cases} 3^{m+1} - 3 & \text{if } \mu_3 \subset F, \\ 3^m + \frac{3^{m+1}}{2} - \frac{3}{2} & \text{if } \mu_3 \not\subset F. \end{cases}$$

Theorem 1.2. *Let F be a finite extension over $\mathbb{Q}_p$ with $[F : \mathbb{Q}_p] = m$, $\nu(F, G)$ the number of the Galois extensions K/F with $\text{Gal}(K/F) = G$.*

(1) *Suppose the prime $p \geq 3$; then*

$$\nu(F, S_4) = \nu(F, A_4) = 0.$$

(2) *Suppose $p = 2$; then*

$$\nu(F, A_4) = \begin{cases} 4(2^{2m} - 1)/3 & \text{if } \mu_3 \subset F, \\ (2^{2m} - 1)/3 & \text{if } \mu_3 \not\subset F; \end{cases}$$

$$\nu(F, S_4) = \begin{cases} 0 & \text{if } \mu_3 \subset F, \\ 2^{2m+1} - 1 & \text{if } \mu_4 \not\subset F \text{ and } m \text{ is even and } f = 1, \\ 2^{2m} - 1 & \text{otherwise.} \end{cases}$$

2. SOME LEMMAS

The number of S_2 -extensions and A_3 -extensions of local fields is specified by well-known results of local class field theory. So we only need to calculate the number of Galois extensions over F with Galois group S_3 , S_4 and A_4 . The following lemma plays an important role in our calculation.

Lemma 1. *Let K be a Galois extension over F with the Galois group G . For any subgroup A of G , let F_A be the field fixed by A . Then the Galois closure $\text{cl}(F_A)$ of F_A is a subfield of K and $\text{Gal}(K/\text{cl}(F_A)) = \bigcap_{g \in G} gAg^{-1}$.*

We can get some Galois extensions from some non-Galois extensions by taking their Galois closure. For example, if $G = S_3$, D_8 , A_4 and S_4 , we can choose A to be isomorphic to $\mathbb{Z}/2\mathbb{Z}$, $\mathbb{Z}/2\mathbb{Z}$, $\mathbb{Z}/3\mathbb{Z}$ and S_3 , which is the non-normal subgroup of G respectively, and where D_8 is the 2-sylow subgroup of S_4 . By the above lemma, the Galois extensions of F with Galois group S_3 can be gotten by the Galois closure of extensions of degree 3 of F , and the Galois extensions of F with Galois group D_8 , A_4 and S_4 can be gotten by the Galois closure of extensions of degree 4 of F .

Let $M(n)$ denote the set of all extensions of degree n of F . Let $Ab(n)$ denote the set of abelian extensions of degree n of F . Also, let $M(G)$ denote the set of Galois extensions of F with the Galois group G . Let K be the Galois closure of an extension of degree n of F . The Galois group $\text{Gal}(K/F)$ is a subgroup of S_n . Obviously the order of $\text{Gal}(K/F)$ must be divided by n . So there are the following two maps:

$$f : M(3) \rightarrow Ab(3) \cup M(S_3)$$

and

$$g: M(4) \rightarrow Ab(4) \cup M(D_8) \cup M(A_4) \cup M(S_4)$$

by

$$L \rightarrow cl(L).$$

The two maps are surjective. Any inverse image L of an element K in $M(G)$ is a subfield of K , and L is not a Galois extension of F if G is not an abelian group. In these cases, the Galois group $\text{Gal}(K/L)$ is not a normal subgroup of G . For $G = S_3, D_8, A_4$ and S_4 , we consider respectively the number of non-normal subgroups isomorphic to $\mathbb{Z}/2\mathbb{Z}, \mathbb{Z}/2\mathbb{Z}, \mathbb{Z}/3\mathbb{Z}$ and S_3 . The subgroups are non-normal except there is an order-2 normal subgroup in D_8 . So the number of inverse images of any element in $M(S_3), M(D_8), M(A_4)$ and $M(S_4)$ are 3, 4, 4, 4 respectively. Let $|S|$ denote cardinality of a finite set S . Let $\nu(F, G)$ denote the number of $M(G)$. So there is the following result.

Lemma 2.

$$\begin{aligned} |M(3)| &= |Ab(3)| + 3\nu(F, S_3), \\ |M(4)| &= |Ab(4)| + 4\nu(F, D_8) + 4\nu(F, A_4) + 4\nu(F, S_4). \end{aligned}$$

3. THE PROOF OF THEOREM 1.1

In the following, we denote $m = [F : \mathbb{Q}_p]$, and e is the absolute ramification index of F and $q = p^f$ is the number of elements of the residue field of F .

Proof. (1) For $p \neq 3$,

$$M(3) = \{K \mid [K : F] = 3, K \text{ is a tamely ramified extension of } F\}.$$

(i) If 3rd roots of unity are contained in F , then $M(3) = Ab(3)$. By Lemma 2

$$\nu(F, S_3) = 0.$$

(ii) If 3rd roots of unity are not contained in F , then $|M(3)| = 10$, $|Ab(3)| = 1$. By Lemma 2

$$\nu(F, S_3) = 3.$$

(2) For $p = 3$, by Krasner's theorem [3],

$$|M(3)| = 3q^e + 6(q-1)\left(\sum_{a=0}^{e-1} q^a\right) + 1 = 9q^e - 5.$$

Suppose $\mu_3 \not\subset F$; then

$$|Ab(3)| = \frac{1}{2}\left(\frac{3}{2}\right)^{m+1}(3^{m+1} - 1) = \frac{3^{m+1} - 1}{2} = \frac{3q^e - 1}{2}.$$

Suppose $\mu_3 \subset F$; then

$$|Ab(3)| = 4.$$

By Lemma 2

$$\nu(F, S_3) = \begin{cases} \frac{5q^e - 3}{2} & \text{if } \mu_3 \not\subset F, \\ 3q^e - 3 & \text{if } \mu_3 \subset F. \end{cases}$$

□

4. THE PROOF OF THEOREM 1.2

First we give some propositions.

Proposition 4.1. *Let the prime $p \geq 3$. Then*

$$\nu(F, S_4) = \nu(F, A_4) = 0.$$

Proof. Suppose K is a Galois extension over F with Galois group S_4 . There must exist intermediate fields F^{tr} and F^{ur} such that $\text{Gal}(K/F^{tr})$ is a p -group, and $\text{Gal}(F^{tr}/F^{ur})$ and $\text{Gal}(F^{ur}/F)$ are cyclic groups. By Galois theory, there is a p -group S' which is a normal subgroup of S_4 . Since $p \geq 3$, S' must be (1). Since S_4 does not have a cyclic normal subgroup S such that S_4/S is also cyclic, this is a contradiction.

Similarly we get $\nu(F, A_4) = 0$. $\square$

Proposition 4.2. *Let $p = 2$. Then*

$$\nu(F, A_4) = \begin{cases} 4(2^{2m} - 1)/3 & \text{if } \mu_3 \subset F, \\ (2^{2m} - 1)/3 & \text{if } \mu_3 \not\subset F. \end{cases}$$

Proof. Let K be an A_4 -extension over F . Since K_4 is a normal subgroup of A_4 , there exists a (unique) Galois subfield F' of degree 3 over F , where $K_4 \cong \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$. By [2],

$$\begin{aligned} |F'^*/(F'^*)^2| &= 4q^{3e}, \\ |F^*/(F^*)^2| &= 4q^e. \end{aligned}$$

It is clear that the natural map of $F^*/(F^*)^2 \rightarrow F'^*/(F'^*)^2$ is an injection since $[F' : F] = 3$. We consider the action on $F'^*/(F'^*)^2$ of the Galois group $\text{Gal}(F'/F)$.

(1) Denote $G' = \text{Gal}(F'/F)$. Then the following result holds:

$$(F'^*/(F'^*)^2)^{G'} \cong F^*/(F^*)^2.$$

As we've already noted injectivity, it remains to show that the natural map is surjective. Let $a \in F'^*/(F'^*)^2$ be a fixed point of $\text{Gal}(F'/F)$ and $a \neq 1$ in $F'^*/(F'^*)^2$. Then $F'(\sqrt{a'})$ is a Galois extension over F , where a' represents a lifting of a in F'^* . There isn't an order-2 normal subgroup in S_3 , so

$$\text{Gal}(F'(\sqrt{a'})/F) \cong \mathbb{Z}/6\mathbb{Z}.$$

Let F'' be the fixed field of the normal subgroup $\mathbb{Z}/3\mathbb{Z}$. There exists an element $b \in F^*/(F^*)^2$ such that

$$F'' = F(\sqrt{b}).$$

Then

$$F'(\sqrt{a'}) = F'(\sqrt{b}).$$

So $a = b$ in $F'^*/(F'^*)^2$.

(2) Let σ be a generator of G' . Assume $x \in F'^*/(F'^*)^2 - F^*/(F^*)^2$; then there are the following two cases:

$$(i) \ N_{F'/F}(x) = 1 \text{ in } F^*/(F^*)^2,$$

$$(ii) \ N_{F'/F}(x) \neq 1 \text{ in } F^*/(F^*)^2.$$

In (i), the field $F'(\sqrt{x}, \sqrt{\sigma x})$ is an A_4 -extension over F since the Galois group of $F'(\sqrt{x}, \sqrt{\sigma x})/F$ is isomorphic to $K_4 \rtimes \mathbb{Z}/3\mathbb{Z}$, where $K_4 \cong \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$.

In (ii),

$$F'(\sqrt{x}, \sqrt{\sigma x}, \sqrt{\sigma^2 x}) = F'(\sqrt{x/\sigma(x)}, \sqrt{\sigma x/\sigma^2 x}, \sqrt{N_{F'/F}(x)}),$$

So $F'(\sqrt{x}, \sqrt{\sigma x}, \sqrt{\sigma^2 x})$ is an $(A_4 \times \mathbb{Z}/2\mathbb{Z})$ -extension over F . Any $(A_4 \times \mathbb{Z}/2\mathbb{Z})$ -extension K is generated by an A_4 -extension over F and an extension of degree 2 over F . Denote the unique extension of degree 3 by F' ; then there exist $x \in F'^*/(F'^*)^2$ and $a \in F^*/(F^*)^2$ satisfying $x \notin F^*/(F^*)^2$, $N_{F'/F}(x) \in (F^*)^2$ and $a \notin (F^*)^2$, such that $K = F'(\sqrt{x}, \sqrt{\sigma x}, \sqrt{a})$. It is easy to see that

$$K = F'(\sqrt{ax/\sigma x}, \sqrt{a\sigma x/\sigma^2 x}, \sqrt{a\sigma^2 x/x})$$

since $N_{F'/F}(x) = x\sigma x\sigma^2 x = 1$ in $F^*/(F^*)^2$. Then $\sigma x/\sigma^2 x = x\sigma^2 x/\sigma^2 x = x$ in $F'^*/(F'^*)^2$. Consider K as an extension over F' ; there exist 7 subfields with order 2 over F' which are one-to-one correspondents to $\{y, \sigma y, \sigma^2 y, y\sigma y, \sigma y\sigma^2 y, y\sigma^2 y, N_{F'/F}(y)\}$, where $y = ax/\sigma x$. And $N_{F'/F}(y) = a \neq 1$ in $F^*/(F^*)^2$. The $\text{Gal}(F'/F)$ -orbits are $\{y, \sigma y, \sigma^2 y\}$, $\{y\sigma y, \sigma y\sigma^2 y, y\sigma^2 y\}$, and $\{N_{F'/F}(y)\}$. So any $(A_4 \times \mathbb{Z}/2\mathbb{Z})$ -extension over F is in form of (ii) and

$$\nu(F, A_4 \times \mathbb{Z}/2\mathbb{Z}) = \nu(F, A_4)\nu(F, \mathbb{Z}/2\mathbb{Z}).$$

(i) Suppose $\mu_3 \not\subset F$, F' is the unique unramified extension of degree 3 over F , so

$$\begin{aligned} 3\nu(F, A_4) + 3\nu(F, A_4 \times \mathbb{Z}/2\mathbb{Z}) &= 4q^{3e} - 4q^e, \\ \nu(F, A_4 \times \mathbb{Z}/2\mathbb{Z}) &= \nu(F, A_4)(4q^e - 1). \end{aligned}$$

Then

$$\nu(F, A_4) = (q^{2e} - 1)/3.$$

(ii) Suppose $\mu_3 \subset F$, F' is the unique unramified extension and 3 totally ramified extensions of degree 3 over F , so

$$\begin{aligned} 3\nu(F, A_4) + 3\nu(F, A_4 \times \mathbb{Z}/2\mathbb{Z}) &= 4(4q^{3e} - 4q^e), \\ \nu(F, A_4 \times \mathbb{Z}/2\mathbb{Z}) &= \nu(F, A_4)(4q^e - 1). \end{aligned}$$

Then

$$\nu(F, A_4) = 4(q^{2e} - 1)/3.$$

□

Proposition 4.3. *Let $p = 2$. Then*

$$\nu(F, S_4) = \begin{cases} 0 & \text{if } \mu_3 \subset F, \\ 2^{2m+1} - 1 & \text{if } \mu_4 \not\subset F \text{ and } m \text{ is even and } f = 1, \\ 2^{2m} - 1 & \text{otherwise.} \end{cases}$$

Proof. By Krasner's theorem [3],

$$|M(4)| = 16q^{3e} - 4q^{2e} - 5.$$

By the local class field theory and the dual theory of the finite abelian group, the following equation holds:

$$|Ab(4)| = |\{S : S \text{ is the subgroup of order 4 of } F^*/(F^*)^4\}|.$$

Let T_1 be the set consisting of the elements of order ≤ 2 in $F^*/(F^*)^4$, and let T_2 be the set consisting of the elements of order 4 in $F^*/(F^*)^4$. The sequence

$$0 \rightarrow T_1 \rightarrow F^*/(F^*)^4 \rightarrow (F^*)^2/(F^*)^4 \rightarrow 0$$

is exact. The third map is $a \mapsto a^2$. So

$$|T_1| = |F^*/(F^*)^4|/|(F^*)^2/(F^*)^4|.$$

Suppose $\mu_4 \not\subset F$; then

$$\begin{aligned} |T_1| &= 4q^e, \\ |T_2| &= 8q^{2e} - 4q^e. \end{aligned}$$

Then

$$|Ab(4)| = |T_2|/2 + (|T_1| - 1)(|T_1| - 2)/6 = 20q^{2e}/3 - 4q^e + 1/3.$$

Suppose $\mu_4 \subset F$; then

$$\begin{aligned} |T_1| &= 4q^e, \\ |T_2| &= 16q^{2e} - 4q^e. \end{aligned}$$

Then

$$|Ab(4)| = |T_2|/2 + (|T_1| - 1)(|T_1| - 2)/6 = 32q^{2e}/3 - 4q^e + 1/3.$$

Since D_8 is a 2-group, by Theorem 2.2 of [7],

$$\nu(F, D_8) = \begin{cases} q^e(q^e - 1)(4q^e - 1) & \text{if } \mu_4 \subset F \text{ or } \mu_4 \not\subset F \text{ and } m \text{ is even and } f = 1, \\ q^e(2q^e - 1)^2 & \text{otherwise.} \end{cases}$$

By Lemma 2,

(i) If $\mu_4 \subset F$, then

$$\nu(F, S_4) = (|M(4)| - |Ab(4)|)/4 - \nu(F, A_4) - \nu(F, D_8) = 4(q^{2e} - 1)/3 - \nu(F, A_4).$$

(ii) If $\mu_4 \not\subset F$, and m is odd or m is even and $f \geq 2$, then

$$\nu(F, S_4) = (|M(4)| - |Ab(4)|)/4 - \nu(F, A_4) - \nu(F, D_8) = 4(q^{2e} - 1)/3 - \nu(F, A_4).$$

(iii) If $\mu_4 \not\subset F$ and m is even and $f = 1$, then

$$\nu(F, S_4) = (|M(4)| - |Ab(4)|)/4 - \nu(F, A_4) - \nu(F, D_8) = (7q^{2e} - 4)/3 - \nu(F, A_4).$$

By Proposition 4.2, we have

$$\nu(F, S_4) = \begin{cases} 0 & \text{if } \mu_3 \subset F, \\ 2q^{2e} - 1 & \text{if } \mu_4 \not\subset F \text{ and } n \text{ is even and } f = 1, \\ q^{2e} - 1 & \text{otherwise.} \end{cases}$$

□

Remark. Since K_4 is a normal subgroup of S_4 and $S_4/K_4 \cong S_3$, there exists an S_3 -subextension in an S_4 -extension of F by Galois theory. If $\mu_3 \subset F$ and $p \neq 3$, then $\nu(F, S_3) = 0$. So $\nu(F, S_4) = 0$. This gives another proof for a case of $\nu(F, S_4)$.

Using these propositions, the proof of Theorem 1.2 is obtained. This completes the proof of Theorem 1.2.

5. EXAMPLES

Example 5.1. Let $F = \mathbb{Q}_p$. Then

(1)

$$\nu(\mathbb{Q}_p, S_3) = \begin{cases} 6 & \text{if } p = 3, \\ 0 & \text{if } p \equiv 1 \pmod{3}, \\ 3 & \text{if } p \equiv 2 \pmod{3}. \end{cases}$$

(2)

$$\nu(\mathbb{Q}_p, A_4) = \begin{cases} 1 & \text{if } p = 2, \\ 0 & \text{if } p > 2. \end{cases}$$

(3)

$$\nu(\mathbb{Q}_p, S_4) = \begin{cases} 3 & \text{if } p = 2, \\ 0 & \text{if } p > 2. \end{cases}$$

(4)

$$\nu(\mathbb{Q}_p, S_n) = \nu(\mathbb{Q}_p, A_n) = 0 \quad (n \geq 5).$$

REFERENCES

- [1] M. Krasner, Nombre des extensions d'un degré donné d'un corps p -adique, les Tendances Géométriques en algèbre et Théorie des Nombres, ED. CNRS, Paris, 1966, 143-169. MR0225756 (37:1349)
- [2] S. Lang, Algebraic Number Theory, Second edition. GTM, 110. Springer-Verlag, New York, 1994. MR1282723 (95f:11085)
- [3] S. Pauli and X. F. Roblot, On the computation of all extensions of a p -adic field of a given degree, Math. Comp. **70** (2001), no. 236, 1641-1659. MR1836924 (2002e:11166)
- [4] I. R. Šafarevič, On p -extensions, Mat. Sb. **20**(62)(1947), 351-363; English transl., Amer. Math. Soc. Transl. Ser. 2 **4** (1956), 59-72. MR0020546 (8:560e)
- [5] J. -P. Serre, Corps Locaux, Hermann, Paris, 1963. MR0354618 (50:7096)
- [6] J. -P. Serre, Une "formule de masse" pour les extensions totalement ramifiées de degré donné d'un corps local, C. R. Acad. Sci. Paris Sér. A-B **286** (1978), A1031-A1036. MR500361 (80a:12018)
- [7] M. Yamagishi, On the number of Galois p -extensions of a local field, Proc. Amer. Math. Soc. **123** (1995), 2373-2380. MR1264832 (95j:11109)

DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF SCIENCE AND TECHNOLOGY OF CHINA, HEFEI, PEOPLE'S REPUBLIC OF CHINA 230026

E-mail address: dshwei@ustc.edu

DEPARTMENT OF MATHEMATICS, NANJING NORMAL UNIVERSITY, NANJING, PEOPLE'S REPUBLIC OF CHINA 210097

E-mail address: cgji@njnu.edu.cn