ON LITTLEWOOD-PALEY FUNCTIONS

LESLIE C. CHENG

(Communicated by Michael T. Lacey)

ABSTRACT. We prove that, for a compactly supported L^q function Φ with vanishing integral on $\mathbf{R}^n$, the corresponding square function operator S_Φ is bounded on L^p for $|1/p - 1/2| < \min\{(q-1)/2, 1/2\}$.

1. INTRODUCTION

Let $n \geq 1$ and $\mathbf{R}^n$ denote the n -dimensional Euclidean space. For a function $\Phi \in L^1(\mathbf{R}^n)$ which satisfies

$$(1.1) \quad \int_{\mathbf{R}^n} \Phi(x) dx = 0,$$

we define the square function operator S_Φ by

$$(1.2) \quad (S_\Phi f)(x) = \left(\int_0^\infty |\Phi_t * f(x)|^2 \frac{dt}{t} \right)^{1/2},$$

where $\Phi_t(x) = t^{-n}\Phi(x/t)$. The operator S_Φ is often called a square function or a Littlewood-Paley function. Such operators have long played important roles in harmonic analysis. The main problem under investigation concerns the boundedness of these operators on various L^p spaces.

It has been well known that, if the function Φ is sufficiently nice (in terms of decaying and smoothness properties), the corresponding operator S_Φ is bounded on $L^p(\mathbf{R}^n)$ for $1 < p < \infty$. The following result is due to Benedek, Calderón and Panzone ([2]):

Theorem A. *Suppose that Φ satisfies (1.1) and for some positive α ,*

$$(1.3) \quad |\Phi(x)| \leq C(1 + |x|)^{-n-\alpha}, \quad \int_{\mathbf{R}^n} |\Phi(x-y) - \Phi(x)| dx \leq C|y|^\alpha.$$

Then S_Φ is bounded on $L^p(\mathbf{R}^n)$ for $1 < p < \infty$.

Examples of functions satisfying (1.3) include the Schwartz functions, as well as the following which arise from the Poisson kernel on $\mathbf{R}^n$:

$$\Phi_0(x) = \frac{\partial}{\partial t} \left(\frac{t}{(|x|^2 + t^2)^{(n+1)/2}} \right) \Big|_{t=1}$$

Received by the editors June 27, 2006.

2000 *Mathematics Subject Classification.* Primary 42B25.

©2007 American Mathematical Society

and

$$\Phi_j(x) = \frac{\partial}{\partial x_j} \left(\frac{1}{(|x|^2 + 1)^{(n+1)/2}} \right).$$

When Φ is given by

$$(1.4) \quad \Phi(x) = |x|^{-n+1} \Omega(x) \chi_{[0,1]}(|x|),$$

where Ω is homogeneous of degree 0 and has mean value zero on $\mathbf{S}^{n-1}$, then S_Φ becomes the Marcinkiewicz integral operator ([13]). See also [1], [14], [16] and the extensive list of references given in the survey [6].

In [12] S. Sato proved that, among other things, the conclusion of Theorem A is still true if the smoothness condition (1.3) is eliminated (for L^2 this had been known earlier; see [5], [10]).

In this paper we shall study the L^p boundedness of S_Φ without imposing conditions (1.2) or (1.3) on Φ , or the assumption that Φ be given as in (1.4). The following is a known result:

Theorem B. *Suppose that Φ satisfies (1.1) and is compactly supported.*

- (i) *If $\Phi \in L^q(\mathbf{R}^n)$ for some $q \geq 2$, then S_Φ is a bounded operator on $L^p(\mathbf{R}^n, w)$ for $p > q'$ and $w \in A_{p/q'}$.*
- (ii) *If $\Phi \in L^2(\mathbf{R}^n)$, then S_Φ is a bounded operator on $L^p(\mathbf{R}^n)$ for $1 < p < \infty$.*

In the above statement, $L^p(\mathbf{R}^n, w)$ represents the weighted L^p space with weight w (the definition of the weight class A_p can be found in [11], [4] or [14]). When $w \equiv 1$ we write $L^p(\mathbf{R}^n, w)$ as $L^p(\mathbf{R}^n)$.

Part (i) of Theorem B is due to S. Sato (Theorem 3 in [12]). Part (ii) follows from (i) by using duality and interpolation (with $w \equiv 1$).

Theorem B (ii) covers the cases $\Phi \in L^q(\mathbf{R}^n)$, $q > 2$ as well because of the compact support assumption. However, the approach used in [12] does not appear to work when $q < 2$ (see also [9], page 241). The main purpose of the present paper is to establish the following theorem dealing with the case where $\Phi \in L^q(\mathbf{R}^n)$ for $q < 2$.

Theorem C. *Suppose that Φ is a compactly supported function satisfying (1.1). If $\Phi \in L^q(\mathbf{R}^n)$ for some $q \in (1, 2]$, then S_Φ is a bounded operator on $L^p(\mathbf{R}^n)$ for $|1/p - 1/2| < (q - 1)/2$.*

Remarks 1. (i) The range of p given by $|1/p - 1/2| < (q - 1)/2$ is the same as $2/q < p < 2/(2 - q)$, which becomes $(1, \infty)$ when $q = 2$.

(ii) A result relevant to the theorems mentioned above is Theorem 1 in [7]. While in general an L^q function Φ does not satisfy the pointwise decay condition imposed on its Fourier transform in Theorem 1 of [7], a modification of the proof given in [7] can yield the L^p boundedness of S_Φ for $|1/p - 1/2| < (q - 1)/(2q)$ under the conditions in Theorem C. Since $q > 1$, the range of p given by $|1/p - 1/2| < (q - 1)/2$ in our theorem is considerably better.

(iii) While the inequality $|1/p - 1/2| < (q - 1)/2$ gives the full range $1 < p < \infty$ when $q = 2$, it would be an interesting problem to determine whether it also represents the best possible range for p when $q < 2$.

The proof of Theorem C will be given in Section 2. Section 3 contains a result on the boundedness of S_Φ when the compact support condition is replaced by some other conditions.

2. PROOF OF THEOREM C

Lemma 2.1. *Let Ψ be a compactly supported function in $L^q(\mathbf{R}^n)$ for some $q \in (1, 2]$. Then, for every p satisfying $|1/p - 1/2| < (q - 1)/2$, there exists a $C_p > 0$ such that*

$$(2.1) \quad \left\| \left(\int_0^\infty |(\Psi_t * F^t)(x)|^2 \frac{dt}{t} \right)^{1/2} \right\|_{L^p(\mathbf{R}^n)} \leq C_p \left\| \left(\int_0^\infty |F^t(x)|^2 \frac{dt}{t} \right)^{1/2} \right\|_{L^p(\mathbf{R}^n)}$$

holds for every measurable function $F^t(x) = F(t, x)$ on $(0, \infty) \times \mathbf{R}^n$.

Proof. By duality and interpolation we may assume that $p > 2$. We shall also assume that $\text{supp}(\Psi)$ is contained in $B(0, 1)$, where $B(x_0, t) = \{x \in \mathbf{R}^n : |x - x_0| \leq t\}$. Let T be the operator acting on functions defined on $(0, \infty) \times \mathbf{R}^n$ given by

$$(2.2) \quad T(F)(t, x) = (\Psi_t * F^t)(x),$$

where $F^t(y) = F(t, y)$ for $(t, y) \in (0, \infty) \times \mathbf{R}^n$. For $1 \leq p, q < \infty$ we shall use the following notation:

$$(2.3) \quad \|F\|_{L^p(L^q(t^{-1}dt), dx)} = \left(\int_{\mathbf{R}^n} \left(\int_0^\infty |F(t, x)|^q \frac{dt}{t} \right)^{p/q} dx \right)^{1/p}.$$

Thus we have

$$\begin{aligned} \|T(F)\|_{L^1(L^1(t^{-1}dt), dx)} &= \int_{\mathbf{R}^n} \left(\int_0^\infty |(\Psi_t * F^t)(x)| t^{-1} dt \right) dx \\ &= \int_0^\infty \|\Psi_t * F^t\|_{L^1(\mathbf{R}^n)} t^{-1} dt \\ &\leq \|\Psi\|_{L^1(\mathbf{R}^n)} \int_0^\infty \|F^t\|_{L^1(\mathbf{R}^n)} t^{-1} dt \\ (2.4) \quad &= \|\Psi\|_{L^1(\mathbf{R}^n)} \|F\|_{L^1(L^1(t^{-1}dt), dx)}. \end{aligned}$$

It follows from $1/2 - 1/p < (q - 1)/2$ that

$$p'q/2 > 1.$$

Let $r = (p'q/2)'$. By $p > 2$ we have $r > q'$. Thus, for any F that satisfies

$$\|T(F)\|_{L^r(L^{q'}(t^{-1}dt), dx)} < \infty,$$

there exists a function $h \in L^{(r/q')'}(\mathbf{R}^n)$ such that

$$\|h\|_{L^{(r/q')'}(\mathbf{R}^n)} = 1$$

and

$$(2.5) \quad \|T(F)\|_{L^r(L^{q'}(t^{-1}dt), dx)}^{q'} = \int_{\mathbf{R}^n} \left(\int_0^\infty |(\Psi_t * F^t)(x)|^{q'} t^{-1} dt \right) h(x) dx.$$

By Hölder's inequality,

$$\begin{aligned} |\Psi_t * F^t(x)|^{q'} &= t^{-nq'} \left| \int_{\mathbf{R}^n} \Psi\left(\frac{x-y}{t}\right) F(t, y) dy \right|^{q'} \\ (2.6) \quad &\leq \|\Psi\|_{L^q(\mathbf{R}^n)}^{q'} |B(0, t)|^{-1} \int_{\mathbf{R}^n} |F(t, y)|^{q'} \chi_{B(0, t)}(x - y) dy. \end{aligned}$$

Let M denote the Hardy-Littlewood maximal operator on $\mathbf{R}^n$. Then by (2.5) and (2.6),

$$\begin{aligned}
 \|T(F)\|_{L^r(L^{q'}(t^{-1}dt),dx)}^{q'} &\leq \|\Psi\|_{L^q(\mathbf{R}^n)}^{q'} \int_{\mathbf{R}^n} \left(\int_0^\infty |F(t,y)|^{q'} t^{-1} dt \right) (Mh)(y) dy \\
 &\leq \|\Psi\|_{L^q(\mathbf{R}^n)}^{q'} \left\| \left(\int_0^\infty |F(t,y)|^{q'} t^{-1} dt \right) \right\|_{L^{(r/q')'}(\mathbf{R}^n)}^{q'} \|Mh\|_{L^{(r/q')'}(\mathbf{R}^n)} \\
 (2.7) \qquad &\leq C_{r,q} \|\Psi\|_{L^q(\mathbf{R}^n)}^{q'} \|F\|_{L^r(L^{q'}(t^{-1}dt),dx)}^{q'}.
 \end{aligned}$$

Since

$$\begin{aligned}
 \frac{1}{2} &= \frac{\theta}{q'} + \frac{(1-\theta)}{1}, \\
 \frac{1}{p} &= \frac{\theta}{r} + \frac{(1-\theta)}{1}
 \end{aligned}$$

hold with $\theta = q/2$, by interpolating between (2.4) and (2.7) (see, for example, [3]) we obtain

$$\|T(F)\|_{L^p(L^2(t^{-1}dt),dx)} \leq C_p \|F\|_{L^p(L^2(t^{-1}dt),dx)},$$

which proves (2.1) for $|1/p - 1/2| < (q-1)/2$.

For $s > 0$ we let $s^{\pm\alpha} = \min\{s^\alpha, s^{-\alpha}\}$. □

Lemma 2.2. *Let $\Psi \in \mathcal{S}(\mathbf{R}^n)$ and $\Phi \in L^q(\mathbf{R}^n)$ for some $q \in (1, 2]$. Suppose that Φ is compactly supported and $\text{supp}(\widehat{\Psi}) \subset \{1/2 < |\xi| < 2\}$. Then for every p satisfying $|1/p - 1/2| < (q-1)/2$, there exist $C_p, \alpha_p > 0$ such that*

$$(2.8) \qquad \left\| \left(\int_0^\infty |\Phi_t * \Psi_{st} * f|^2 \frac{dt}{t} \right)^{1/2} \right\|_{L^p(\mathbf{R}^n)} \leq C_p (s^{\pm\alpha_p}) \|f\|_{L^p(\mathbf{R}^n)}$$

for all $f \in L^p(\mathbf{R}^n)$ and $s > 0$.

Proof. Let

$$T_s f(x) = \left(\int_0^\infty |\Phi_t * \Psi_{st} * f(x)|^2 \frac{dt}{t} \right)^{1/2}.$$

Then, by Lemma 2.1 and Theorem A, for each p satisfying $|1/p - 1/2| < (q-1)/2$,

$$\begin{aligned}
 \|T_s f\|_{L^p(\mathbf{R}^n)} &\leq C_p \|S_\Psi f\|_{L^p(\mathbf{R}^n)} \\
 (2.9) \qquad &\leq C_p \|f\|_{L^p(\mathbf{R}^n)}.
 \end{aligned}$$

On the other hand, by Plancherel's Theorem,

$$\begin{aligned}
 \|T_s f\|_{L^2(\mathbf{R}^n)}^2 &= \int_0^\infty \int_{\mathbf{R}^n} |\Phi_t * \Psi_{st} * f(x)|^2 dx t^{-1} dt \\
 (2.10) \qquad &= \int_{\mathbf{R}^n} |\hat{f}(\xi)|^2 \left(\int_0^\infty |\widehat{\Phi}(t\xi)|^2 |\widehat{\Psi}(st\xi)|^2 \frac{dt}{t} \right) d\xi.
 \end{aligned}$$

It follows from Lemmas 2, 3 of [12] and (1.1) that

$$(2.11) \qquad \int_{1/2}^2 |\widehat{\Phi}(t\xi)|^2 dt \leq C |\xi|^{\pm 1/(2q')}.$$

Let $\xi' = |\xi|^{-1}\xi$ for $\xi \neq 0$. By (2.10) and (2.11) we have

$$\begin{aligned} \|T_s f\|_{L^2(\mathbf{R}^n)}^2 &\leq \int_{\mathbf{R}^n} |\hat{f}(\xi)|^2 \left(\int_{1/2}^2 |\widehat{\Phi}(ts^{-1}\xi')|^2 \frac{dt}{t} \right) d\xi \\ (2.12) \qquad \qquad \qquad &\leq C(s^{\pm 1/(2q')}) \|f\|_{L^2(\mathbf{R}^n)}^2. \end{aligned}$$

By (2.9), (2.12) and interpolation we conclude that (2.8) holds for $|1/p - 1/2| < (q-1)/2$. $\square$

Proof of Theorem C. It suffices to establish

$$\|S_\Phi f\|_{L^p(\mathbf{R}^n)} \leq C_p \|f\|_{L^p(\mathbf{R}^n)}$$

for $f \in \mathcal{S}(\mathbf{R}^n)$ and $|1/p - 1/2| < (q-1)/2$.

Let $\eta \in C^\infty(\mathbf{R})$ such that $\text{supp}(\eta) \subset (1/4, 4)$ and

$$(2.13) \qquad \qquad \qquad \int_0^\infty \eta(s) \frac{ds}{s} = 2.$$

Define the Schwartz function Ψ on $\mathbf{R}^n$ by

$$\widehat{\Psi}(\xi) = \eta(|\xi|^2)$$

for $\xi \in \mathbf{R}^n$. Then by (2.13) we have

$$\int_0^\infty \widehat{\Psi}(s\xi) \frac{ds}{s} = 1$$

and

$$(2.14) \qquad \qquad \qquad \Phi_t * f = \int_0^\infty (\Phi_t * \Psi_{st} * f) \frac{ds}{s}$$

for all $f \in \mathcal{S}(\mathbf{R}^n)$ and $t > 0$. By Minkowski's inequality we have

$$|S_\Phi f(x)| \leq \int_0^\infty \left(\int_0^\infty |\Phi_t * \Psi_{st} * f(x)|^2 \frac{dt}{t} \right)^{1/2} \frac{ds}{s}.$$

By Lemma 2.2, for every p satisfying $|1/p - 1/2| < (q-1)/2$,

$$\begin{aligned} \|S_\Phi f\|_{L^p(\mathbf{R}^n)} &\leq C_p \|f\|_{L^p(\mathbf{R}^n)} \int_0^\infty s^{\pm \alpha_p} \frac{ds}{s} \\ (2.15) \qquad \qquad \qquad &= 2\alpha_p^{-1} C_p \|f\|_{L^p(\mathbf{R}^n)}. \end{aligned}$$

The proof of Theorem C is now complete. $\square$

3. FURTHER RESULTS

For Φ 's which are not necessarily compactly supported, one can easily deduce the following result from Theorem 2 of [12] by using duality and interpolation:

Theorem 3.1. *Let $\Phi \in L^1(\mathbf{R}^n)$ and satisfy (1.1). Suppose that*

- (i) $\int_{|x| < 1} |\Phi(x)|^{1+\varepsilon} dx + \int_{|x| > 1} |\Phi(x)||x|^\varepsilon dx < \infty$ for some $\varepsilon > 0$;
- (ii) $|\Phi(x)| \leq h(|x|)\Omega(x')$, where

$$(ii.a) \text{ } h \text{ is non-negative, non-increasing and satisfies } \int_0^\infty h(r)r^{n-1}dr < \infty;$$

$$(ii.b) \text{ } \Omega \in L^q(\mathbf{S}^{n-1}) \text{ for some } q \geq 2.$$

Then S_Φ is bounded on $L^p(\mathbf{R}^n)$ for $1 < p < \infty$.

Using the above theorem one can see that the condition (1.4) in Theorem A is redundant, as observed in [12].

Below we shall show that the requirement “ $q \geq 2$ ” in (ii.b) of the above theorem can be lowered to $q > 1$ without affecting the validity of the claim.

Theorem 3.2. *If the condition (ii.b) in Theorem 3.1 is replaced by the weaker condition (ii.b)': $\Omega \in L^q(\mathbf{S}^{n-1})$ for some $q > 1$, while all other conditions in Theorem 3.1 remain unchanged, then S_Φ is bounded on $L^p(\mathbf{R}^n)$ for $1 < p < \infty$.*

Proof. By Lemmas 1–3 in [12], we see that (2.11) still holds (note that $q > 1$ is needed when applying Lemma 2 of [12]). Thus, it suffices to show that (2.1) holds for all $p \in (1, \infty)$.

For each $y' \in \mathbf{S}^{n-1}$ and $x \in \mathbf{R}^n$, let

$$M_{y'}f(x) = \sup_{s>0} \left(\frac{1}{s} \int_0^s |f(x - sy')| ds \right).$$

For $t > 0$ and $x \in \mathbf{R}^n$,

$$\begin{aligned} |\Phi_t * f(x)| &\leq \int_{\mathbf{S}^{n-1}} |\Omega(y')| \left(\int_0^\infty |f(x - tr y')| h(r) r^{n-1} dr \right) d\sigma(y') \\ &\leq \int_{\mathbf{S}^{n-1}} |\Omega(y')| \left(\sum_{j=-\infty}^\infty 2^{n(j+1)} h(2^j) M_{y'}f(x) \right) d\sigma(y') \\ (3.1) \quad &\leq 2^{2n} \left(\int_0^\infty h(r) r^{n-1} dr \right) \int_{\mathbf{S}^{n-1}} |\Omega(y')| M_{y'}f(x) d\sigma(y'). \end{aligned}$$

By (3.1) and the uniform boundedness of the operators $M_{y'}$ on $L^p(\mathbf{R}^n)$ for $1 < p < \infty$, we have

$$(3.2) \quad \left\| \sup_{t>0} |\Phi_t * f| \right\|_{L^p(\mathbf{R}^n)} \leq C_p \|\Omega\|_1 \|f\|_{L^p(\mathbf{R}^n)}$$

for $1 < p < \infty$. It follows from (3.2) and the proof of the lemma on p. 544 of [8] (after some trivial modifications) that (2.1) holds for $1 < p < \infty$. $\square$

REFERENCES

- [1] Al-Salman, A., Al-Qassem, H., Cheng, L. C., Pan, Y., *L^p Bounds for the Function of Marcinkiewicz*, Math. Res. Lett. **9** (2002), 697–700. MR1906071 (2004d:42020)
- [2] Benedek, A., Calderón, A. and Panzone, R., *Convolution operators on Banach space valued functions*, Proc. Natl. Acad. Sci. USA **48** (1962), 356–365. MR0133653 (24:A3479)
- [3] Bergh, J. and Löfström, J., *Interpolation Spaces*, Springer-Verlag, 1976.
- [4] Coifman, R. and Fefferman, C., *Weighted norm inequalities for maximal functions and singular integrals*, Studia Math. **51** (1974), 241–250. MR0358205 (50:10670)
- [5] Coifman, R. and Meyer, Y., *Au delà des opérateurs pseudo-différentiels*, Astérisque **57** (1978). MR518170 (81b:47061)
- [6] Ding, Y., *On Marcinkiewicz integral*, Proc. of the conference “Singular Integrals and Related Topics, III”, Osaka, Japan (2001), 28–38.
- [7] Ding, Y., Fan, D. and Pan, Y., *On Littlewood-Paley functions and singular integrals*, Hokkaido Math. J. **29** (2000), 537–552. MR1795491 (2001m:42038)
- [8] Duoandikoetxea, J. and Rubio de Francia, J. L., *Maximal and singular integral operators via Fourier transform estimates*, Invent. Math. **84** (1986), 541–561. MR837527 (87f:42046)
- [9] Duoandikoetxea, J. and Seijo, E., *Weighted inequalities for rough square functions through extrapolation*, Studia Math. **149** (2002), 239–252. MR1890732 (2003a:42022)
- [10] Journé, J.-L., *Calderón-Zygmund operators, pseudo-differential operators and the Cauchy integral of Calderón*, Lecture Notes in Math. **994** (1983), Springer-Verlag. MR706075 (85i:42021)

- [11] Muckenhoupt, B., *Weighted norm inequalities for the Hardy maximal functions*, Trans. Amer. Math. Soc. **165** (1972), 207–226. MR0293384 (45:2461)
- [12] Sato, S., *Remarks on square functions in the Littlewood-Paley theory*, Bull. Austral. Math. Soc. **58** (1998), 199–221. MR1642027 (99g:42024)
- [13] Stein, E.M., *On the functions of Littlewood-Paley, Lusin and Marcinkiewicz*, Trans. Amer. Math. Soc. **88** (1958), 430–466. MR0112932 (22:3778)
- [14] Stein, E.M., *Harmonic Analysis: Real-variable methods, orthogonality, and oscillatory integrals*, Princeton University Press, 1993. MR1232192 (95c:42002)
- [15] Stein, E.M., *Singular integrals and differentiability properties of functions*, Princeton University Press, 1970. MR0290095 (44:7280)
- [16] Walsh, T., *On the function of Marcinkiewicz*, Studia Math. **44** (1972), 203–217. MR0315368 (47:3917)

DEPARTMENT OF MATHEMATICS, BRYN MAWR COLLEGE, BRYN MAWR, PENNSYLVANIA 19010
E-mail address: lcheng@brynmawr.edu