AN EXAMPLE ON ORDERED BANACH ALGEBRAS

GERD HERZOG AND CHRISTOPH SCHMOEGER

(Communicated by Joseph A. Ball)

ABSTRACT. Let $\mathcal{B}$ be a complex unital Banach algebra. We consider the Banach algebra $\mathcal{A} = \mathcal{B} \times \mathbb{C}$ ordered by the algebra cone $K = \{(a, \xi) \in \mathcal{A} : \|a\| \leq \xi\}$, and investigate the connection between results on ordered Banach algebras and the right bound of the numerical range of elements in $\mathcal{B}$.

1. ORDERED BANACH ALGEBRAS

The aim of this paper is to stress the aspect of the applicability of the ordered Banach algebra theory, within a wider scope of general Banach algebras. To this end we will embed a given (non-ordered) Banach algebra $\mathcal{B}$ into a certain ordered Banach algebra $\mathcal{A}$; see section 2. In particular Theorems 6 and 7 in section 5 are results strictly in terms of the originally given Banach algebra $\mathcal{B}$, while the proofs involve results on ordered Banach algebras applied to $\mathcal{A}$. We start with some notations and results on general ordered Banach algebras.

Let $\mathcal{A}$ be a complex unital Banach algebra with unit $\mathbf{1}$, and assume that $\mathcal{A}$ is ordered by an algebra cone K , that is, K is a closed convex subset of $\mathcal{A}$ with $\lambda K \subseteq K$ ($\lambda \geq 0$), $K \cap (-K) = \{0\}$, $\mathbf{1} \in K$, $K \cdot K \subseteq K$, and $a \leq b : \iff b - a \in K$.

For a general survey on ordered Banach algebras we refer to [4], [9], and the references given there.

The spectrum of $a \in \mathcal{A}$ is denoted by $\sigma(a)$, and $r(a)$ denotes its spectral radius. Moreover let $\tau(a)$ denote the right spectral bound of a , that is,

$$\tau(a) := \max\{\operatorname{Re} \lambda : \lambda \in \sigma(a)\}.$$

Next, we consider the sets

$$Q_+ := \{a \in \mathcal{A} : \exp(ta) \geq 0 \ (t \geq 0)\}, \quad Q_{\pm} = Q_+ \cap (-Q_+).$$

Amongst other things the following properties of these sets are known. Note that $\emptyset \neq W \subseteq \mathcal{A}$ is called a wedge if W is closed, convex and $\lambda W \subseteq W$ ($\lambda \geq 0$).

Theorem 1. *Under the assumptions above*

1. Q_+ is a wedge, and $\overline{K + \mathbb{R}\mathbf{1}} = Q_+$;
2. $Q_{\pm}$ is a closed real subspace of $\mathcal{A}$;
3. $Q_{\pm}$ is a real Lie algebra;
4. $a \in Q_{\pm} \implies a^2 \in Q_+$;

Received by the editors September 22, 2006 and, in revised form, November 6, 2006.

2000 *Mathematics Subject Classification.* Primary 47H05, 47A12, 47B60.

Key words and phrases. Ordered Banach algebras, bounds of numerical range, fractional powers.

©2007 American Mathematical Society
 Reverts to public domain 28 years from publication

5. $a \in Q_+, \tau(a) < 0 \implies a^{-1} \leq 0$;
 6. $a \in Q_+, \tau(a) < 0 \implies a(\lambda \mathbf{1} - a)^{-1} \in Q_+ \ (\lambda \geq 0)$.

Proof. Parts 1, 2, 4 and 5 of Theorem 1 have been proved in [4] for real Banach algebras (where then in part 5 $\tau(a)$ is obtained by complexification), but hold obviously in the complex case by considering $\mathcal{A}$ as a real Banach algebra (by restricting the scalar field to $\mathbb{R}$). Part 3 of Theorem 1 follows easily from the commutator formula for the exponential function; see [5, Lemma I.2.13]. To prove part 6 note that $\tau(a - \lambda \mathbf{1}) < 0 \ (\lambda \geq 0)$ and consider

$$a(\lambda \mathbf{1} - a)^{-1} = (\lambda \mathbf{1} + a - \lambda \mathbf{1})(\lambda \mathbf{1} - a)^{-1} = -\lambda(a - \lambda \mathbf{1})^{-1} - \mathbf{1}.$$

Thus $a(\lambda \mathbf{1} - a)^{-1} \in K + \mathbb{R}\mathbf{1}$ according to part 5; hence it is in Q_+ by 1. $\square$

In particular part 6 of Theorem 1 can be used to find roots of elements of $-Q_+$ in Q_+ by using the formula of Balakrishnan [10, p.260].

Theorem 2. *Let $a \in Q_+$ with $\tau(a) < 0$, and let $\alpha \in (0, 1)$. Then*

$$-(-a)^\alpha := \frac{\sin(\alpha\pi)}{\pi} \int_0^\infty \lambda^{\alpha-1} a(\lambda \mathbf{1} - a)^{-1} d\lambda \in Q_+.$$

Proof. By means of part 6 of Theorem 1 we see that the integrand is in Q_+ for each $\lambda \in (0, \infty)$. Since Q_+ is a wedge we conclude that $-(-a)^\alpha \in Q_+$. $\square$

For our third theorem, which is on majorization, let the unital Banach algebra $\mathcal{A} \times \mathcal{A}$ (endowed with coordinatewise multiplication and the norm $\|(a, b)\| = \max\{\|a\|, \|b\|\}$) be ordered by the algebra cone

$$\tilde{K} = \{(a, b) \in \mathcal{A} \times \mathcal{A} : -b \leq a \leq b\},$$

and let $\tilde{Q}_+$ denote the corresponding set of exponentially nonnegative elements. The following theorem is proved in [3] for operators on ordered Banach spaces and holds with an analog proof for Banach algebras in the following version:

Theorem 3. *Let $a, b \in \mathcal{A}$ be such that $a \leq b$ and $a + b \in Q_+$. Then $(a, b) \in \tilde{Q}_+$.*

2. SPECIAL ORDERED BANACH ALGEBRAS

Now, let $\mathcal{B}$ be any complex unital Banach algebra with unit $\mathbf{1}$ and let $\mathcal{A}$ denote the Banach algebra $\mathcal{B} \times \mathbb{C}$, endowed with multiplication and norm

$$(a, \xi) \cdot (b, \eta) = (ab, \xi\eta), \quad \|(a, \xi)\| = \max\{\|a\|, |\xi|\},$$

respectively. Then $\mathcal{A}$ is a unital Banach algebra with unit $(\mathbf{1}, 1)$, and we consider the ordering defined by the cone

$$K = \{(a, \xi) \in \mathcal{A} : \|a\| \leq \xi\}.$$

We will characterize Q_+ and $Q_\pm$ in this case, and we will obtain results on the right bound of the numerical range of elements of $\mathcal{B}$, by means of Theorem 1.

3. BOUNDS OF THE NUMERICAL RANGE

Let $\mathcal{B}$ be a complex unital Banach algebra and consider the one-sided directional derivatives of the norm at $\mathbf{1}$:

$$n_+[a] = \lim_{h \rightarrow 0+} \frac{\|\mathbf{1} + ha\| - 1}{h}, \quad n_-[a] = \lim_{h \rightarrow 0-} \frac{\|\mathbf{1} + ha\| - 1}{h} \quad (a \in \mathcal{B}).$$

For each $a \in \mathcal{B}$ let $V(a)$ denote the numerical range of a , that is,

$$V(a) = \{\varphi(a) : \varphi \in \mathcal{B}^*, \|\varphi\| = \varphi(\mathbf{1}) = 1\}.$$

The set $V(a)$ is bounded closed and convex, and $\sigma(a) \subseteq V(a)$ for each $a \in \mathcal{B}$; see [2]. The number $n_+[a]$ is the right bound of the numerical range of a , that is,

$$n_+[a] = \max\{\operatorname{Re} \lambda : \lambda \in V(a)\},$$

according to Mazur's representation of directional derivatives of sublinear functionals [8]; see also [2], [7]. In particular $\tau(a) \leq n_+[a]$.

Note the following properties of n_+ and n_- : The function $a \mapsto n_+[a]$ is sublinear, and

$$n_-[a] \leq n_+[a], \quad n_-[-a] = -n_+[a], \quad n_\pm[a + \lambda \mathbf{1}] = n_\pm[a] + \lambda$$

for all $a \in \mathcal{B}$ and $\lambda \in \mathbb{R}$.

Next, let $H(\mathcal{B})$ denote the set of all hermitian elements of $\mathcal{B}$, that is,

$$H(\mathcal{B}) = \{a \in \mathcal{B} : \|\exp(ita)\| = 1 \ (t \in \mathbb{R})\}.$$

For the following properties of hermitian elements see [2, 5. Lemma 2] for part 1 and [1] for part 2.

Proposition 1. *We have*

1. $a \in H(\mathcal{B}) \iff V(a) \subseteq \mathbb{R} \iff n_+[ia] = n_-[ia] = 0;$
2. $a \in H(\mathcal{B}) \iff \sigma(a) \subseteq \mathbb{R} \text{ and } r(\alpha \mathbf{1} + \beta a) = \|\alpha \mathbf{1} + \beta a\| \ (\alpha, \beta \in \mathbb{C}).$

Now, in the sequel let $\mathcal{A} = \mathcal{B} \times \mathbb{C}$ be as in section 2.

4. REPRESENTATION OF Q_+ AND $Q_\pm$

Note that

$$Q_+ = \{(a, \xi) \in \mathcal{A} : \|\exp(ta)\| \leq \exp(t\xi) \ (t \geq 0)\}$$

in our case.

Theorem 4. *We have*

1. $(a, \xi) \in Q_+$ if and only if $n_+[a] \leq \xi$.
2. $(a, \xi) \in Q_\pm$ if and only if one of the following equivalent conditions is valid:
 - (a) $\|\exp(ta)\| = \exp(t\xi) \ (t \in \mathbb{R});$
 - (b) $n_+[a] = n_-[a] = \xi;$
 - (c) $\xi \in \mathbb{R} \text{ and } i(a - \xi \mathbf{1}) \in H(\mathcal{B});$
 - (d) $\operatorname{Re} \lambda = \xi \ (\lambda \in \sigma(a)) \text{ and } r(\alpha \mathbf{1} + \beta a) = \|\alpha \mathbf{1} + \beta a\| \ (\alpha, \beta \in \mathbb{C}).$

Proof. 1. First note that for each $a \in \mathcal{B}$

$$\|\exp(ta)\| \leq \exp(tn_+[a]) \quad (t \geq 0);$$

see [2, 3. Theorem 4]. Thus, $n_+[a] \leq \xi$ implies $(a, \xi) \in Q_+$. On the other hand, $(a, \xi) \in Q_+$ implies

$$\frac{\|\exp(ta)\| - 1}{t} \leq \frac{\exp(t\xi) - 1}{t} \quad (t > 0)$$

and as $t \rightarrow 0+$ we obtain $n_+[a] \leq \xi$, since $\exp(ta) = 1 + ta + O(t^2)$ ($t \rightarrow 0+$).

2. Let $(a, \xi) \in Q_{\pm}$. Then $\|\exp(ta)\| \leq \exp(t\xi)$ ($t \in \mathbb{R}$). From

$$\|\exp(ta)\| \geq \frac{1}{\|(\exp(ta))^{-1}\|} = \frac{1}{\|\exp(-ta)\|} \geq \frac{1}{\exp(-t\xi)} = \exp(t\xi)$$

we obtain $\|\exp(ta)\| = \exp(t\xi)$ ($t \in \mathbb{R}$).

Next, by means of part 1 we have

$$n_-[a] \leq n_+[a] \leq \xi, \quad -n_-[a] = n_+[-a] \leq -\xi,$$

thus

$$\xi \leq n_-[a] \leq n_+[a] \leq \xi \Rightarrow n_-[a] = n_+[a] = \xi.$$

In particular, $\xi \in \mathbb{R}$ and

$$n_{\pm}[a - \xi \mathbf{1}] = n_{\pm}[a] - \xi = 0.$$

Thus

$$\lim_{h \rightarrow 0} \frac{\|1 + h(a - \xi \mathbf{1})\| - 1}{h} = 0,$$

and according to part 1 of Proposition 1, $i(a - \xi \mathbf{1}) \in H(\mathcal{B})$.

Since clearly each element (a, ξ) with the property in (a) or (b) is in $Q_{\pm}$ and since (c) implies (b) according to part 1 of Proposition 1, we have proved that $(a, \xi) \in Q_{\pm}$ if and only if (a, ξ) satisfies (a), (b) or (c), respectively

To prove the equivalence of (c) and (d) first let (a, ξ) be such that $\xi \in \mathbb{R}$ and $b := i(a - \xi \mathbf{1}) \in H(\mathcal{B})$. According to part 2 of Proposition 1

$$\sigma(b) \subseteq \mathbb{R} \Rightarrow \sigma(a) \subseteq \{\xi + it : t \in \mathbb{R}\}.$$

For $\alpha, \beta \in \mathbb{C}$ we obtain

$$r(\alpha \mathbf{1} + \beta a) = r((\alpha + \xi \beta) \mathbf{1} - i\beta b) = \|(\alpha + \xi \beta) \mathbf{1} - i\beta b\| = \|\alpha \mathbf{1} + \beta a\|.$$

Vice versa, let (a, ξ) satisfy (d), and set $b := i(a - \xi \mathbf{1})$. Then $\xi \in \mathbb{R}$, and $\sigma(b) \subseteq \mathbb{R}$.

For $\alpha, \beta \in \mathbb{C}$ we have

$$\alpha \mathbf{1} + \beta b = (\alpha - i\xi \beta) \mathbf{1} + i\beta a,$$

thus

$$r(\alpha \mathbf{1} + \beta b) = \|\alpha \mathbf{1} + \beta a\|.$$

Once more using part 2 of Proposition 1 we conclude $b \in H(\mathcal{B})$. □

5. APPLICATIONS

By application of Theorem 1 to the algebra $\mathcal{A}$ we obtain:

Theorem 5. *Let $a, b \in \mathcal{B}$.*

1. *If $a, b \in H(\mathcal{B})$, then $i(ab - ba) \in H(\mathcal{B})$; compare [2, 5. Lemma 4].*
2. *If $a \in H(\mathcal{B})$, then $n_+[-a^2] \leq 0$.*
3. *$\tau((a, n_+[a])) = n_+[a]$.*
4. *If $n_+[a] < 0$, then $\|a^{-1}\| \leq -(n_+[a])^{-1}$; compare [6, Proposition 1.1(3)].*
5. *If $n_+[a] < 0$, then*

$$n_+[a(\lambda \mathbf{1} - a)^{-1}] \leq \frac{n_+[a]}{\lambda - n_+[a]} \quad (\lambda \geq 0).$$

Proof. 1. According to Theorem 4 we have $(ia, 0), (ib, 0) \in Q_{\pm}$. Hence $(-ab + ba, 0) \in Q_{\pm}$, and again Theorem 4 proves $i(ab - ba) \in H(\mathcal{B})$.

2. Since $(ia, 0) \in Q_{\pm}$ we have $(-a^2, 0) \in Q_{+}$. Hence $n_{+}[-a^2] \leq 0$.

3. If $\operatorname{Re} \lambda > n_{+}[a]$, then $(a, n_{+}[a]) - \lambda(\mathbf{1}, 1)$ is invertible, since $\tau(a) \leq n_{+}[a]$ and $n_{+}[a] - \lambda \neq 0$. Moreover $(a, n_{+}[a]) - n_{+}[a](\mathbf{1}, 1) = (a - n_{+}[a]\mathbf{1}, 0)$ is not invertible. Thus $n_{+}[a] \in \sigma((a, n_{+}[a]))$.

4. We have $(a, n_{+}[a]) \in Q_{+}$, and $\tau((a, n_{+}[a])) = n_{+}[a] < 0$. Thus

$$-(a, n_{+}[a])^{-1} = (-a^{-1}, (-n_{+}[a])^{-1}) \geq 0,$$

which means $\|a^{-1}\| \leq -(n_{+}[a])^{-1}$.

5. Since $(a, n_{+}[a]) \in Q_{+}$ and since $\tau((a, n_{+}[a])) = n_{+}[a] < 0$ we have

$$\begin{aligned} & (a, n_{+}[a])(\lambda(\mathbf{1}, 1) - (a, n_{+}[a]))^{-1} \\ &= (a(\lambda\mathbf{1} - a)^{-1}, n_{+}[a](\lambda - n_{+}[a])^{-1}) \in Q_{+} \end{aligned}$$

for each $\lambda \geq 0$. According to Theorem 4

$$n_{+}[a(\lambda\mathbf{1} - a)^{-1}] \leq \frac{n_{+}[a]}{\lambda - n_{+}[a]} \quad (\lambda \geq 0).$$

□

Analogously we can apply Theorem 2 and derive one sided estimates for fractional powers:

Theorem 6. Let $a \in \mathcal{B}$ with $n_{+}[a] < 0$, and $\alpha \in (0, 1)$. Then

$$n_{+}[(-(-a))^{\alpha}] \leq -(-n_{+}[a])^{\alpha}.$$

Proof. Since $(a, n_{+}[a]) \in Q_{+}$ we have

$$-(-(a, n_{+}[a]))^{\alpha} = (-(-a)^{\alpha}, -(-n_{+}[a])^{\alpha}) \in Q_{+}.$$

□

Remark. For results on the numerical range of roots in Banach algebras see [6] and the references given there.

Next, Theorem 3 leads to the following inequalities for the exponential function in Banach algebras:

Theorem 7. Let $a, b \in \mathcal{B}$. Then

$$\|\exp(b) - \exp(a)\| \leq 2 \sinh\left(\frac{\|b - a\|}{2}\right) \exp\left(\frac{n_{+}[a + b]}{2}\right)$$

and

$$\|\exp(b) + \exp(a)\| \leq 2 \cosh\left(\frac{\|b - a\|}{2}\right) \exp\left(\frac{n_{+}[a + b]}{2}\right).$$

Proof. We set

$$\alpha = \frac{n_{+}[a + b] - \|b - a\|}{2}, \quad \beta = \frac{n_{+}[a + b] + \|b - a\|}{2}.$$

Then

$$\|b - a\| = \beta - \alpha \Rightarrow (a, \alpha) \leq (b, \beta)$$

and

$$n_{+}[a + b] = \alpha + \beta \Rightarrow (a, \alpha) + (b, \beta) \in Q_{+}.$$

According to Theorem 3 we have $((a, \alpha), (b, \beta)) \in \tilde{Q}_+$. Hence

$$-(\exp(tb), \exp(t\beta)) \leq (\exp(ta), \exp(t\alpha)) \leq (\exp(tb), \exp(t\beta)) \quad (t \geq 0).$$

For $t = 1$ we obtain

$$\|\exp(b) - \exp(a)\| \leq \exp(\beta) - \exp(\alpha) = 2 \sinh\left(\frac{\|b - a\|}{2}\right) \exp\left(\frac{n_+[a + b]}{2}\right),$$

and the second inequality in an analog way. $\square$

Remark. The inequalities in Theorem 7 are quite obvious if a and b commute. Then, for example,

$$\begin{aligned} \|\exp(b) - \exp(a)\| &= \left\| 2 \sinh\left(\frac{b - a}{2}\right) \exp\left(\frac{a + b}{2}\right) \right\| \\ &\leq 2 \sinh\left(\frac{\|b - a\|}{2}\right) \left\| \exp\left(\frac{a + b}{2}\right) \right\| \leq 2 \sinh\left(\frac{\|b - a\|}{2}\right) \exp\left(\frac{n_+[a + b]}{2}\right). \end{aligned}$$

REFERENCES

- [1] Baillet, M.: *Sur les éléments hermitiens d'une algèbre de Banach et les dilatations de certains opérateurs normaloïdes*. C. R. Acad. Sci., Paris, Ser. **A** **281**, 1039-1042 (1975). MR0390822 (52:11645)
- [2] Bonsall, F.F.; Duncan, J.: *Numerical ranges of operators on normed spaces and of elements of normed algebras*. London: Cambridge University Press. II (1971). MR0288583 (44:5779)
- [3] Herzog, G.; Kunstmann, P.C.: *Majorization of C_0 -semigroups in ordered Banach spaces*. Comment. Math. Univ. Carolin. **47**, 47-54 (2006). MR2223966
- [4] Herzog, G.; Lemmert, R.: *On quasipositive elements in ordered Banach algebras*. Stud. Math. **129**, 59-65 (1998). MR1611855 (99g:46061)
- [5] Hilgert, J.; Neeb, K.-H.: *Lie-Gruppen und Lie-Algebren*. Braunschweig: Vieweg (1991).
- [6] Li, C.-K.; Rodman, L.; Spitkovsky, I.M.: *On numerical ranges and roots*. J. Math. Anal. Appl. **282**, 329-340 (2003). MR2000347 (2004g:47009)
- [7] Martin, R.H.: *Nonlinear operators and differential equations in Banach spaces*. Pure and Applied Mathematics. New York etc.: John Wiley&Sons XI (1976). MR0492671 (58:11753)
- [8] Mazur, S.: *Über konvexe Mengen in linearen normierten Räumen*. Stud. Math. **4**, 70-84 (1933).
- [9] Raubenheimer, H.; Rode, S.: *Cones in Banach algebras*. Indag. Math., New Ser. **7**, 489-502 (1996). MR1620116 (99i:46035)
- [10] Yosida, K.: *Functional analysis*. Berlin-Göttingen-Heidelberg: Springer-Verlag. XI (1965).

INSTITUT FÜR ANALYSIS, UNIVERSITÄT KARLSRUHE, D-76128 KARLSRUHE, GERMANY
E-mail address: Gerd.Herzog@math.uni-karlsruhe.de

INSTITUT FÜR ANALYSIS, UNIVERSITÄT KARLSRUHE, D-76128 KARLSRUHE, GERMANY
E-mail address: christoph.schmoeger@math.uni-karlsruhe.de