THE GLOBAL ATTRACTIVITY OF THE RATIONAL DIFFERENCE EQUATION $y_n = A + \left(\frac{y_{n-k}}{y_{n-m}}\right)^p$

KENNETH S. BERENHAUT, JOHN D. FOLEY, AND STEVO STEVIĆ

(Communicated by Carmen C. Chicone)

ABSTRACT. This paper studies the behavior of positive solutions of the recursive equation

$$y_n = A + \left(\frac{y_{n-k}}{y_{n-m}}\right)^p, \quad n = 0, 1, 2, \dots,$$

with $y_{-s}, y_{-s+1}, \dots, y_{-1} \in (0, \infty)$ and $k, m \in \{1, 2, 3, 4, \dots\}$, where $s = \max\{k, m\}$. We prove that if $\gcd(k, m) = 1$, and $p \leq \min\{1, (A+1)/2\}$, then y_n tends to $A+1$. This complements several results in the recent literature, including the main result in K. S. Berenhaut, J. D. Foley and S. Stević, The global attractivity of the rational difference equation $y_n = 1 + \frac{y_{n-k}}{y_{n-m}}$, *Proc. Amer. Math. Soc.*, 135 (2007) 1133–1140.

1. INTRODUCTION

This paper studies the behavior of positive solutions of the recursive equation

$$(1) \quad y_n = A + \left(\frac{y_{n-k}}{y_{n-m}}\right)^p, \quad n = 0, 1, \dots,$$

with $y_{-s}, y_{-s+1}, \dots, y_{-1} \in (0, \infty)$ and $k, m \in \{1, 2, 3, 4, \dots\}$, where $s = \max\{k, m\}$.

The case $k = m$ is trivial, so from now on we will assume that $k \neq m$.

Note that if $g = \gcd(k, m) > 1$, then $\{y_i\}$ can be separated into g different equations of the form

$$(2) \quad y_n^{(j)} = A + \left(\frac{y_{n-\frac{k}{g}}^{(j)}}{y_{n-\frac{m}{g}}^{(j)}}\right)^p,$$

where $j \in \{1, 2, \dots, g\}$. Hence, we may assume that $\gcd(k, m) = 1$.

The study of properties of rational and nonlinear difference equations has been an area of intense interest in recent years; cf. [1]–[25] and the references therein.

There is a relatively long history in studying equation (1). For example, for $p = 1$, the case $k = 2, m = 1$ was studied in [2] by Amleh *et al.*, the case $k \in \mathbb{N}, m = 1$ was studied by DeVault *et al.* in [11], and the case $A > 1, k = 1, m \in \mathbb{N}$ was studied by Stević in [20]. The investigation of global stability and periodicity of

Received by the editors April 18, 2006 and in revised form, July 31, 2006.

2000 *Mathematics Subject Classification.* Primary 39A10, 39A11.

Key words and phrases. Rational difference equation, stability.

The first author acknowledges financial support from a Sterge Faculty Fellowship.

©2007 American Mathematical Society
 Reverts to public domain 28 years from publication

positive solutions of equation (1), for the case $p = A = 1$, $k, m \in \mathbb{N}$ was completed by results in [3] and [15]; see also [17] and [21].

The study of the case $p > 1$ was suggested in [14], where the authors noted that some results from [2] for the case $p = 1$, $k = 2$, $m = 1$, can be translated to the case $p > 1$, $k = 2$, $m = 1$. The first results for the case $p < 1$ were given in [23]. The existence of monotone solutions, for the case $p > 0$ and $A > -1$ was shown in [5] by developing the technique from [6, 7, 8, 9, 10, 24] and [25]. Equations in papers [4] and [12] were investigated by transforming them into some special cases of equation (1).

The linearized equation associated with equation (1) for the case $k = 2$ and $m = 1$ is

$$(A + 1)z_n + pz_{n-1} - pz_{n-2} = 0,$$

and its characteristic roots are

$$\lambda_1 = \frac{-p + \sqrt{p^2 + 4p(A+1)}}{2(A+1)} \quad \text{and} \quad \lambda_2 = \frac{-p - \sqrt{p^2 + 4p(A+1)}}{2(A+1)}.$$

By some simple calculation we obtain

$$|\lambda_1| = \frac{2p}{p + \sqrt{p^2 + 4p(A+1)}} < 1,$$

for every $p, A > 0$.

On the other hand, we have that

$$|\lambda_2| < 1 \iff 2p < A + 1.$$

Hence, when $2p < A + 1$ equation (1) for the case $k = 2$ and $m = 1$ is locally asymptotically stable by the Linearized Stability Theorem.

Motivated by this local stability result, in [22] Stević has posed the following conjecture.

Conjecture 1. *If $k = 2$, $m = 1$ and $p, A \in (0, 1)$ are such that $p < (A + 1)/2$, then every positive solution of equation (1) converges to the unique equilibrium $A + 1$.*

Among other results, here we confirm the conjecture by proving that the following holds true for every $k, m \in \mathbb{N}$, $0 < A < 1$ and $0 < p \leq (A + 1)/2$.

Theorem 1. *Suppose that $m, k \geq 1$, and that p, A are positive numbers satisfying $0 < A < 1$ and $0 < p \leq (A + 1)/2$. If the sequence $\{y_i\}$ satisfies (1) with $y_{-s}, y_{-s+1}, \dots, y_{-1} \in (0, \infty)$ where $s = \max\{m, k\}$, then, $\{y_i\}$ converges to the unique equilibrium $A + 1$.*

Remark 1 (The case k even and m odd). Note that the general characteristic equation associated with the linearized equation for equation (1) is

$$(3) \quad (A + 1)\lambda^s + p\lambda^{s-m} - p\lambda^{s-k} = 0,$$

and for k even and m odd, equation (3) has a real root $\lambda_0 < -1$, when $p > (A + 1)/2$. To see this, suppose that $p > (A + 1)/2$, and set

$$(4) \quad f(\lambda) = A + 1 + \frac{p}{\lambda^m} - \frac{p}{\lambda^k}.$$

Now, note that $f(-1) = A + 1 - 2p < 0$, and for $\lambda < -1$,

$$(5) \quad f(\lambda) = A + 1 - \frac{p}{|\lambda|^m} - \frac{p}{|\lambda|^k} > A + 1 - \frac{2p}{|\lambda|^{\min\{m, k\}}} > 0,$$

for sufficiently large $|\lambda|$.

Hence, by the continuity of the function f on the interval $(-\infty, -1)$ it follows that $f(\lambda) = 0$ for some $\lambda \in (-\infty, -1)$, as required.

Thus, by the Linearized Stability Theorem, the positive equilibrium $\bar{y} = A + 1$ of equation (1) is not stable, in this case. This fact in conjunction with Theorem 1 gives a full characterization of stability for the case k even and m odd, for $A, p \in (0, 1)$. $\square$

The paper proceeds as follows. In Section 2, we introduce some preliminary lemmas and notation. Section 3 is devoted to global stability, where among other results we give a proof of Theorem 1.

2. PRELIMINARIES AND NOTATION

In this section, we introduce some preliminary lemmas and notation.

First, consider the simple transformed sequence $\{z_i\}$ defined by $z_n = y_n - A$, for $n \geq -s$. Then, equation (1) becomes

$$(6) \quad z_n = \left(\frac{A + z_{n-k}}{A + z_{n-m}} \right)^p,$$

for $n \geq 0$.

Now, define $\{z_i^*\}$ by

$$(7) \quad z_i^* = \begin{cases} z_i, & \text{if } z_i \geq 1, \\ \frac{1}{z_i}, & \text{otherwise.} \end{cases}$$

The following elementary lemma will be useful.

Lemma 1. *If $x > 1$ and $0 < A < 1$, then*

$$(8) \quad \left(\frac{A + x}{A + 1/x} \right)^{\frac{A+1}{2}} \leq x,$$

with the inequality if and only if $x = 1$, and if $x \geq 1$ and $A > 1$, then the reverse inequality to inequality (8) holds.

Proof. Assume first that $A \in (0, 1)$. Then the inequality in (8) is equivalent to

$$(9) \quad g_A(x) \stackrel{\text{def}}{=} (A + 1) \ln \left(\frac{A + x}{Ax + 1} \right) - (1 - A) \ln x \leq 0.$$

Note that

$$(10) \quad \lim_{x \rightarrow +\infty} g_A(x) = -\infty \text{ and } g_A(1) = 0.$$

By some simple calculations we obtain that

$$(11) \quad g'_A(x) = -\frac{A(x-1)^2(1-A)}{(A+x)(Ax+1)x} < 0,$$

when $x \neq 1$, since $A \in (0, 1)$.

Hence, $g_A(x)$ is decreasing, and thus by (10) is negative on the interval $(1, \infty)$.

Now, assume that $A > 1$. Then $\lim_{x \rightarrow +\infty} g_A(x) = +\infty$ and $g_A(1) = 0$. On the other hand, by (11) it follows that $g'_A(x) > 0$, from which the desired inequality follows. $\square$

Remark 2. Note that if $A = 1$ the inequality in (8) becomes equality.

Next we prove a contraction lemma which will be helpful in showing convergence of solutions in the transformed space obtained through (7).

Lemma 2. *Suppose $\{z_i\}$ satisfies (6) with $p \leq (A+1)/2$ and $A \in (0, 1]$. Then,*

$$(12) \quad 1 \leq z_n^* \leq \max\{z_{n-k}^*, z_{n-m}^*\},$$

for all $n \geq s$.

Proof. Suppose that $z_{n-k} > z_{n-m}$ and set $x = \max\{z_{n-k}^*, z_{n-m}^*\}$. Note that if $z_{n-k} \geq 1$, then $1 \leq z_{n-k} \leq x$ and consequently

$$(13) \quad 1/x \leq z_{n-k} \leq x,$$

and if $z_{n-k} < 1$, then $1/z_{n-k} = z_{n-k}^* \leq x$ from which (13) also holds. Similarly, we have that

$$(14) \quad 1/x \leq z_{n-m} \leq x.$$

Then, from (6), (13) and (14), for $n \geq s$, we have that

$$(15) \quad z_n^* = z_n = \left(\frac{A + z_{n-k}}{A + z_{n-m}} \right)^p \leq \left(\frac{A + z_{n-k}}{A + z_{n-m}} \right)^{\frac{A+1}{2}} \leq \left(\frac{A + x}{A + \frac{1}{x}} \right)^{\frac{A+1}{2}} \leq x,$$

where the final inequality in (15) follows from Lemma 1. Similarly, if $z_{n-k} \leq z_{n-m}$, then

$$(16) \quad z_n^* = \frac{1}{z_n} = \left(\frac{A + z_{n-m}}{A + z_{n-k}} \right)^p \leq \left(\frac{A + z_{n-m}}{A + z_{n-k}} \right)^{\frac{A+1}{2}} \leq \left(\frac{A + x}{A + \frac{1}{x}} \right)^{\frac{A+1}{2}} \leq x,$$

and the lemma is proved. $\square$

Now, set

$$(17) \quad D_n = \max_{n-s \leq i \leq n-1} \{z_i^*\},$$

for $n \geq s$.

The following result is a simple consequence of Lemma 2 and (17).

Lemma 3. *The sequence $\{D_i\}$ is monotonically nonincreasing in i , for $i \geq s$.*

Since $D_i \geq 1$ for $i \geq s$, Lemma 3 implies that, as i tends to infinity, the sequence $\{D_i\}$ converges to some limit, say D , where $D \geq 1$.

Next, we have the following lemma concerning boundedness of solutions to equation (1).

Lemma 4. *If $p \in (0, 1)$, then every positive solution of equation (1) is bounded.*

Proof. First, note that each $n \in \mathbb{N}_0$ can be written in the form $lk + i$, for some $l \in \mathbb{N}_0$ and $i \in \{0, 1, \dots, k-1\}$. Let $l_0 = l_0(i)$ be the smallest element of $\mathbb{N}_0$ such that $l_0k + i \geq m$. From (1) and since $y_n > A$ for every $n \geq 0$, we have that

$$(18) \quad y_{lk+i} = A + \frac{y_{(l-1)k+i}^p}{y_{lk+i-m}^p} < A + \frac{y_{(l-1)k+i}^p}{A^p},$$

for every $l \in \mathbb{N}_0$ and $i \in \{0, 1, \dots, k-1\}$ such that $lk + i \geq m$. Let $(u_l^{(i)})_{l \in \mathbb{N}}$ be the solution of the difference equation

$$u_l^{(i)} = A + \frac{u_{l-1}^{(i)}}{A^p}, \quad u_{l_0}^{(i)} = y_{i(l_0-1)+i}.$$

By (18) and induction we see that $y_{(l-1)k+i} \leq u_l^{(i)}$, $l \geq l_0$. Hence it is enough to prove that the sequences $(u_l^{(i)})_{l \geq l_0}$, $i \in \{0, 1, \dots, k-1\}$, are bounded.

Since the function

$$f(x) = A + \frac{x^p}{A^p}, \quad x \in (0, \infty),$$

is increasing and concave for $p \in (0, 1)$ it follows that there is a unique fixed point $\bar{x}$ of the equation $f(x) = x$ and that the function f satisfies the condition

$$(f(x) - x)(x - \bar{x}) < 0, \quad x \in (0, \infty).$$

Using this fact it is easy to see that if $u_l^{(i)} \in (0, \bar{x}]$ the sequence is nondecreasing and bounded above by $\bar{x}$ and if $u_l^{(i)} \geq \bar{x}$, it is nonincreasing and bounded below by $\bar{x}$. Hence for every $u_{l_0}^{(i)} \in (0, \infty)$, each of the sequences $u_l^{(i)}$, $i \in \{0, 1, \dots, k-1\}$, is bounded, from which the result follows. $\square$

3. CONVERGENCE OF SOLUTIONS TO EQUATION (1)

In this section, we study the global attractivity of the positive solutions of equation (1). First, we give a proof of Theorem 1.

Proof of Theorem 1. Note that it suffices to show that the transformed sequence $\{z_i^*\}$ converges to 1.

By the definition in (17), the values of D_i are taken on by entries in the sequence $\{z_j^*\}$, and as well, by Lemma 2, $z_i^* \in [1, D_i]$ for $i \geq s$. Now, for any $\epsilon > 0$, we can find an N such that

$$(19) \quad z_N^* \in [D, D + \epsilon],$$

and for $i \geq N - s$,

$$(20) \quad z_i^* \in [1, D + \epsilon].$$

Note that, similar to (13) and (14), (20) implies that

$$(21) \quad \frac{1}{D + \epsilon} \leq z_{N-m}, z_{N-k} \leq D + \epsilon.$$

We will show that $D = 1$, and from this, (7), (17) and the definition of D , the result follows.

Now, suppose $D > 1$, and note that (19) implies that $z_N \neq 1$.

First, consider the case $z_N > 1$. Then, from (19), we have that

$$(22) \quad z_N = z_N^* \in [D, D + \epsilon].$$

Solving for z_{n-k} in (6), and employing (22) and (21), gives

$$\begin{aligned} D + \epsilon &\geq z_{N-k} = z_N^{1/p} (A + z_{N-m}) - A \\ &\geq D^{1/p} \left(A + \frac{1}{D + \epsilon} \right) - A \\ (23) \quad &\geq D^{\frac{2}{A+1}} \left(A + \frac{1}{D + \epsilon} \right) - A. \end{aligned}$$

This implies that

$$(24) \quad \left(\frac{A + D + \epsilon}{A + \frac{1}{D + \epsilon}} \right) \geq D^{\frac{2}{A+1}}.$$

Assume now that $z_N < 1$. Then,

$$(25) \quad \frac{1}{z_N} = z_N^* \in [D, D + \epsilon].$$

From (6), and employing (21) and (25), it follows that

$$(26) \quad \begin{aligned} D + \epsilon &\geq z_{N-m} = (z_N^*)^{1/p} (A + z_{N-k}) - A \\ &\geq D^{1/p} \left(A + \frac{1}{D + \epsilon} \right) - A \\ &\geq D^{\frac{2}{A+1}} \left(A + \frac{1}{D + \epsilon} \right) - A. \end{aligned}$$

From (26) we have that (24) holds in this case, as well. Since $\epsilon > 0$ was arbitrary and $D > 1$, by Lemma 1 we arrive at a contradiction, which implies that $D = 1$, and the theorem follows. $\square$

Remark 3. Note that the above argument breaks down when $p > (A + 1)/2$. In particular, we have that for $p > (A + 1)/2$,

$$(27) \quad \left(\frac{A + x}{A + 1/x} \right)^p > x,$$

for $x = 1 + \epsilon$, for sufficiently small $\epsilon > 0$. To see this, similar to (9), set

$$(28) \quad h_{A,p}(x) = p(\ln(A + x) - \ln(Ax + 1)) - (1 - p)\ln x,$$

and note that the condition $h_{A,p}(x) > 0$ is equivalent to (27). Now, $h_{A,p}(1) = 0$ and

$$(29) \quad h'_{A,p}(1) = \frac{2p - (A + 1)}{(A + 1)} > 0.$$

Hence for sufficiently small $\epsilon > 0$, $h_{A,p}(1 + \epsilon) > 0$.

The next theorem is devoted to the case $p \in (0, 1)$ and $A \geq 1$. It is simpler than Theorem 1 and is essentially a consequence of the boundedness result in Lemma 4.

Theorem 2. *Suppose that $m, k \geq 1$, $p \in (0, 1)$ and $A \geq 1$. If the sequence $\{y_i\}$ satisfies (1) with $y_{-s}, y_{-s+1}, \dots, y_{-1} \in (0, \infty)$ where $s = \max\{m, k\}$, then, $\{y_i\}$ converges to the unique equilibrium $A + 1$.*

Proof. By Lemma 4, every solution $\{y_n\}$ of equation (1) is bounded which implies that there are finite $\liminf y_n = \lambda$ and $\limsup y_n = \Lambda$. Assume to the contrary that $\lambda < \Lambda$. Taking the $\liminf$ and $\limsup$ in (1) it follows that

$$A + \frac{\lambda^p}{\Lambda^p} \leq \lambda < \Lambda \leq A + \frac{\Lambda^p}{\lambda^p}.$$

From this and since $p \in (0, 1)$, it follows that

$$A\Lambda^p + \lambda^p \leq \Lambda^p\lambda < \Lambda\lambda^p \leq A\lambda^p + \Lambda^p;$$

i.e.,

$$(A - 1)\Lambda^p < (A - 1)\lambda^p.$$

Since $A \geq 1$, this is impossible. Therefore we have that $\lambda = \Lambda$, which implies the result. $\square$

REFERENCES

- [1] R. M. ABU-SARIS AND R. DEVAULT, Global stability of $y_{n+1} = A + \frac{y_n}{y_{n-k}}$, *Appl. Math. Lett.* **16** (2) (2003), 173-178. MR1962312 (2004c:39037)
- [2] A. M. AMLEH, E. A. GROVE, G. LADAS AND D. A. GEORGIU, On the recursive sequence $y_{n+1} = \alpha + \frac{y_{n-1}}{y_n}$, *J. Math. Anal. Appl.* **233** (1999), 790-798. MR1689579 (2000f:39002)
- [3] K. S. BERENHAUT, J. D. FOLEY AND S. STEVIĆ, The global attractivity of the rational difference equation $y_n = 1 + \frac{y_{n-k}}{y_{n-m}}$, *Proc. Amer. Math. Soc.*, **135** (2007), 1133-1140. MR2262916 (2007f:39006)
- [4] K. S. BERENHAUT AND S. STEVIĆ, A note on the difference equation $x_{n+1} = \frac{1}{x_n x_{n-1}} + \frac{1}{x_{n-3} x_{n-4}}$, *J. Differ. Equations Appl.* **11** (14) (2005), 1225-1228. MR2182249
- [5] K. S. BERENHAUT AND S. STEVIĆ, On Positive Nonoscillatory Solutions of the Difference Equation $x_{n+1} = \alpha + \frac{x_{n-k}^p}{x_n^p}$, *J. Differ. Equations Appl.*, in press (2005).
- [6] L. BERG, Asymptotische Darstellungen und Entwicklungen, Dt. Verlag Wiss., Berlin, 1968. MR0241873 (39:3210)
- [7] L. BERG, On the asymptotics of nonlinear difference equations, *Z. Anal. Anwendungen* **21**, No.4, (2002), 1061-1074. MR1957315 (2004b:39015)
- [8] L. BERG, Inclusion theorems for non-linear difference equations with applications, *J. Differ. Equations Appl.* **10** (4) (2004), 399-408. MR2047219 (2004m:39007)
- [9] L. BERG, Corrections to "Inclusion theorems for non-linear difference equations with applications," from [3], *J. Differ. Equations Appl.* **11** (2) (2005), 181-182. MR2114324 (2005h:39007)
- [10] L. BERG AND L. V. WOLFERSDORF, On a class of generalized autoconvolution equations of the third kind, *Z. Anal. Anwendungen* **24**, No. 2 (2005), 217-250. MR2174021 (2006j:45020)
- [11] R. DEVAULT, C. KENT AND W. KOSMALA, On the recursive sequence $x_{n+1} = p + \frac{x_{n-k}}{x_n}$, *J. Differ. Equations Appl.* **9** (8) (2003), 721-730. MR1992905 (2004f:39014)
- [12] R. DEVAULT, G. LADAS AND S.W. SCHULTZ, On the recursive sequence $x_{n+1} = \frac{A}{x_n x_{n-1}} + \frac{1}{x_{n-3} x_{n-4}}$, *J. Differ. Equation Appl.* **6** (4) (2000), 481-483. MR1785161
- [13] H. EL-METWALLY, E. A. GROVE, G. LADAS, AND H. D. VOULOV, On the global attractivity and the periodic character of some difference equations, *J. Diff. Eqn. Appl.* **7** (2001), 837-850. MR1870725 (2003e:39006)
- [14] H. M. EL-OWAIDY, A. M. AHMED AND M. S. MOUSA, On asymptotic behaviour of the difference equation $x_{n+1} = \alpha + \frac{x_{n-1}^p}{x_n^p}$, *J. Appl. Math. & Computing* **12** (1-2) (2003), 31-37. MR1976801 (2004a:39011)
- [15] E. A. GROVE, AND G. LADAS, *Periodicities in Nonlinear Difference Equations*, Chapman & Hall/CRC Press, Boca Raton, (2004). MR2193366 (2006j:39002)
- [16] V. KOCIĆ AND G. LADAS, *Global behavior of nonlinear difference equations of higher order with applications*, Mathematics and its Applications, 256. Kluwer Academic Publishers Group, Dordrecht, 1993. MR1247956 (94k:39005)
- [17] W. T. PATULA AND H. D. VOULOV, On the oscillation and periodic character of a third order rational difference equation. *Proc. Amer. Math. Soc.* **131** (2003), no. 3, 905-909. MR1937429 (2003j:39008)
- [18] S. STEVIĆ, Asymptotic behaviour of a sequence defined by iteration with applications, *Colloq. Math.* **93** (2) (2002), 267-276. MR1930804 (2003k:39012)
- [19] S. STEVIĆ, Asymptotic behaviour of a nonlinear difference equation, *Indian J. Pure Appl. Math.* **34** (12) (2003), 1681-1687. MR2030114 (2005a:39029)
- [20] S. STEVIĆ, On the recursive sequence $x_{n+1} = \frac{A}{\prod_{i=0}^k x_{n-i}} + \frac{1}{\prod_{j=k+2}^{2(k+1)} x_{n-j}}$, *Taiwanese J. Math.* **7** (2) (2003), 249-259. MR1978014 (2004c:39030)
- [21] S. STEVIĆ, A note on periodic character of a difference equation, *J. Differ. Equations Appl.* **10** (10) (2004), 929-932. MR2079642 (2005b:39011)
- [22] S. STEVIĆ, Some open problems and conjectures on difference equations, <http://www.mi.sanu.ac.yu>, April 29, 2004.
- [23] S. STEVIĆ, On the recursive sequence $x_{n+1} = \alpha + \frac{x_{n-1}^p}{x_n^p}$, *J. Appl. Math & Computing* **18** (1-2) (2005), 229-234. MR2137703 (2005m:39032)

- [24] S. STEVIĆ, Global stability and asymptotics of some classes of rational difference equations, *J. Math. Anal. Appl.* **316** (2006), 60-68. MR2201749 (2006i:39026)
- [25] S. STEVIĆ, On positive solutions of a $(k + 1)$ -th order difference equation, *Appl. Math. Lett.* **19** (5) (2006), 427-431. MR2213143

DEPARTMENT OF MATHEMATICS, WAKE FOREST UNIVERSITY, WINSTON-SALEM, NORTH CAROLINA 27109

E-mail address: `berenhks@wfu.edu`

DEPARTMENT OF MATHEMATICS, WAKE FOREST UNIVERSITY, WINSTON-SALEM, NORTH CAROLINA 27109

E-mail address: `folejd4@wfu.edu`

MATHEMATICAL INSTITUTE OF THE SERBIAN ACADEMY OF SCIENCE, KNEZ MIHAILOVA 35/I 11000 BEOGRAD, SERBIA

E-mail address: `sstevic@ptt.yu`

E-mail address: `sstevo@matf.bg.ac.yu`