SUBMANIFOLDS OF REAL ALGEBRAIC VARIETIES

W. KUCHARZ

(Communicated by Paul Goerss)

ABSTRACT. By the Nash-Tognoli theorem, each compact smooth manifold M is diffeomorphic to a nonsingular real algebraic set, called an algebraic model of M . We construct algebraic models X of M with controlled behavior of the group of cohomology classes represented by algebraic subsets of X .

1. INTRODUCTION

A compact smooth (of class C^∞) manifold is said to be a *boundary* if it is diffeomorphic to the boundary of a compact smooth manifold with boundary. By convention, the empty manifold is a boundary. The goal of the present paper is to demonstrate that boundaries play a surprisingly interesting role in real algebraic geometry (cf. [4, Theorem 1.4] for the first result of this kind).

All real algebraic varieties in this paper are assumed to be affine (that is, isomorphic to an algebraic subset of $\mathbb{R}^n$ for some n). For background material on real algebraic geometry the reader may consult [2]. Every real algebraic variety carries also the Euclidean topology, induced by the usual metric topology on $\mathbb{R}$. Unless explicitly stated otherwise, all topological notions related to real algebraic varieties will refer to the Euclidean topology.

Given a compact nonsingular real algebraic variety X , we denote by $H_d^{alg}(X, \mathbb{Z}/2)$ the subgroup of $H_d(X, \mathbb{Z}/2)$ of homology classes represented by d -dimensional algebraic subsets of X ; cf. [1, 2, 3, 5]. For technical reasons it is more convenient to work with cohomology groups. We set

$$H_{alg}^c(X, \mathbb{Z}/2) := D_X^{-1}(H_d^{alg}(X, \mathbb{Z}/2)),$$

where $c + d = \dim X$ and $D_X : H^c(X, \mathbb{Z}/2) \rightarrow H_d(X, \mathbb{Z}/2)$ is the Poincaré duality isomorphism. The groups $H_{alg}^c(-, \mathbb{Z}/2)$ are of fundamental interest in real algebraic geometry and will also be in the center of our attention here. Their basic properties and applications are surveyed in [3].

Let M be a compact smooth manifold of dimension m . Denote by $[M]$ the fundamental class of M in $H_m(M, \mathbb{Z}/2)$. For any smooth d -dimensional submanifold N of M (submanifolds are always assumed to be closed subsets), we let $[N]_M$ denote the homology class in $H_d(M, \mathbb{Z}/2)$ represented by N . We set $[N]_M = 0$ if N is empty.

Received by the editors September 10, 2005 and, in revised form, September 28, 2006.
 2000 *Mathematics Subject Classification*. Primary 14P05, 14C25.

Given a finite collection $\mathcal{F}$ of smooth submanifolds of M , we define

$$G^k(\mathcal{F}) := \{u \in H^k(M, \mathbb{Z}/2) \mid \langle u, [N]_M \rangle = 0 \text{ for all } N \text{ in } \mathcal{F} \text{ with } \dim N = k\}.$$

Here, as usual, $\langle \cdot, \cdot \rangle$ stands for scalar product (Kronecker index). If there is no k -dimensional submanifold in $\mathcal{F}$, then $G^k(\mathcal{F}) := H^k(M, \mathbb{Z}/2)$. Clearly, $G^k(\mathcal{F})$ is a subgroup of $H^k(M, \mathbb{Z}/2)$.

Recall that $\mathcal{F}$ is said to be in *general position* if for each point x in M , the collection $\mathcal{F}_x = \{N \in \mathcal{F} \mid x \in N\}$ is either empty or otherwise $\text{codim}(\bigcap_{N \in \mathcal{F}_x} T_x N) = \sum_{N \in \mathcal{F}_x} \text{codim} T_x N$, where $T_x N$ is regarded as a subspace of the tangent space $T_x M$. If $\mathcal{F}$ is in general position and $N_1, \dots, N_s$ are in $\mathcal{F}$, then $N_1 \cap \dots \cap N_s$ is a smooth submanifold of M .

Definition 1.1. A finite collection $\mathcal{F}$ of smooth submanifolds of M is said to be admissible if:

- (i) $\dim N < \dim M$ for all N in $\mathcal{F}$;
- (ii) $\mathcal{F}$ is in general position;
- (iii) if $N_1, \dots, N_s$ are in $\mathcal{F}$, then $N_1 \cap \dots \cap N_s$ is in $\mathcal{F}$;
- (iv) each submanifold in $\mathcal{F}$ has trivial normal vector bundle.

If M' is another smooth manifold and if $\varphi : M' \rightarrow M$ is a smooth diffeomorphism, then

$$\varphi^* \mathcal{F} := \{\varphi^{-1}(N) \mid N \in \mathcal{F}\}$$

is a collection of smooth submanifolds of M' . The collection $\varphi^* \mathcal{F}$ is admissible, provided $\mathcal{F}$ is admissible.

Theorem 1.2. *Let X be a compact nonsingular real algebraic variety and let $\mathcal{F}$ be an admissible collection of smooth submanifolds of X . If*

$$H_{alg}^k(X, \mathbb{Z}/2) \subseteq G^k(\mathcal{F})$$

for some nonnegative integer k , then each k -dimensional manifold in $\mathcal{F}$ is a boundary.

Theorem 1.2 is easy to prove. Its purpose is to provide motivation for our main result, Theorem 1.3, below. By Tognoli's theorem [14] (cf. also [2, Theorem 14.1.10] and, for a weaker but influential result, [12]), each compact smooth manifold M is diffeomorphic to a nonsingular real algebraic variety, called an *algebraic model* of M . Constructing algebraic models satisfying some additional desirable conditions is an interesting and active area of research; cf. [1, 2, 3, 8, 9].

Theorem 1.3. *Let M be a compact smooth manifold and let $\mathcal{F}$ be an admissible collection of smooth submanifolds of M . If each manifold in $\mathcal{F}$ is a boundary, then there exist an irreducible algebraic model X of M and a smooth diffeomorphism $\varphi : X \rightarrow M$ such that*

$$H_{alg}^k(X, \mathbb{Z}/2) \subseteq G^k(\varphi^* \mathcal{F})$$

for all nonnegative integers k .

Theorem 1.3 is useful for constructing examples such that $H_{alg}^k(-, \mathbb{Z}/2) \neq H^k(-, \mathbb{Z}/2)$.

Example 1.4. The n -fold product $T^n = S^1 \times \cdots \times S^1$ of the unit circle S^1 has an algebraic model X with $H_{alg}^k(X, \mathbb{Z}/2) \neq H^k(X, \mathbb{Z}/2)$ for $1 \leq k \leq n-1$. Indeed, suppose $n \geq 2$, since for $n = 1$ the assertion is void. Choose $n-1$ distinct points $p_1, \dots, p_{n-1}$ in S^1 and set

$$N_i = \{(x_1, \dots, x_n) \in T^n \mid x_1 = \cdots = x_i = p_i\}$$

for $1 \leq i \leq n-1$. Then $\mathcal{F} := \{N_1, \dots, N_{n-1}\}$ is an admissible collection of smooth submanifolds of T^n . Moreover, $\text{codim}_{T^n} N = i$ and N_i is a boundary for $1 \leq i \leq n-1$. Hence the assertion follows from Theorem 1.3.

Theorems 1.2 and 1.3 will be proved in the next section.

2. PROOFS

As usual, the i th Stiefel-Whitney class of a smooth manifold M will be denoted by $w_i(M)$. Given a smooth submanifold N of M , we set

$$[N]^M := D_M^{-1}([N]_M).$$

Henceforth, both the homology class $[N]_M$ and cohomology class $[N]^M$ will be used. Basic properties of the products $\langle \cdot, \cdot \rangle, \cap, \cup$ familiar from algebraic topology [6] will be used without further explanation.

Lemma 2.1. *Let M be a compact smooth manifold and let N be a k -dimensional smooth submanifold of M . If the normal vector bundle of N in M is trivial, then the following conditions are equivalent:*

- (a) N is a boundary,
- (b) $\langle w_{i_1}(M) \cup \cdots \cup w_{i_r}(M), [N]_M \rangle = 0$ for all nonnegative integers $i_1, \dots, i_r$ with $i_1 + \cdots + i_r = k$,
- (c) $\langle w_{i_1}(M) \cup \cdots \cup w_{i_r}(M) \cup [N]^M, [M] \rangle = 0$ for all nonnegative integers $i_1, \dots, i_r$ with $i_1 + \cdots + i_r = k$.

Proof. Let $e : N \hookrightarrow M$ be the inclusion map. Then

$$[N]^M \cap [M] = D_M([N]^M) = [N]_M = e_*([N]).$$

Setting $w = w_{i_1}(M) \cup \cdots \cup w_{i_r}(M)$, we have

$$\begin{aligned} \langle w \cup [N]^M, [M] \rangle &= \langle w, [N]^M \cap [M] \rangle \\ &= \langle w, [N]_M \rangle \\ &= \langle w, e_*([N]) \rangle \\ &= \langle e^*(w), [N] \rangle \\ &= \langle e^*(w_{i_1}(M)) \cup \cdots \cup e^*(w_{i_r}(M)), [N] \rangle \\ &= \langle w_{i_1}(N) \cup \cdots \cup w_{i_r}(N), [N] \rangle, \end{aligned}$$

where the last equality holds since the triviality of the normal vector bundle of N implies that $e^*(w_i(M)) = w_i(N)$ for all $i \geq 0$. The proof is complete since, by [13, Théorèmes IV.3, IV.10], N is a boundary if and only if $\langle w_{i_1}(N) \cup \cdots \cup w_{i_r}(N), [N] \rangle = 0$ for all nonnegative integers $i_1, \dots, i_r$ with $i_1 + \cdots + i_r = k$. $\square$

Proof of Theorem 1.2. Every cup product $w_{i_1}(X) \cup \cdots \cup w_{i_r}(X)$ is in $H_{alg}^k(X, \mathbb{Z}/2)$, where $k = i_1 + \cdots + i_r$; cf. [1, 2, 3, 5]. By assumption, for any k -dimensional submanifold N in $\mathcal{F}$, we have

$$\langle w_{i_1}(X) \cup \cdots \cup w_{i_r}(X), [N]_X \rangle = 0,$$

which completes the proof in view of Lemma 2.1. $\square$

The proof of Theorem 1.3 requires further preparations. Given a smooth manifold P , let $\mathfrak{N}_*(P)$ denote the unoriented bordism group of P ; cf. [7]. The following fundamental result will play a crucial role.

Theorem 2.2. *Let P be a smooth manifold. Two smooth maps $f : M \rightarrow P$ and $g : N \rightarrow P$, where M and N are compact smooth manifolds of dimension d , represent the same bordism class in $\mathfrak{N}_*(P)$ if and only if for every nonnegative integer q and every cohomology class v in $H^q(P, \mathbb{Z}/2)$, one has*

$$\langle w_{i_1}(M) \cup \dots \cup w_{i_r}(M) \cup f^*(v), [M] \rangle = \langle w_{i_1}(N) \cup \dots \cup w_{i_r}(N) \cup g^*(v), [N] \rangle$$

for all nonnegative integers $i_1, \dots, i_r$ with $i_1 + \dots + i_r = d - q$.

Reference for the proof. [7, (17.3)]. □

If W is a nonsingular real algebraic variety, then a bordism class in $\mathfrak{N}_*(W)$ is said to be *algebraic* if it can be represented by a regular map $h : Y \rightarrow W$, where Y is a compact nonsingular real algebraic variety.

We will also make use of a certain construction from real algebraic geometry. Let X be a compact nonsingular real algebraic variety. Define $Alg^l(X)$ to be the subset of $H^l(X, \mathbb{Z}/2)$ that consists of all elements v for which there exist a compact irreducible nonsingular real algebraic variety T (depending on v), two points t_0 and t_1 in T , and a cohomology class z in $H_{alg}^l(X \times T, \mathbb{Z}/2)$ such that

$$v = i_{t_1}^*(z) - i_{t_0}^*(z).$$

Here given t in T , we let $i_t : X \rightarrow X \times T$ denote the map defined by $i_t(x) = (x, t)$ for all x in X . For properties and an alternative definition of $Alg^l(-)$, the reader may refer to [10, 11]. Below we will need the following facts. The set $Alg^l(X)$ is a subgroup of $H_{alg}^l(X, \mathbb{Z}/2)$, which has the expected functorial property: if $f : X \rightarrow Y$ is a regular map between compact nonsingular real algebraic varieties, then the induced homomorphism $f^* : H^l(Y, \mathbb{Z}/2) \rightarrow H^l(X, \mathbb{Z}/2)$ satisfies

$$f^*(Alg^l(Y)) \subseteq Alg^l(X);$$

cf. [1, p. 114].

Proposition 2.3. *Let X be a compact nonsingular real algebraic variety. If u is in $H_{alg}^k(X, \mathbb{Z}/2)$ and v is in $Alg^l(X)$, where $k + l = \dim X$, then $\langle u \cup v, [X] \rangle = 0$.*

References for the proof. [10, Theorem 2.1] (cf. also [11, Theorem 4.4]). □

Example 2.4. Given a positive integer l let B^l be an irreducible nonsingular real algebraic variety with precisely two connected components B_0^l and B_1^l , each diffeomorphic to the unit l -sphere S^l (for example $B^l = \{(x_0, \dots, x_l) \in \mathbb{R}^{l+1} \mid x_0^4 - 4x_0^2 + 1 + x_1^2 + \dots + x_l^2 = 0\}$). Let $B = B^{l_1} \times \dots \times B^{l_s}$ and $B_0 = B_0^{l_1} \times \dots \times B_0^{l_s}$. Then

$$H^q(B_0, \mathbb{Z}/2) = \delta^*(H^q(B, \mathbb{Z}/2)) = \delta^*(Alg^q(B))$$

for all $q \geq 0$, where $\delta : B_0 \hookrightarrow B$ is the inclusion map. This assertion is a minor generalization of [11, Example 4.5].

For convenience we introduce the following notation. For any finite collection $\mathcal{F}$ of smooth submanifolds of a compact smooth manifold M , denote by $A^l(\mathcal{F})$ the subgroup of $H^l(M, \mathbb{Z}/2)$ generated by $\{[N]^M \mid N \in \mathcal{F}, \text{codim}_M N = l\}$. If $k + l = \dim M$, then

$$G^k(\mathcal{F}) = \{u \in H^k(M, \mathbb{Z}/2) \mid \langle u \cup v, [M] \rangle = 0 \text{ for all } v \text{ in } A^l(\mathcal{F})\}.$$

This follows immediately from the equality $\langle u \cup v, [M] \rangle = \langle u, D_M(v) \rangle$.

Proof of Theorem 1.3. Observe that $G^0(\mathcal{F})$ contains the subgroup of $H^0(M, \mathbb{Z}/2)$ generated by 1. On the other hand, if Y is a compact irreducible nonsingular real algebraic variety, then $H_{alg}^0(Y, \mathbb{Z}/2)$ is the subgroup of $H^0(Y, \mathbb{Z}/2)$ generated by 1. Hence, without loss of generality, we can enlarge the collection $\mathcal{F}$ by adding to it finitely many 0-dimensional submanifolds of M , each consisting of an even number of points. Thus we can assume that $\mathcal{F}$ has the following property: if M_1 and M_2 are distinct connected components of M , then $\{x_1, x_2\}$ is in $\mathcal{F}$ for some points x_1 in M_1 and x_2 in M_2 .

Let $\mathcal{F} = \{N_1, \dots, N_s\}$ and $l_i = \text{codim}_M N_i$ for $1 \leq i \leq s$. Note that $l_i \geq 1$. We will use notation introduced in Example 2.4. There is a smooth map $f_i : M \rightarrow B^{l_i}$ such that $f_i(M) \subseteq B_0^{l_i}$ and $f_i^*(H^{l_i}(B^{l_i}, \mathbb{Z}/2))$ is the subgroup of $H^{l_i}(M, \mathbb{Z}/2)$ generated by $[N_i]^M$. This assertion follows from a well-known fact that if the normal vector bundle of a smooth submanifold N of M is trivial and $\text{codim}_M N = l \geq 1$, then one can find a smooth map $h : M \rightarrow S^l$ with $h^*(\lambda) = [N]^M$, where λ is the unique generator of $H^l(S^l, \mathbb{Z}/2) \cong \mathbb{Z}/2$; cf. [13, Théorème II.2]. Setting $f = (f_1, \dots, f_s) : M \rightarrow B = B^{l_1} \times \dots \times B^{l_s}$, we have

$$(1) \quad f^*(H^q(B, \mathbb{Z}/2)) = A^q(\mathcal{F}) \quad \text{for all } q \geq 1.$$

Let $c : M \rightarrow B$ be a constant map, whose single value is a point in B_0 . Since M has an algebraic model, the bordism class of c in $\mathfrak{N}_*(B)$ is algebraic. We claim that for every nonnegative integer q and every cohomology class b in $H^q(B, \mathbb{Z}/2)$,

$$(2) \quad \langle w_{i_1}(M) \cup \dots \cup w_{i_r}(M) \cup f^*(b), [M] \rangle = \langle w_{i_1}(M) \cup \dots \cup w_{i_r}(M) \cup c^*(b), [M] \rangle$$

for all nonnegative integers $i_1, \dots, i_r$ with $i_1 + \dots + i_r = \dim M - q$. The claim is obvious if $q = 0$. For $q \geq 1$ condition (2) is equivalent to

$$\langle w_{i_1}(M) \cup \dots \cup w_{i_r}(M) \cup f^*(b), [M] \rangle = 0.$$

The last equality holds in view of (1) and Lemma 2.1, and hence (2) is proved.

It follows from (2) and Theorem 2.2 that f and c represent the same bordism class in $\mathfrak{N}_*(B)$. In particular, the bordism class of f in $\mathfrak{N}_*(B)$ is algebraic. By [1, Theorem 2.8.4], there exist an algebraic model X of M , a smooth diffeomorphism $\varphi : X \rightarrow M$, and a regular map $g : X \rightarrow B$ such that g is homotopic to $f \circ \varphi$. It remains to show that X and φ have the properties stated in the theorem. This can be done as follows. Since g is homotopic to $f \circ \varphi$, we get $\varphi^* \circ f^* = (f \circ \varphi)^* = g^*$ in cohomology, and hence for all $q \geq 0$,

$$(3) \quad \varphi^*(f^*(H^q(B, \mathbb{Z}/2))) = g^*(H^q(B, \mathbb{Z}/2)) = g^*(Alg^q(B)),$$

where the last equality follows from the inclusion $g(X) \subseteq B_0$ and Example 2.4. We also have

$$(4) \quad g^*(Alg^q(B)) \subseteq Alg^q(X),$$

the map $g : X \rightarrow B$ being regular. Conditions (1), (3), and (4) combined yield

$$(5) \quad \varphi^*(A^q(\mathcal{F})) \subseteq Alg^q(X) \text{ for all } q \geq 1.$$

Let k and l be integers satisfying $k \geq 0$, $l \geq 0$, and $k + l = m$, where $m = \dim M = \dim X$. Since $\varphi^*(A^l(\mathcal{F})) = A^l(\varphi^*\mathcal{F})$ and

$$G^k(\varphi^*\mathcal{F}) = \{u \in H^k(X, \mathbb{Z}/2) \mid \langle u \cup v, [X] \rangle = 0 \text{ for all } v \in A^l(\varphi^*\mathcal{F})\},$$

it follows from (5) and Proposition 2.3 that

$$(6) \quad H_{alg}^k(X, \mathbb{Z}/2) \subseteq G^k(\varphi^*\mathcal{F})$$

for $0 \leq k \leq m-1$. In fact (6) holds for all $k \geq 0$, the equality $G^k(\varphi^*\mathcal{F}) = H^k(X, \mathbb{Z}/2)$ being automatically satisfied if $k \geq m$.

In order to complete the proof, we need to demonstrate that X is irreducible. This is obvious if M is connected. Suppose then that M is disconnected and X is reducible. Let X_1 and X_2 be distinct irreducible components of X . There are points p_1 in X_1 and p_2 in X_2 such that $P = \{p_1, p_2\}$ is in $\varphi^*\mathcal{F}$ (see the beginning of the proof). In particular, $[P]^X$ belongs to $A^m(\varphi^*\mathcal{F}) = \varphi^*(A^m(\mathcal{F}))$, which in view of (5) implies that $[P]^X$ is in $Alg^m(X)$. If $e : X_1 \hookrightarrow X$ is the inclusion map, then $e^*(Alg^m(X)) \subseteq Alg^m(X_1)$, and hence $e^*([P]^X) = [\{p_1\}]^{X_1}$ is in $Alg^m(X_1)$. However, $\langle [\{p_1\}]^{X_1}, [X_1] \rangle = 1$, which contradicts Proposition 2.3. Thus X is irreducible, as required. $\square$

REFERENCES

- [1] S. Akbulut and H. King, *Topology of Real Algebraic Sets*, Math. Sci. Research Institute Publ. 25, Springer, 1992. MR1225577 (94m:57001)
- [2] J. Bochnak, M. Coste and M.-F. Roy, *Real Algebraic Geometry*, Ergebnisse der Math. und ihrer Grenzgeb. Folge (3), Vol. 36, Springer, 1998. MR1659509 (2000a:14067)
- [3] J. Bochnak and W. Kucharz, *On homology classes represented by real algebraic varieties*, Banach Center Publications 44 (1998), 21-35. MR1677394 (2000b:14080)
- [4] J. Bochnak and W. Kucharz, *On approximation of smooth submanifolds by nonsingular real algebraic subvarieties*, Ann. Sci. Ec. Norm. Sup. 36 (2003), 685-690. MR2032984 (2004j:14063)
- [5] A. Borel and A. Haefliger, *La classe d'homologie fondamentale d'un espace analytique*, Bull. Soc. Math. France 89 (1961), 461-513. MR0149503 (26:6990)
- [6] G. Bredon, *Topology and Geometry*, New York, Berlin, Heidelberg, Springer, 1997. MR1700700 (2000b:55001)
- [7] P. Conner, *Differentiable Periodic Maps*, Lecture Notes in Math. Vol. 738, Springer, 1979. MR548463 (81f:57018)
- [8] J. Huisman and F. Mangolte, *Every orientable Seifert 3-manifold is a real component of a uniruled algebraic variety*, Topology 44 (2005), 63-71. MR2104001 (2005h:14139)
- [9] J. Kollár, *The topology of real and complex algebraic varieties*, Taniguchi Conference on Mathematics, Nara 1998, Advanced Studies in Pure Mathematics, Vol. 31, Mathematical Society of Japan, Tokyo, 2001, pp. 127-145. MR1865090 (2003a:14033)
- [10] W. Kucharz, *Algebraic equivalence and homology classes of algebraic cycles*, Math. Nachr. 180 (1996), 135-140. MR1397672 (97e:14009)
- [11] W. Kucharz, *Algebraic cycles and algebraic models of smooth manifolds*, J. Algebraic Geometry 11 (2002), 101-127. MR1865915 (2002j:14065)
- [12] J. Nash, *Real algebraic manifolds*, Ann. of Math. 56 (1952), 405-421. MR0050928 (14:403b)
- [13] R. Thom, *Quelques propriétés globales des variétés différentiables*, Comment. Math. Helv. 28 (1954), 17-86. MR0061823 (15:890a)
- [14] A. Tognoli, *Su una congettura di Nash*, Ann. Scuola Norm. Sup. Pisa, Sci. Fis. Mat. 27 (3) (1973), 167-185. MR0396571 (53:434)

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF NEW MEXICO, ALBUQUERQUE, NEW MEXICO 87131-1141

E-mail address: kucharz@math.unm.edu