SLICE KNOTS WITH DISTINCT OZSVÁTH-SZABÓ AND RASMUSSEN INVARIANTS

CHARLES LIVINGSTON

(Communicated by Daniel Ruberman)

ABSTRACT. As proved by Hedden and Ording, there exist knots for which the Ozsváth-Szabó and Rasmussen smooth concordance invariants, τ and s , differ. The Hedden-Ording examples have nontrivial Alexander polynomials and are not topologically slice. It is shown in this note that a simple manipulation of the Hedden-Ording examples yields a topologically slice Alexander polynomial one knot for which τ and s differ. Manolescu and Owens have previously found a concordance invariant that is independent of both τ and s on knots of polynomial one, and as a consequence have shown that the smooth concordance group of topologically slice knots contains a summand isomorphic to $\mathbf{Z} \oplus \mathbf{Z}$. It thus follows quickly from the observation in this note that this concordance group contains a summand isomorphic to $\mathbf{Z} \oplus \mathbf{Z} \oplus \mathbf{Z}$.

The Ozsváth-Szabó and Rasmussen knot concordance invariants [OS, Ra], τ and s , are each powerful invariants, sufficient to resolve the Milnor conjecture on the 4-genus of torus knots. It had been conjectured that $\tau = s/2$, but recently Hedden and Ording [HO] provided a counterexample, showing that certain doubled knots, including $D_+(T_{2,3}, 2)$ and $D_+(T_{2,5}, 4)$, satisfy $\tau = 0$ and $s = 2$. Here $D_+(T_{p,q}, t)$ denotes the t -twisted positive double of the (p, q) -torus knot. These examples demonstrate the richness of these new invariants. However, in and of themselves, they do not reveal any new structure of the concordance group. For instance, although some of these examples are algebraically slice, all can be shown not to be even topologically slice using Casson-Gordon invariants [CG]. (See [Gi] for techniques that resolve the nonsliceness of these particular doubled knots, and [K] for a general discussion of Casson-Gordon invariants and doubled knots.)

In this note it will be shown that the basic Hedden-Ording examples can be manipulated to yield a knot with Alexander polynomial one (and thus, by Freedman [F], a topologically slice knot) for which τ and $s/2$ differ.

The smooth concordance group contains a subgroup $\mathcal{S}$ consisting of topologically slice knots. In [L2] it was shown that τ and $s/2$ agree and are nonzero on some polynomial one knots: both invariants take value 1 on $D_+(T_{2,3}, 0)$ and $D_+(T_{2,5}, 0)$. It followed that $\mathcal{S}$ contains a $\mathbf{Z}$ summand. Manolescu and Owens [MO] developed a concordance invariant δ and showed $\delta(D_+(T_{2,3}, 0)) \neq \delta(D_+(T_{2,5}, 0))$, and thus

Received by the editors April 12, 2006.

2000 *Mathematics Subject Classification.* Primary 57M25.

Key words and phrases. Slice knot, Ozsváth-Szabó invariant, Rasmussen invariant, polynomial one.

The author's research was supported by the NSF.

©2007 American Mathematical Society
 Reverts to public domain 28 years from publication

showed $\mathcal{S}$ contains a summand isomorphic to $\mathbf{Z} \oplus \mathbf{Z}$. The example here shows the independence of τ and $s/2$ on $\mathcal{S}$ and it follows from a quick determinant calculation that $\tau, s/2$, and δ yield a summand isomorphic to $\mathbf{Z} \oplus \mathbf{Z} \oplus \mathbf{Z}$. (Note that these are *summands*. Gompf [Go] showed that there are Alexander polynomial one knots of infinite order in smooth concordance, using the work of Donaldson [D]. This was extended by Endo [E], who showed that there is a subgroup of infinite rank generated by Alexander polynomial one knots.)

1. AN ALEXANDER POLYNOMIAL ONE KNOT WITH $\tau \neq s/2$

The doubled knot $D_+(K, t)$ bounds a Seifert surface built by adding two bands to a disk, one with framing -1 and the other with framing t . With respect to the corresponding basis of the first homology of the surface, the Seifert matrix is

$$\begin{pmatrix} -1 & 1 \\ 0 & t \end{pmatrix}.$$

It follows from [HO] that the connected sum

$$K = D_+(T_{2,3}, 2) \# D_+(T_{2,3}, 2) \# D_+(T_{2,5}, 4)$$

satisfies $\tau(K) = 0$ and $s(K) = 6$. With respect to the natural Seifert surface and bases for homology, the Seifert form of K is given by the matrix V_1 below:

$$V_1 = \begin{pmatrix} -1 & 1 & 0 & 0 & 0 & 0 \\ 0 & 2 & 0 & 0 & 0 & 0 \\ 0 & 0 & -1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 2 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 & 1 \\ 0 & 0 & 0 & 0 & 0 & 4 \end{pmatrix}, \quad V_2 = \begin{pmatrix} 3 & 0 & 1 & 0 & 0 & 4 \\ -1 & 2 & 0 & 0 & 0 & 0 \\ 1 & 0 & -1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 2 & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 & 1 \\ 4 & 0 & 0 & 0 & 0 & 4 \end{pmatrix}.$$

If the first band in the Seifert surface is cut and reattached after twisting and linking with the other bands, a knot J with the Seifert matrix V_2 above can be constructed.

For a knot with Seifert matrix V , the Alexander polynomial $\Delta_K(t)$ is given by $\det(V - tV^t)$; using this, a calculation gives $\Delta_J(t) = t^3$, and thus J has a trivial Alexander polynomial.

According to Livingston and Naik [LN], cutting a band in a Seifert surface for a knot and reattaching it can change τ by at most ± 1 and can change s by at most ± 2 . It follows that $\tau(J) \in \{-1, 0, 1\}$ and $s(J) \in \{4, 6\}$. (Note: s is even and is bounded by twice the genus.) Clearly, $\tau(J) \neq s(J)/2$.

REFERENCES

- [CG] A. Casson, C. McA. Gordon, *Cobordism of classical knots*, in *À la recherche de la topologie perdue*, ed. by Guillou and Marin, Progress in Mathematics, Volume 62, 1986. (Originally published as an Orsay Preprint, 1975.) MR900252
- [D] S. K. Donaldson, *An application of gauge theory to four-dimensional topology*, J. Differential Geom. 18 (1983), no. 2, 279–315. MR710056 (85c:57015)
- [E] H. Endo, *Linear independence of topologically slice knots in the smooth cobordism group*, Topology Appl. 63 (1995), no. 3, 257–262. MR1334309 (96c:57010)
- [F] M. Freedman, *The topology of four-dimensional manifolds*, J. Differential Geom. 17 (1982), 357–453. MR679066 (84b:57006)
- [Gi] P. Gilmer, *Slice knots in S^3* , Quart. J. Math. Oxford Ser. (2) 34 (1983), 305–322. MR711523 (85d:57004)

- [Go] R. Gompf, *Smooth concordance of topologically slice knots*, Topology 25 (1986), no. 3, 353–373. MR842430 (87i:57004)
- [HO] M. Hedden, P. Ording, *The Ozsváth-Szabó and Rasmussen concordance invariants are not equal* (2005), [arxiv.org/math/0512348](https://arxiv.org/abs/math/0512348).
- [K] S.-G. Kim, *Polynomial splittings of Casson-Gordon invariants*, Math. Proc. Cambridge Philos. Soc. 138 (2005), 59–78. MR2127228 (2005m:57018)
- [L1] C. Livingston, *Splitting the concordance group of algebraically slice knots*, Geom. Topol. 7 (2003), 641–643. MR2026553 (2004j:57009)
- [L2] C. Livingston, *Computations of the Ozsváth-Szabó knot concordance invariant*, Geom. Topol. 8 (2004), 735–742. MR2057779 (2005d:57019)
- [LN] C. Livingston, S. Naik, *Ozsváth-Szabó and Rasmussen invariants of doubled knots*, Algebr. Geom. Topol. 6 (2006), 651–657. MR2240910 (2007d:57023)
- [MO] C. Manolescu, B. Owens, *A concordance invariant from the Floer homology of double branched covers* (2005), [arxiv.org/math.GT/0508065](https://arxiv.org/abs/math.GT/0508065).
- [OS] P. Ozsváth, Z. Szabó, *Knot Floer homology and the four-ball genus*, Geom. Topol. 7 (2003), 615–639. MR2026543 (2004i:57036)
- [Ra] J. A. Rasmussen, *Khovanov homology and the slice genus* (2003), [arxiv.org/math/0306378](https://arxiv.org/abs/math/0306378).

DEPARTMENT OF MATHEMATICS, INDIANA UNIVERSITY, BLOOMINGTON, INDIANA 47405

E-mail address: livingst@indiana.edu