MINIMAL BOUNDED INDEX SUBGROUP FOR DEPENDENT THEORIES

SAHARON SHELAH

(Communicated by Julia Knight)

ABSTRACT. For a dependent theory T , in $\mathfrak{C}_T$ for every type definable group G , the intersection of type definable subgroups with bounded index is a type definable subgroup with bounded index.

§0. INTRODUCTION

Assume that T is a dependent (complete first-order) theory, $\mathfrak{C}$ is a $\bar{\kappa}$ -saturated model of T (a monster), G is a type definable (in $\mathfrak{C}$) group in $\mathfrak{C}$ (of course we consider only types of cardinality $< \bar{\kappa}$).

A type definable subgroup H of G is called bounded if the index $(G : H)$ is $< \bar{\kappa}$. We prove that there is a minimal bounded definable subgroup. The first theorem on this line for T stable is due to Baldwin and Saxl [BaSx76].

Recently Hrushovski, Peterzil and Pillay [HPP0x] investigated definable groups, o-minimality and measure. In an earlier work on definable subgroups in o-minimal T in Berarducci, Otero, Peterzil and Pillay [BOPP05] the minimal type-definable bounded index theorem and more results are proved for o-minimal theories.

Hrushovski, in a lecture at the Hebrew University, mentioned that he, Peterzil and Pillay had observed the main result of the current paper, but assuming in addition the existence of an invariant measure on the group in question, and Hrushovski asked if the measure assumption could be removed. So we answer it positively. The current version of their paper [HPP0x] includes an exposition of our proof.

Recent works of the author on dependent theories are [Sh:783] (see §3, §4 on groups) [Sh:863] (e.g., the first-order theory of the p -adics is strongly¹ dependent but not strongly² dependent, see end of §1; on strongly² dependent fields, see §5) and [Sh:F705]. This work is continued in [Sh:F753] (getting mainly a parallel result for G abelian and $\mathbb{L}_{\infty, \bar{\kappa}}$ -definable subgroups).

I thank Aviv Tatarski and Ilay Kaplan for their very careful checking.

Received by the editors January 22, 2006 and, in revised form, July 19, 2006, July 30, 2006, and October 19, 2006.

2000 *Mathematics Subject Classification*. Primary 03C45; Secondary 03C60.

The author would like to thank the Israel Science Foundation for partial support of this research (Grant No. 242/03), Publication 876. I would like to thank Alice Leonhardt for the beautiful typing.

§1

1.1 Theorem. Assume T is a dependent (complete first-order) theory, $\mathfrak{C}$ a $\bar{\kappa}$ -saturated model for T . We consider types of cardinality $< \bar{\kappa}$.

1) If $\circledast$ below holds, then:

- (α) $q(\mathfrak{C})$ is a subgroup of $p(\mathfrak{C})$,
- (β) $q(\mathfrak{C})$ is of index $< \bar{\kappa}$,
- (γ) essentially $q(x) \setminus p(x)$ is of cardinality $\leq \lambda := |T|^{\aleph_0}$ (i.e., for some $q'(x) \subseteq q(x)$ of cardinality $\leq \lambda$, $q(x)$ is equivalent to $p(x) \cup q'(x)$),
- (δ) we can strengthen (α), (β) to $q(\mathfrak{C})$ is a subgroup of index $\leq 2^\lambda$,

where

- $\circledast$ (a) $p(x)$ is a type such that $p(\mathfrak{C})$ is a group which we call G (with some definable operations xy, x^{-1} and the identity e_G which is constant here),
- (b) $q(x) = p(x) \cup \bigcup \{r(x) : r(x) \in \mathbf{R}\}$, where
- (c) $\mathbf{R} = \{r(x) : r(x) \text{ a type such that } (p \cup r)(\mathfrak{C}) \text{ is a subgroup of } p(\mathfrak{C}) \text{ of index } < \bar{\kappa}\}$.

2) There exists some $q' \subseteq q$ over $\text{Dom}(p)$, equivalent to q and such that $|q'| \leq |T| + |\text{Dom}(p)|$. So $(p(\mathfrak{C}) : q(\mathfrak{C})) \leq 2^{|\text{Dom}(p)| + |T|}$.

3) If $r_i(x) \in \mathbf{R}$ for $i < \lambda^+$, then for some $\alpha < (|T|^{\aleph_0})^+$ we have $(p(x) \cup \bigcup \{r_i(x) : i < \alpha\})(\mathfrak{C}) = (p(x) \cup \bigcup \{r_i(x) : i < \lambda^+\})(\mathfrak{C})$.

Proof. 1) Note

- $\circledast_1$ $\mathbf{R}$ is closed under unions of $< \bar{\kappa}$.
- $\circledast_2$ (a) If $r(x) \in \mathbf{R}$, $r'(x) \subseteq r(x)$ is countable, then there is a countable $r''(x) \subseteq r(x)$ including $r'(x)$ which belongs to $\mathbf{R}$.
- (b) If $p(x) \subseteq r(x) \in \mathbf{R}$ and $r(x)$ is closed under conjunctions and $r'(x) \subseteq r(x)$ is countable, then we can find $\psi_n(x, \bar{b}_n)$ for $n < \omega$ such that
- (α) $\psi_n(x, \bar{b}_n) \in r(x)$,
- (β) $\psi_{n+1}(x, \bar{b}_{n+1}) \vdash \psi_n(x, \bar{b}_n)$,
- (γ) $\psi_{n+1}(x, \bar{b}_{n+1}), \psi_{n+1}(y, \bar{b}_{n+1}) \vdash \psi_n(x^{-1}y, \bar{b}_n) \wedge \psi_n(x^{-1}, \bar{b}_n) \wedge \psi_n(xy, \bar{b}_n)$,
- (δ) $\psi_n(x, \bar{b}_{n+1}) \vdash \varphi_n(x, \bar{a}_n)$, where $\{\varphi_n(x, \bar{a}_n) : n < \omega\}$ list r' ,
- (ε) $\mathfrak{C} \models \psi_n(e_G, \bar{b}_n)$, actually follows from clause (α).

[Why? Let $r'(x) = \{\varphi_n(x, \bar{a}_n) : n < \omega\}$ (can use $\varphi_n = (x = x)$). Without loss of generality, $r(x)$ is closed under conjunctions and also $r'(x)$ is. Now we choose $\psi_n(x, \bar{b}_n)$ by induction on $n < \omega$ such that $\psi_{n+1}(x, \bar{b}_{n+1}) \wedge \psi_{n+1}(y, \bar{b}_{n+1}) \vdash \psi_n(xy^{-1}, \bar{b}_n) \wedge \varphi_n(x, \bar{a}_n) \wedge \psi_n(xy, \bar{b}_n)$; notice that trivially $e_G \in \varphi_n(\mathfrak{C}, \bar{a}_n) \cap \psi_n(\mathfrak{C}, \bar{b}_n)$. Such a formula exists as $(p(x) \cup r(x)) \cup (p(y) \cup r(y)) \vdash \psi_n(xy^{-1}, \bar{b}_n) \wedge \varphi_n(x, \bar{a}_n) \wedge \psi_n(xy, \bar{b}_n)$.

Now $r''(x) = \{\varphi_n(x, \bar{a}_n), \psi_n(x, \bar{b}_n) : n < \omega\}$ is as required in clause (a), $\langle \psi_n(x, \bar{b}_n) : n < \omega \rangle$ in (b).]

In the conclusion of Theorem 1.1, Clause (α) is obvious.

Assume toward a contradiction that the conclusion (β) + (γ) fails. So we can choose (c_α, r_α) by induction on $\alpha < \lambda^+$ such that

- $\circledast_3$ (a) $c_\alpha \in (p(x) \cup \bigcup \{r_\beta : \beta < \alpha\})(\mathfrak{C})$,
- (b) $r_\alpha = \{\psi_n^\alpha(x, \bar{b}_n^\alpha) : n < \omega\} \subseteq q$ and $\bar{b}_n^\alpha \triangleleft \bar{b}_{n+1}^\alpha$,
- (c) $r_\alpha \in \mathbf{R}$, and $\psi_{n+1}(x, \bar{b}_{n+1}^\alpha) \vdash \psi_n(x, \bar{b}_n^\alpha)$,
- (d) c_α does not realize r_α , in fact $\mathfrak{C} \models \neg \psi_0^\alpha(c_\alpha, \bar{b}_0^\alpha)$, hence $c_\alpha \notin q(\mathfrak{C})$.

[Why? Arriving to α , we try to let $q_\alpha = p(x) \cup \{r_\beta : \beta < \alpha\}$, so $q_\alpha \subseteq q$ and by cardinality considerations, $q_\alpha(\mathfrak{C})$ is a subgroup of G of index $< \bar{\kappa}$. So by our assumption toward a contradiction, $q_\alpha(x) \not\models q(x)$; hence there is $r_\alpha^*(x) \in \mathbf{R}$ such that $q_\alpha(x) \not\models r_\alpha^*$, so $q_\alpha(x) \not\models \vartheta_\alpha(x, \bar{d}_\alpha)$ for some $\vartheta_\alpha(x, \bar{d}_\alpha) \in r_\alpha^*$. Let $r_\alpha(x) = \{\psi_n^\alpha(x, \bar{b}_n^\alpha) : n < \omega\} \subseteq r_\alpha^*(x)$ belong to $\mathbf{R}$ and be such that $\vartheta_\alpha(x, \bar{d}_\alpha) = \psi_0^\alpha(x, \bar{b}_0^\alpha)$; it exists by $\otimes_2(\alpha)$ above.

Without loss of generality, we can assume $\otimes_4 - \otimes_8$:

- $\otimes_4$ e_G is an individual constant, $\bar{y}_n^\alpha \leq \bar{y}_{n+1}^\alpha$ and $\psi_{n+1}^\alpha(x, \bar{y}_{n+1}^\alpha) \vdash \psi_n^\alpha(x, \bar{y}_n^\alpha)$ and $(\psi_{n+1}^\alpha(x_1, \bar{y}_{n+1}^\alpha) \wedge \psi_{n+1}^\alpha(x_2, \bar{y}_{n+1}^\alpha)) \vdash (\psi_n^\alpha(e_G, \bar{y}_n^\alpha) \wedge \psi_n^\alpha(x_1 x_2^{-1}, \bar{y}_n^\alpha))$.

[Why? As in the proof of $\otimes_2$ above, i.e., during the induction in the proof of $\otimes_3$, we use $\otimes_2(\beta)$ and get $\psi_n^\alpha(x, \bar{y}_n^\alpha), \bar{b}_n^\alpha$ and without loss of generality $\bar{b}_n^\alpha \leq \bar{b}_{n+1}^\alpha, \bar{y}_n^\alpha \triangleleft \bar{y}_{n+1}^\alpha$ (as we can change the order and name the free variable and add dummy variables). Now we define $\psi_n^{\alpha,*}(x, \bar{y}_n^\alpha)$ by induction on n by

$$\begin{aligned} \psi_0^{\alpha,*}(x, \bar{y}_0^\alpha) &= \psi_0^\alpha(x, \bar{y}_0^\alpha), \psi_{n+1}^{\alpha,*}(x, \bar{y}_{n+1}^\alpha) = \psi_{n+1}^\alpha(x, \bar{y}_{n+1}^\alpha) \\ &\wedge \bigwedge_{m \leq n} (\forall z) [\psi_{m+1}^\alpha(z, \bar{y}_{m+1}^\alpha) \rightarrow \psi_m^\alpha(z, \bar{y}_m^\alpha)] \\ &\wedge \bigwedge_{m \leq n} (\forall z_1, z_2) [\psi_{m+1}^\alpha(z_1, \bar{y}_{m+1}^\alpha) \wedge \psi_{m+1}^\alpha(z_2, \bar{y}_{m+1}^\alpha) \\ &\quad \rightarrow \psi_m^\alpha(e_G, \bar{y}_m^\alpha) \wedge \psi_m^\alpha(z_1 z_2^{-1}, \bar{y}_m^\alpha)]. \end{aligned}$$

So clearly $\langle \psi_n^{\alpha,*}(x, \bar{y}_n^\alpha) : n < \omega \rangle$ satisfies $\otimes_4$ and $\mathfrak{C} \models (\forall x) [\psi_n^{\alpha,*}(x, \bar{b}_n^\alpha) \equiv \psi_n^\alpha(x, \bar{b}_n^\alpha)]$. So renaming we are done.]

- $\otimes_5$ $\psi_n^\alpha(x, \bar{y}_n^\alpha) = \psi_n(x, \bar{y}_n)$ and $\psi_{n+1}^\alpha(x, \bar{y}_{n+1}^\alpha) \vdash \psi_n(x, \bar{y}_n)$.

[Why? By the pigeon hole principle.]

- $\otimes_6$ $\langle c_\alpha \bar{\mathbf{a}}_\alpha : \alpha < \lambda^+ \rangle$ is an indiscernible sequence over $\text{Dom}(p)$ where $\bar{\mathbf{a}}_\alpha = \cup \{b_n^{-\alpha} : n < \omega\} \bar{b}_0^\alpha \wedge \bar{b}_1^\alpha \wedge \bar{b}_2^\alpha \wedge \dots$; note that by clause (b) of $\otimes_3$ we have $\bar{b}_n^\alpha = \bar{\mathbf{a}}_\alpha \upharpoonright k_n$.

[Why? By the Ramsey theorem and compactness.]

- $\otimes_7$ If $\alpha < \beta < \gamma$, then $c_\alpha c_\beta^{-1} \in r_\gamma(\mathfrak{C})$.

[Why? By the indiscernibility, without loss of generality, γ is infinite, so $\gamma \geq \omega$ and $\langle c_i : i < \gamma \rangle$ is an indiscernible sequence over $\text{Dom}(p) \cup \bar{b}_\gamma$ of elements of $p(\mathfrak{C})$ pairwise nonequivalent modulo the subgroup $G_\gamma = (p \cup r_\gamma)(\mathfrak{C})$. Then we can extend it to $\langle \bar{c}_i : i < \bar{\kappa} \rangle$, an indiscernible sequence over $\text{Dom}(p) \cup \bar{b}_\gamma$ and arrive at $\alpha < \beta \Rightarrow c_\alpha c_\beta^{-1} \notin G_\gamma \Rightarrow c_\beta c_\alpha^{-1} \notin G_\gamma$ so $\langle c_\alpha G_\gamma : \alpha < \bar{\kappa} \rangle$ are pairwise distinct (equivalently $\langle G_\gamma c_\alpha : \alpha < \bar{\kappa} \rangle$ are pairwise distinct), a contradiction.]

- $\otimes_8$ $c_\alpha \in r_\beta(\mathfrak{C})$ iff $\alpha \neq \beta$.

[Why? Let

$$\begin{aligned} c_\alpha^* &= c_{2\alpha+1} \cdot (c_{2\alpha})^{-1}, \\ r_\alpha^* &= r_{2\alpha}. \end{aligned}$$

So:

- (i) If $\beta < \alpha$, then $c_\alpha^* \in (p \cup r_\beta^*)(\mathfrak{C})$, because $c_{2\alpha+1}, c_{2\alpha}$ belong to the subgroup $(p \cup r_{2\beta})(\mathfrak{C})$ by clause (a) of $\otimes_3$.
- (ii) If $\beta > \alpha$, then c_α^* belongs to $(p \cup r_\beta^*)(\mathfrak{C})$ by $\otimes_7$.
- (iii) If $\beta = \alpha$, then c_α^* does not belong to $(p \cup r_\beta^*)(\mathfrak{C})$ because:

- (α) it is a subgroup,
- (β) $c_{2\alpha+1}$ belongs to it by clause (a) of $\otimes_3$ and
- (γ) $c_{2\alpha}$ does not belong to it by clause (e) of $\otimes_3$.

Let $\bar{a}_\alpha^* = \bar{a}_{2\alpha+1}$, $\bar{b}_n^{\alpha,*} = \bar{b}_n^{2\alpha}$ retaining the same ψ 's. So we have obtained an example as required in $\otimes_8$ (not losing the other demands).]

$\otimes_9$ For some $n < \omega$ for every α , we have if $d_1, d_2 \in (p \cup r_\alpha)(\mathfrak{C})$, then $d_1 c_\alpha d_2 \notin \psi_n(\mathfrak{C}, \bar{b}_n^\alpha)$; without loss of generality, $n = 1$.

[Why? Fix α . If this holds for some $\psi_n(-, \bar{b}_n^\alpha)$, by indiscernibility, renaming the φ_i 's, this is O.K. Otherwise for each $n < \omega$ there are $d_1^n, d_2^n \in (p \cup r_\alpha)(\mathfrak{C})$ such that $\mathfrak{C} \models \psi_n(d_1^n c_\alpha d_2^n, \bar{b}_n^\alpha)$. By compactness for some $d_1^*, d_2^* \in (p \cup r_\alpha)(\mathfrak{C})$ we have $\models \psi_n[d_1^* c_\alpha d_2^*, \bar{b}_n^\alpha]$ for every $n < \omega$. So $d_1^* c_\alpha d_2^*$ belongs to the subgroup $(p \cup r_\alpha)(\mathfrak{C})$, but also d_1^*, d_2^* belongs to it; hence c_α belongs, a contradiction. Alternatively, note that $n = 2$ is O.K.: let $c' = d_1 c d_2$ and assume toward a contradiction that $c' \in \psi_2(\mathfrak{C}, \bar{b}_2^\alpha)$ and let $d'_1 = (d_1)^{-1}, d'_2 = (d_2)^{-1}$, so clearly $d_1, d'_2 \in (p \cup r_\alpha)(\mathfrak{C}) \subseteq \psi_2(\mathfrak{C}, \bar{b}_2^\alpha)$. Now by $\otimes_4$ as $d'_1, c' \in \psi_2(\mathfrak{C}, \bar{b}_1^\alpha)$ it follows that $d_2(c')^{-1} \in \psi_1(\mathfrak{C}, \bar{b}_1^\alpha)$. As $d_1, d'_2(c')^{-1} \in \psi_1(\mathfrak{C}, \bar{b}_1^\alpha)$ by $\otimes_4$ we have $d'_1(d_2(c')^{-1})^{-1} \in \psi_0(\mathfrak{C}, \bar{b}_0^\alpha)$. But $c' = d_1 c_\alpha d_2$; hence $c_\alpha = d'_1((d'_2)^{-1}(c')^{-1})^{-1} = d'_1(d_2(c')^{-1})^{-1}$, but $d'_1(d_2(c')^{-1})^{-1} \in \psi_0(\mathfrak{C}, \bar{b}_0^\alpha)$ by the previous sentence, whereas $c_\alpha \notin \psi_0(\mathfrak{C}, \bar{b}_1^\alpha)$ by $\otimes_3(e)$, a contradiction.]

$\otimes_{10}$ If $w = \{i_1, \dots, i_n\}, i_1 < \dots < i_n < \lambda^+$, and $d_w := c_{i_1} c_{i_2} \dots c_{i_n} \in G$ and $\alpha < \lambda^+$, then $\models \varphi_1[d_w, \bar{b}_\alpha^1] \Leftrightarrow \alpha \notin w$.

[Why? If $\alpha \in w$, let k be such that $\alpha = i_k$, so $c_{i_1}, \dots, c_{i_{k-1}} \in (p \cup r_\alpha)(\mathfrak{C})$ by $\otimes_8$ and similarly $c_{i_{k+1}} \dots c_{i_n} \in (p \cup r_\alpha)(\mathfrak{C})$; hence

$$d_w = (c_{i_1} \dots c_{i_k}) c_{i_k} (c_{i_{k+1}} \dots c_{i_n}) \notin (p \cup r_\alpha)(\mathfrak{C})$$

by $\otimes_9$.

Second, if $\alpha \notin w$, this holds by $\otimes_8$ as $\{c_{i_\ell} : \ell < n\}$ is included in the subgroup $(p \cup r_\alpha)(\mathfrak{C})$.]

So we get a contradiction to “ T is dependent”; hence clauses (β), (γ) hold. Also clause (δ) follows by the following observation:

Observation: If $r(x) \in \mathbf{R}$ and $|r(x)| \leq \theta$, then $(p(\mathfrak{C}) : (p \cup r)(\mathfrak{C})) \leq 2^\theta$ (except for being just finite when θ is finite).

Proof. If θ is finite, then the proof follows by compactness. If θ is infinite, then without loss of generality, r is closed under conjunctions. Let $r = \{\varphi_i(x, \bar{\mathbf{b}}) : i < \theta\}$, where $\bar{\mathbf{b}}$ is possibly infinite.

Let u be a set of ordinals ($< \bar{\kappa}$) such that $\bar{\kappa} > |u| > (p(\mathfrak{C}) : (p \cup r)(\mathfrak{C}))$. Now for each $i < \theta$, let $\Gamma_{i,u} = \bigcup \{p(x_\alpha) : \alpha \in u\} \cup \{\neg \varphi_i(x_\alpha x_\beta^{-1}, \bar{\mathbf{b}}) : \alpha < \beta \text{ from } u\}$. So for some finite $u_i^* \subseteq u$, Γ_{i,u_i^*} is contradictory, so Γ_{i,n_i} is contradictory, when $n_i = |u_i|$. It suffices to use $(2^\theta)^+ \rightarrow (\dots n_i \dots)_{i < \theta}$ (why? let $\langle c_\alpha : \alpha < (2^\theta)^+ \rangle$ exemplify the failure and let $\zeta_{\alpha,\beta} = \text{Min}\{i : \models \neg \varphi_i(c_\alpha c_\beta^{-1}, \bar{\mathbf{b}})\}$).

This finishes the proof of part (1). We still need to prove 2), 3).

2) Let $q'(x) \subseteq q(x)$ have cardinality $\leq |T|^{\aleph_0}$ and be such that $q(\mathfrak{C}) = (p \cup q')(\mathfrak{C})$; $q'(x)$ exists by part (1). Observe that every automorphism of $\mathfrak{C}$ fixing $\text{Dom}(p)$ maps $p(\mathfrak{C})$ onto itself and therefore maps $q(\mathfrak{C})$ onto itself.

It follows that if $c_1, c_2 \in p(\mathfrak{C})$ are such that $\text{tp}(c_1, \text{Dom}(p)) = \text{tp}(c_2, \text{Dom}(p))$, then $c_1 \in q(\mathfrak{C})$ if and only if $c_2 \in q(\mathfrak{C})$. Let $\mathbf{P} := \{\text{tp}(b, \text{Dom}(p)) : b \in q(\mathfrak{C})\}$, $\mathbf{P}(\mathfrak{C}) := \bigcup \{r(\mathfrak{C}) : r \in \mathbf{P}\}$. Then by the above explanation, $\mathbf{P}(\mathfrak{C}) \subseteq q(\mathfrak{C})$. By definition, $q(\mathfrak{C}) \subseteq p(\mathfrak{C})$, so they are equal. Let $q_{**} = \bigcap \{r : r \in \mathbf{P}\}$, so we have $q(\mathfrak{C}) \subseteq q_{**}(\mathfrak{C})$.

If they are equal, then we are done. Otherwise take $c_1 \in q_{**}(\mathfrak{C}) \setminus q(\mathfrak{C})$. Without loss of generality, let $\psi(x, \bar{d}) \in q$ be such that $\models \neg\psi(c_1, \bar{d})$.

By definition of $\mathbf{P}$ and c_1 , for each $\theta(x, \bar{e}) \in \text{tp}(c_1, \text{Dom}(p))$ there exists some $p_{\theta(x, \bar{e})} \in \mathbf{P}$ such that $\theta(x, \bar{e}) \in p_{\theta(x, \bar{e})}$ and therefore some $c_{\theta(x, \bar{e})} \in q(\mathfrak{C})$ satisfies $\theta(x, \bar{e})$. So $\text{tp}(c_1, \text{Dom}(p)) \cup q'(x)$ is finitely satisfiable and is therefore realized by some c_2 . Thus $\text{tp}(c_1, \text{Dom}(p)) = \text{tp}(c_2, \text{Dom}(p))$, but $c_1 \notin q'(\mathfrak{C}) = q(\mathfrak{C})$ and $c_2 \in (p \cup q')(\mathfrak{C})$, a contradiction.

3) By the proof of part (1). □_{1.1}

REFERENCES

- [BaSx76] John T. Baldwin and Jan Saxl. Logical stability in group theory. *J. Austral. Math. Soc. Ser. A*, **21**:267–276, 1976. MR0407151 (53:10934)
- [BOPP05] Alessandro Berarducci, Margarita Otero, Yaa'cov Peterzil, and Anand Pillay. A descending chain condition for groups definable in o-minimal structures. *Annals of Pure and Applied Logic*, **134**:303–313, 2005. MR2139910 (2006a:03052)
- [HPP0x] Ehud Hrushovski, Ya'acov Peterzil, and Anand Pillay. Groups, measures, and the NIP. *preprint*, 2005.
- [Sh:F753] Saharon Shelah. Definable Groups for Dependent Theories.
- [Sh:783] Saharon Shelah. Dependent first order theories, continued. *Israel Journal of Mathematics*, **accepted**. math.LO/0406440.
- [Sh:863] Saharon Shelah. Strongly dependent theories. *Israel Journal of Mathematics*, **submitted**. math.LO/0504197.
- [Sh:F705] Shelah, Saharon. Representation over orders of elementary classes.

THE HEBREW UNIVERSITY OF JERUSALEM, EINSTEIN INSTITUTE OF MATHEMATICS, EDMOND J. SAFRA CAMPUS, GIVAT RAM, JERUSALEM 91904, ISRAEL

Current address: Department of Mathematics, Hill Center-Busch Campus, Rutgers, The State University of New Jersey, 110 Frelinghuysen Road, Piscataway, New Jersey 08854-8019