A KHINCHIN SEQUENCE

THOMAS WIETING

(Communicated by Jonathan M. Borwein)

ABSTRACT. We prove that the geometric and harmonic means of the sequence Z_2 of positive integers proposed by Bailey, Borwein, and Crandall converge to the corresponding Khinchin Constants.

1. KHINCHIN SEQUENCES

One defines the *Khinchin Constant* K by the following relation:

$$\log(K) = \frac{1}{\log(2)} \sum_{k=1}^{\infty} \log(k) \log\left(\frac{(k+1)^2}{k(k+2)}\right) = \log(2.685452001 \dots).$$

For any sequence $A = (a_j)$,

$$A : a_1, a_2, a_3, \dots, a_j, \dots$$

of positive integers, let us refer to A as a *Khinchin Sequence* iff the geometric means of A converge to K :

$$\lim_{n \rightarrow \infty} \left(\prod_{j=1}^n a_j \right)^{1/n} = K.$$

That is,

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{j=1}^n \log(a_j) = \log(K).$$

For any irrational number x in the interval $(0, 1)$, let us refer to x as a *Khinchin Number* iff the continued fraction expansion $A(x) = (a_j(x))$

$$A(x) : a_1(x), a_2(x), a_3(x), \dots, a_j(x), \dots$$

for x is a Khinchin Sequence. In this paper, our objective is to prove that the particular sequence $C = (c_j)$

$$C : 2, 5, 1, 11, 1, 3, 1, 22, 2, 4, 1, 7, 1, 2, 1, 45, 2, 4, 1, 8, \dots, c_j, \dots$$

of positive integers proposed by Bailey, Borwein, and Crandall is a Khinchin Sequence. See reference [1].

Received by the editors January 12, 2007.

2000 *Mathematics Subject Classification.* Primary 11Y65; Secondary 28D05.

Key words and phrases. Khinchin Sequence, continued fraction expansion, geometric mean, harmonic mean.

Thanks to R. C. Crandall for suggesting the subject of this paper.

©2007 American Mathematical Society
 Reverts to public domain 28 years from publication

In the paper just cited, the authors denote C by Z_2 . They define the sequence C in terms of two auxiliary sequences $U = (u_j)$ and $V = (v_k)$, defined in turn as follows. The first sequence, U , is the *van der Corput Sequence*:

$$U: \quad \frac{1}{2}, \quad \frac{1}{4}, \frac{3}{4}, \quad \frac{1}{8}, \frac{5}{8}, \frac{3}{8}, \frac{7}{8}, \quad \frac{1}{16}, \frac{9}{16}, \frac{5}{16}, \frac{13}{16}, \frac{3}{16}, \frac{11}{16}, \frac{7}{16}, \frac{15}{16}, \quad \dots, u_j, \dots$$

Specifically, for each positive integer j , u_j is the dyadic rational number obtained by reflecting the binary representation of j in the binary point. For example, $u_{12} := 0.0011 = 3/16$ because $12 = 1100.0$. See reference [2]. The second sequence, V , describes a particular partition of the interval $(0, 1]$:

$$V: \quad \dots < v_k = \frac{1}{\log(2)} \log \left(\frac{k+1}{k} \right) < \dots < v_3 < v_2 < v_1 = 1.$$

Now, in terms of U and V , Bailey, Borwein, and Crandall define the sequence C as follows:

$$C: \quad c_j = k \iff v_{k+1} < u_j \leq v_k.$$

For example, $c_{12} = 7$ because $v_8 < u_{12} \leq v_7$.

2. MOTIVATION

To set a context for our study of the sequence C , let us describe a special case of the Ergodic Theorem. Let λ stand for Lebesgue Measure, defined as usual on $\mathbb{R}$. Let X be the set of all irrational numbers in the interval $(0, 1)$. Let μ stand for Gauss Measure, defined on X as follows:

$$\mu(E) := \frac{1}{\log(2)} \int_E \frac{1}{1+x} \lambda(dx)$$

where E is any Borel subset of X . Note that $\mu(E) = 0$ iff $\lambda(E) = 0$. Let F be the mapping carrying X to itself defined as follows:

$$F(x) := \frac{1}{x} - \left\lfloor \frac{1}{x} \right\rfloor$$

where x is any number in X . Of course, $\lfloor 1/x \rfloor$ denotes the largest among all integers ℓ for which $\ell \leq 1/x$. Note that F is continuous. One may view the ordered pair (X, F) as a (discrete) dynamical system. For any x in X , one may say that if the system is in state x at time 0, then the system is in state $F(x)$ one unit of time later. By an elementary argument, one can show that μ is invariant under F , in the sense that, for any Borel subset E of X , $\mu(F^{-1}(E)) = \mu(E)$. By a more substantial argument, one can also show that μ is ergodic under F , in the sense that, for any Borel subset E of X , if $F^{-1}(E) = E$, then either $\mu(E) = 0$ or $\mu(E) = 1$. See reference [4]. Let h be the function defined on X as follows:

$$h(x) := \left\lfloor \frac{1}{x} \right\rfloor$$

where x is any number in X . Note that h is continuous and that the values of h are positive integers. One may refer to h as an observable for the given dynamical system.

For any x in X , one obtains the orbit $O(x) = (x_j)$ of x under F ,

$$O(x): \quad x = x_1, x_2, x_3, \dots$$

and one obtains the corresponding (discrete) time sequence $A(x) = (a_j(x))$,

$$A(x) : \quad a_1(x), a_2(x), a_3(x), \dots,$$

where

$$x_j := F^{j-1}(x), \quad a_j(x) := h(x_j).$$

The sequence $A(x)$ is the *Continued Fraction Expansion* for x .

For the assembly X , μ , F , and $\log(h)$, the Ergodic Theorem states that, for almost every x in X ,

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{j=1}^n \log(h(F^{j-1}(x))) = \int_X \log(h(y)) \mu(dy).$$

That is, the time average of $\log(h)$ over $O(x)$ equals the space average of $\log(h)$ over X . See reference [5]. Hence,

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{j=1}^n \log(a_j(x)) &= \int_X \log(h(y)) \mu(dy) \\ &= \sum_{k=1}^{\infty} \log(k) \mu\left(\frac{1}{k+1}, \frac{1}{k}\right) \\ &= \frac{1}{\log(2)} \sum_{k=1}^{\infty} \log(k) \log\left(\frac{(k+1)^2}{k(k+2)}\right) \\ &= \log(K). \end{aligned}$$

Now it is plain that, for almost every irrational number x in the interval $(0, 1)$, x is a Khinchin Number. However, no particular examples of such numbers are known. The beguiling cases of $\pi - 3$ and even of $K - 2$ itself have been studied energetically but to no analytic decision as yet. In reference [3], one may find the optimistic opinion that $\pi - 3$ is a Khinchin Number. The graphs displayed in Figures 1 and 2 suggest a more cautious, though still hopeful, opinion on $\pi - 3$ and on $K - 2$ as well. The graphs are list plots of the following data:

$$\frac{1}{n} \sum_{j=1}^n \log(a_j(x)) - \log(K) \quad (1 \leq n \leq 4096)$$

where $x = \pi - 3$ and $x = K - 2$.

Failing to identify particular Khinchin Numbers, one naturally turns to the design of particular Khinchin Sequences. One might, for instance, design sequences $A = (a_j)$ such that, for each j , a_j equals 2 or 3 and such that the 2s and 3s occur in A in correct “limiting proportions,” specifically, the proportions p and q , where p and q are the positive numbers for which $p + q = 1$ and $\log(K) = p \log(2) + q \log(3)$. However, such a design would be very difficult to implement, since it depends upon the calculation of $\log(K)$ to arbitrary accuracy. In sharp contrast, Bailey, Borwein, and Crandall have proposed a particular candidate for a Khinchin Sequence, namely, the sequence C , which they have defined in constructive and rapidly computable terms. Let us prove formally that C is indeed a Khinchin Sequence.

FIGURE 1. $x = \pi - 3$

FIGURE 2. $x = K - 2$

3. THE FUNCTION ϕ

Let ϕ be the function defined on the interval $J = (0, 1]$ as follows. For each x in J and for any positive integer k ,

$$\phi(x) = \log(k) \iff v_{k+1} < x \leq v_k.$$

In particular, for each positive integer j , $\phi(u_j) = \log(c_j)$. See Figure 3. Clearly,

$$\begin{aligned} \int_J \phi(x) \lambda(dx) &= \frac{1}{\log(2)} \sum_{k=1}^\infty \log(k) \left(\log \left(\frac{k+1}{k} \right) - \log \left(\frac{k+2}{k+1} \right) \right) \\ &= \frac{1}{\log(2)} \sum_{k=1}^\infty \log(k) \log \left(\frac{(k+1)^2}{k(k+2)} \right) \\ &= \log(K). \end{aligned}$$

FIGURE 3. The function ϕ

Now it is plain that C is a Khinchin Sequence iff

$$(1) \quad \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{j=1}^n \phi(u_j) = \int_J \phi(x) \lambda(dx).$$

We proceed to prove relation (1).

4. INTEGRATING SEQUENCES

Let ψ be a real-valued Borel function defined on J and integrable with respect to λ . Let us say that the sequence U *integrates* ψ iff

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{j=1}^n \psi(u_j) = \int_J \psi(x) \lambda(dx).$$

To prove relation (1), we must prove that U integrates ϕ . Obviously, the functions integrated by U comprise a linear space. By an elementary argument, one can show that, for each subinterval I of J , U integrates the characteristic function χ_I of I . One summarizes this property of U by saying that U is *uniformly distributed* in J . We will prove this property in an appendix to this paper. Presuming the property, let us prove that U integrates ϕ . To that end, we require only that:

- (1) ϕ is nonnegative and decreasing;
- (2) for each positive integer p , U integrates the function

$$\phi_p := \chi_{[1/2^p, 1]} \phi.$$

Let n be any positive integer. Let α_n be the average of the values of ϕ at the first n terms of U :

$$\alpha_n := \frac{1}{n} \sum_{j=1}^n \phi(u_j).$$

Let

$$\beta := \int_J \phi(x) \lambda(dx).$$

We must prove that

$$\lim_{n \rightarrow \infty} \alpha_n = \beta.$$

Let p be any positive integer. Let ϕ_p be the function defined on J by truncation of ϕ , as follows:

$$\phi_p := \chi_{[1/2^p, 1]} \phi.$$

That is,

$$\phi_p(x) := \begin{cases} 0 & \text{if } 0 < x < 1/2^p, \\ \phi(x) & \text{if } 1/2^p \leq x \leq 1. \end{cases}$$

Obviously, for each x in J ,

$$\phi_1(x) \leq \phi_2(x) \leq \cdots \leq \phi_p(x) \leq \cdots \uparrow \phi(x).$$

Let $\alpha_{n,p}$ be the corresponding average of the values of ϕ_p at the first n terms of U :

$$\alpha_{n,p} := \frac{1}{n} \sum_{j=1}^n \phi_p(u_j).$$

Let

$$\beta_p := \int_J \phi_p(x) \lambda(dx).$$

Clearly, ϕ_p is a linear combination of characteristic functions of subintervals of J . By our initial remarks, it is plain that U integrates ϕ_p :

$$(2) \quad \lim_{n \rightarrow \infty} \alpha_{p,n} = \beta_p.$$

Now let ϵ be any positive real number. By the Monotone Convergence Theorem, we may introduce a positive integer p such that

$$(3) \quad \beta - \epsilon < \beta_p \leq \beta,$$

from which it follows that

$$(4) \quad \int_{(0, 1/2^p)} \phi(x) \lambda(dx) < \epsilon.$$

By relation (2), we may introduce a positive integer m such that, for every positive integer n , if $m \leq n$, then

$$(5) \quad \beta_p - \epsilon < \alpha_{n,p} < \beta_p + \epsilon.$$

We may as well arrange that $2^p \leq m$. Let n be any positive integer for which $m \leq n$. Let q be the positive integer for which $2^{q-1} - 1 < n \leq 2^q - 1$. Clearly, $p < q$. One may say that the first n terms of U have run through the first $q-1$ “cycles” of U and have at least begun (perhaps even finished) the q -th cycle. The smallest term of the q -th cycle is $1/2^q$. Hence, for each positive integer j , if $1 \leq j \leq n$, then $1/2^q \leq u_j$. Consequently,

$$(6) \quad \alpha_{n,q} = \alpha_n.$$

Now let

$$t_1, t_2, \dots, t_\ell \quad (\ell = 2^{q-p} - 1)$$

be the terms among

$$u_1, u_2, \dots, u_r \quad (r = 2^q - 1)$$

FIGURE 4. Comparison of areas

that are less than $1/2^p$. Figure 4, $p = 2$, $q = 4$, and $\ell = 3$. Since ϕ is nonnegative and decreasing, we find that

$$\begin{aligned}
 \alpha_{n,q} - \alpha_{n,p} &\leq \frac{1}{n} \sum_{j=1}^{\ell} \phi(t_j) \\
 &= \frac{2^q}{n} \frac{1}{2^q} \sum_{j=1}^{\ell} \phi(t_j) \\
 &\leq 4 \int_{(0,1/2^p)} \phi(x) \lambda(dx) && (\text{since } 2 \cdot 2^{q-1} < 2(n+1)) \\
 &< 4\epsilon && (\text{by relation (4)}).
 \end{aligned}$$

Hence, by relations (3) and (5) and by the foregoing computation,

$$\beta - 2\epsilon < \beta_p - \epsilon < \alpha_{n,p} \leq \alpha_{n,q} < \alpha_{n,p} + 4\epsilon < \beta_p + 5\epsilon \leq \beta + 5\epsilon$$

so that, by relation (6), $\beta - 2\epsilon < \alpha_n < \beta + 5\epsilon$. Therefore,

$$(1) \quad \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{j=1}^n \phi(u_j) = \int_J \phi(x) \lambda(dv).$$

5. QUESTIONS

The number x in $(0, 1)$ of which C is the continued fraction expansion is approximately equal to 0.46107049595671951935. Of course, it is a Khinchin Number. Can one identify x in “familiar” terms?

FIGURE 5. $N = 4096$ FIGURE 6. $N = 8192$

In Figures 5 and 6, we display list plots of the following data:

$$\frac{1}{n} \sum_{j=1}^n \log(c_j) - \log(K) \quad (1 \leq n \leq N),$$

where $N = 4096$ and $N = 8192$. Can one explain, in formally precise terms, the apparent self-similarity of the data?

6. HARMONIC MEANS

Let r be any real number for which $r < 1$ and $r \neq 0$. With reference to Section 4, let us define the function ϕ_r on J as follows. For each x in J and for any positive integer k ,

$$\phi_r(x) = k^r \iff v_{k+1} < x \leq v_k.$$

In particular, for each positive integer j , $\phi_r(u_j) = c_j^r$. If $r < 0$, then $1 - \phi_r$ is similar to ϕ , in the sense that it meets the conditions (1) and (2) stated in Section 4. If

$0 < r < 1$, then ϕ_r itself is similar to ϕ . In either case, U integrates ϕ_r . Hence,

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{j=1}^n c_j^r &= \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{j=1}^n \phi_r(u_j) \\ &= \int_J \phi_r(x) \lambda(dx) \\ &= \frac{1}{\log(2)} \sum_{k=1}^{\infty} k^r \log \left(\frac{(k+1)^2}{k(k+2)} \right). \end{aligned}$$

One defines the *Khinchin Constant* K_r by the following relation:

$$K_r = \frac{1}{\log(2)} \sum_{k=1}^{\infty} k^r \log \left(\frac{(k+1)^2}{k(k+2)} \right).$$

We infer that the r -th harmonic means of C converge to K_r :

$$\lim_{n \rightarrow \infty} \left(\frac{1}{n} \sum_{j=1}^n c_j^r \right)^{1/r} = K_r.$$

7. APPENDIX

The van der Corput Sequence U falls into cycles:

$$U : \quad \frac{1}{2}, \quad \frac{1}{4}, \frac{3}{4}, \quad \frac{1}{8}, \frac{5}{8}, \frac{3}{8}, \frac{7}{8}, \quad \frac{1}{16}, \frac{9}{16}, \frac{5}{16}, \frac{13}{16}, \frac{3}{16}, \frac{11}{16}, \frac{7}{16}, \frac{15}{16}, \quad \dots, u_j, \dots$$

For each positive integer p , the first term of the p -th cycle is $1/2^p$ and the length of the p -th cycle is 2^{p-1} . The sum of the lengths of the first p cycles is $2^p - 1$. Moreover,

$$u_{2^p+j} = \frac{1}{2^{p+1}} + u_j \quad (0 < j < 2^p).$$

By these observations, it is plain that, for any dyadic interval I of the form

$$I = [j/2^p, (j+1)/2^p) \quad (p \in \mathbb{Z}^+, 0 < j < 2^p),$$

the sequence U visits I precisely once in the course of its first p cycles. Thereafter, it visits I periodically with period 2^p . Hence, for any positive integer n , if $2^p \leq n$, then

$$\frac{m}{n} \leq \frac{1}{n} \sum_{j=1}^n \chi_I(u_j) \leq \frac{m+1}{n},$$

where m is the positive integer for which

$$m2^p - 1 < n \leq (m+1)2^p - 1.$$

Consequently,

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{j=1}^n \chi_I(u_j) = \frac{1}{2^p} = \lambda(I),$$

which is to say that U integrates χ_I .

In turn, for any subinterval I of the interval $(0, 1)$ and for any positive real number ϵ , we may introduce finite disjoint unions I' and I'' of dyadic intervals of the foregoing form such that $I' \subseteq I \subseteq I''$ and $\lambda(I'' \setminus I') < \epsilon$. Hence,

$$\begin{aligned} \limsup_{n \rightarrow \infty} \frac{1}{n} \sum_{j=1}^n \chi_I(u_j) &\leq \lim_{j \rightarrow \infty} \frac{1}{n} \sum_{j=1}^n \chi_{I''}(u_j) \\ &= \lambda(I'') \\ &< \lambda(I') + \epsilon \\ &= \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{j=1}^n \chi_{I'}(u_j) + \epsilon \\ &\leq \liminf_{n \rightarrow \infty} \frac{1}{n} \sum_{j=1}^n \chi_I(u_j) + \epsilon. \end{aligned}$$

Consequently,

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{j=1}^n \chi_I(u_j) = \lambda(I),$$

which is to say that U integrates χ_I .

REFERENCES

- [1] D. H. Bailey, J. M. Borwein, R. C. Crandall, *On the Khinchin Constant*, Math. Comp. 66 (1997), 417-431. MR1377659 (97c:11119)
- [2] J. G. van der Corput, *Verteilungsfunktionen*, Proc. Ned. Akad. v. Wet. 38 (1935), 813-821.
- [3] D. H. Lehmer, *Note on an Absolute Constant of Khinchin*, Amer. Math. Monthly, 46 (1939), 148-152. MR1524526
- [4] C. Ryll-Nardzewski, *On the Ergodic Theorems (I, II)*, Studia Math. 12 (1951), 65-79. MR0046582 (13:757a); MR0046583 (13:757b)
- [5] P. Walters, *An Introduction to Ergodic Theory*, Springer-Verlag, New York, 1982. MR648108 (84e:28017)

DEPARTMENT OF MATHEMATICS, REED COLLEGE, PORTLAND, OREGON 97202
E-mail address: **wieting@reed.edu**