ON THE COMBINATORICS OF CRYSTAL GRAPHS, II. THE CRYSTAL COMMUTOR

CRISTIAN LENART

(Communicated by Jim Haglund)

ABSTRACT. We present an explicit combinatorial realization of the commutor in the category of crystals which was first studied by Henriques and Kamnitzer. Our realization is based on certain local moves defined by van Leeuwen.

1. INTRODUCTION

We work in the category **g-Crystals** of crystals corresponding to representations of complex semisimple Lie algebras **g**. It is well-known that this is a *monoidal category* with an associative tensor product (e.g., see [6]). The crystals $A \otimes B$ and $B \otimes A$ are isomorphic via maps called *commutors*. The map

$$\text{flip} : A \otimes B \rightarrow B \otimes A, \quad (a, b) \mapsto (b, a)$$

is not a commutor. Henriques and Kamnitzer [6], based on an idea of Berenstein, defined a commutor $\sigma_{A,B}$ based on Lusztig's involution on a crystal. They also proved that the category **g-Crystals** with this commutor is a *coboundary category* (cf. [4]). More recently, Kamnitzer and Tingley [9] proved that the action of $\sigma_{A,B}$ on the highest weight elements (which determines it) is given by *Kashiwara's involution* on the Verma crystal; as remarked in [9], Kashiwara's involution can be realized in terms of *Mirković-Vilonen polytopes* [8]. Both of the above constructions of $\sigma_{A,B}$ depend on some maps of crystals whose explicit construction is nontrivial. In this paper, we present an explicit realization of the commutor $\sigma_{A,B}$ based on some local moves defined by van Leeuwen [22]; the latter generalize the realization of Schützenberger's *jeu de taquin* for Young tableaux (e.g., see [5]) via Fomin's *growth diagrams* [20, Appendix 1]. Our realization is proved to hold, in particular, for all simple Lie algebras **g** with the exception of those of type E_8 , F_4 , and G_2 ; moreover, it is conjectured to hold for all semisimple Lie algebras.

2. BACKGROUND ON CRYSTALS

In this section, we recall some background information on finite root systems and crystal graphs.

Received by the editors February 16, 2007.

2000 *Mathematics Subject Classification*. Primary 20G42; Secondary 17B10, 22E46.

Key words and phrases. Crystals, root operators, coboundary category, commutor, Lusztig's involution, van Leeuwen's jeu de taquin.

The author was supported by National Science Foundation grant DMS-0403029.

©2007 American Mathematical Society
 Reverts to public domain 28 years from publication

2.1. Root systems. Let $\mathfrak{g}$ be a complex semisimple Lie algebra with Cartan subalgebra $\mathfrak{h}$. Let $\Phi \subset \mathfrak{h}^*$ be its finite *root system* with positive roots Φ^+ , and let $\mathfrak{h}_{\mathbb{R}}^* \subset \mathfrak{h}^*$ be the real span of Φ . Let r be the rank of $\mathfrak{h}$, and let $\alpha_i \in \Phi^+$ for $i \in I := \{1, \dots, r\}$ be the corresponding *simple roots*. The nondegenerate scalar product on $\mathfrak{h}_{\mathbb{R}}^*$ induced by the Killing form is denoted by $\langle \cdot, \cdot \rangle$. Given a root α , we have the corresponding *coroot* $\alpha^\vee := 2\alpha/\langle \alpha, \alpha \rangle$. The reflection corresponding to a root α is denoted, as usual, by s_α , and we have the *simple reflections* $s_i := s_{\alpha_i}$ for $i \in I$. Let $W \subset \text{Aut}(\mathfrak{h}_{\mathbb{R}}^*)$ be the Weyl group of Φ . The *length* of an element $w \in W$ is denoted by $\ell(w)$. Let w_o be the longest element of the Weyl group (i.e., $\ell(w_o) = |\Phi^+|$).

We denote the *weight lattice* by Λ , the subset of *dominant weights* by Λ^+ , and the *fundamental weights* by $\Lambda_1, \dots, \Lambda_r$. For each dominant weight λ , there is a finite dimensional irreducible representation of $\mathfrak{g}$ with highest weight λ , which is denoted by V_λ . Let $\mathbb{Z}[\Lambda]$ be the group algebra of the weight lattice Λ , which has a $\mathbb{Z}$ -basis of formal exponents $\{e^\lambda : \lambda \in \Lambda\}$ and multiplication $e^\lambda \cdot e^\mu := e^{\lambda+\mu}$; in other words, $\mathbb{Z}[\Lambda] = \mathbb{Z}[e^{\pm\Lambda_1}, \dots, e^{\pm\Lambda_r}]$ is the algebra of Laurent polynomials in r variables. The formal characters of the modules V_λ are given by $ch(V_\lambda) = \sum_{\mu \in \Lambda} m_\lambda(\mu) e^\mu \in \mathbb{Z}[\Lambda]$, where $m_\lambda(\mu)$ is the multiplicity of the weight μ in V_λ .

A weight λ is called *minuscule* if $\langle \lambda, \alpha^\vee \rangle \in \{0, \pm 1\}$ for all $\alpha \in \Phi$, and *quasi-minuscule* if $\langle \lambda, \alpha^\vee \rangle \in \{0, \pm 1, \pm 2\}$ for all $\alpha \in \Phi$. Since W permutes Φ , every element in the W -orbit of a minuscule (or quasi-minuscule) weight is also minuscule (respectively quasi-minuscule). The irreducible representations corresponding to the dominant minuscule weights are called *minuscule representations*, and are characterized by the fact that their weights form a single W -orbit (i.e., the extremal weights, of multiplicity 1). The dominant minuscule weights have a well-known classification (e.g., see [3, Exercise VI.4.15]). There are minuscule weights for all Cartan-Killing types with the exception of types E_8 , F_4 , and G_2 .

We will need the following result in [21].

Theorem 2.1 ([21]). *If Φ is irreducible, then for every dominant weight λ there is a decomposition $\lambda = \lambda_1 + \dots + \lambda_n$ such that λ_i is minuscule or quasi-minuscule, and $\lambda_1 + \dots + \lambda_i$ is dominant for all $i = 1, \dots, n$. In fact, all λ_i can be chosen to be minuscule if Φ is not of type E_8 , F_4 , and G_2 . Furthermore, all λ_i can be chosen in the orbit of a single minuscule or quasi-minuscule weight unless Φ is of type D_n for n even.*

2.2. Crystals. This section follows [21, Section 2] and [6, Section 2]. We refer to these papers for more details.

Kashiwara [10, 11] defined a directed colored graph, called *crystal graph*, on the *canonical basis* of a representation of the quantum group $U_q(\mathfrak{g})$ [10, 18]. This graph partially encodes the action of the Chevalley generators of $U_q(\mathfrak{g})$ on the canonical basis. We will now define axiomatically the category $\mathfrak{g}\text{-Crystals}$ of crystals corresponding to complex semisimple Lie algebras $\mathfrak{g}$.

The objects of the category $\mathfrak{g}\text{-Crystals}$, which are called *crystals* (or *$\mathfrak{g}$ -crystals*), are 4-tuples $(B, \text{wt}, \varepsilon, \{\tilde{f}_1, \dots, \tilde{f}_r\})$, where

- B is the underlying set of the corresponding crystal;
- wt and ε are maps $B \rightarrow \Lambda$;
- $\tilde{f}_i$ are maps from B to $B \sqcup \{0\}$.

For each $b \in B$, we call $\text{wt}(b)$, $\varepsilon(b)$, and $\phi(b) := \text{wt}(b) + \varepsilon(b)$ the *weight*, *depth*, and *rise* of b . A morphism of crystals is a map of the underlying sets that commutes with all the structure maps.

By a result of Joseph [7], the category **g-Crystals** is uniquely defined by the axioms below.

$$(A1) \quad \varepsilon(b), \phi(b) \in \Lambda^+.$$

We define the depth and rise in the direction of the simple root α_i by $\varepsilon_i(b) := \langle \varepsilon(b), \alpha_i^\vee \rangle$ and $\phi_i(b) := \langle \phi(b), \alpha_i^\vee \rangle$. We also let $\text{wt}_i(b) := \langle \text{wt}(b), \alpha_i^\vee \rangle$.

$$(A2) \quad \tilde{f}_i \text{ is a bijection from } \{b \in B : \phi_i(b) > 0\} \text{ to } \{b \in B : \varepsilon_i(b) > 0\}.$$

We let $\tilde{e}_i := \tilde{f}_i^{-1}$ denote the inverse map, and extend it to a map from B to $B \sqcup \{0\}$ by defining it to be 0 on $\{b \in B : \varepsilon_i(b) = 0\}$.

(A3) We have $\text{wt}(\tilde{f}_i(b)) = \text{wt}(b) - \alpha_i$, and $\varepsilon_i(\tilde{f}_i(b)) = \varepsilon_i(b) + 1$, whenever $\tilde{f}_i(b) \neq 0$.

Hence, we also have $\phi_i(\tilde{f}_i(b)) = \phi_i(b) - 1$. The maps $\tilde{e}_i$ and $\tilde{f}_i$, called *root operators*, act as raising and lowering operators which provide a partition of B into α_i -strings that are closed under the action of $\tilde{e}_i$ and $\tilde{f}_i$. For example, the α_i -string through b is (by definition)

$$\tilde{f}_i^\phi(b), \dots, \tilde{f}_i(b), b, \tilde{e}_i(b), \dots, \tilde{e}_i^\varepsilon(b),$$

where $\varepsilon := \varepsilon_i(b)$ and $\phi := \phi_i(b)$; furthermore, we have $\tilde{f}_i^{\phi+1}(b) = \tilde{e}_i^{\varepsilon+1}(b) = 0$.

We define partial orders on B , one for each $i \in I$, by

$$(2.1) \quad c \preceq_i b \quad \text{if} \quad c = \tilde{f}_i^k(b) \quad \text{for some} \quad k \geq 0.$$

Let $\preceq$ denote the partial order on B generated by all partial orders $\preceq_i$, for $i = 1, \dots, r$. The set of maximal elements of the poset $(B, \preceq)$ is denoted $\max B$.

The *direct sum* of two crystals is defined by the disjoint union of their underlying sets.

(A4) For each dominant weight λ , the category contains an object B_λ such that the poset $(B_\lambda, \preceq)$ has a maximum (i.e., a highest weight element), denoted b_λ , and $\text{wt}(b_\lambda) = \lambda$. Furthermore, the category consists of all crystals isomorphic to a direct sum of crystals B_λ .

The crystal B_λ is called a *highest weight crystal of highest weight λ* , and b_λ is called its highest weight element.

The *tensor product* of two crystals is the crystal corresponding to the tensor product of the corresponding representations; the explicit definition is given in [11]. By iterating this construction, we can define the n -fold tensor product of crystals $A_1 \otimes \dots \otimes A_n$ as follows:

$$(2.2) \quad \begin{aligned} \text{wt}(a_1, \dots, a_n) &:= \text{wt}(a_1) + \dots + \text{wt}(a_n), \\ \varepsilon_i(a_1, \dots, a_n) &:= \max_{1 \leq j \leq n} \varepsilon_i(a_j) - \text{wt}_i(a_1) - \dots - \text{wt}_i(a_{j-1}), \end{aligned}$$

$$(2.3) \quad \phi_i(a_1, \dots, a_n) := \max_{1 \leq k \leq n} \phi_i(a_k) + \text{wt}_i(a_{k+1}) + \dots + \text{wt}_i(a_n).$$

Furthermore, the root operators are defined by

$$\begin{aligned}\tilde{e}_i(a_1, \dots, a_n) &:= (a_1, \dots, \tilde{e}_i(a_j), \dots, a_n), \\ \tilde{f}_i(a_1, \dots, a_n) &:= (a_1, \dots, \tilde{f}_i(a_k), \dots, a_n),\end{aligned}$$

where j and k are the smallest and largest indices for which the maximum is achieved in (2.2) (or, equivalently, in (2.3)).

(A5) For all dominant weights λ, μ , there exists an inclusion of crystals $\iota_{\lambda, \mu} : B_{\lambda+\mu} \hookrightarrow B_\lambda \otimes B_\mu$.

The category $\mathfrak{g}\text{-Crystals}$ is closed under tensor products. Thus, we have

$$(2.4) \quad B_{\lambda_1} \otimes \dots \otimes B_{\lambda_n} \simeq \bigoplus_{(b_1, \dots, b_n) \in \max B_{\lambda_1} \otimes \dots \otimes B_{\lambda_n}} B_{\text{wt}(b_1) + \dots + \text{wt}(b_n)},$$

where $\simeq$ denotes isomorphism of crystals. By (2.2), we have

$$(2.5) \quad \begin{aligned} &\max B_{\lambda_1} \otimes \dots \otimes B_{\lambda_n} \\ &= \{(b_1, \dots, b_n) : \text{wt}(b_1) + \dots + \text{wt}(b_{j-1}) - \varepsilon(b_j) \in \Lambda^+, j = 1, \dots, n\}. \end{aligned}$$

Remark 2.2. (1) If μ is a minuscule weight, then all the summands B_ν in $B_\lambda \otimes B_\mu$ correspond to $\nu = \lambda + \bar{\mu}$ for $\bar{\mu} \in W\mu$, and appear with multiplicity one.

(2) If μ is a quasi-minuscule weight, then all the summands B_ν in $B_\lambda \otimes B_\mu$ correspond to either $\nu = \lambda$, or $\nu = \lambda + \bar{\mu}$ for $\bar{\mu} \in W\mu$; the summands of the second type appear with multiplicity one.

The above axioms imply that any highest weight crystal B_λ has a minimum c_λ with $\text{wt}(c_\lambda) = w_\circ(\lambda)$. Let $i \mapsto i^*$, for $i \in I$, be the Dynkin diagram automorphism specified by $\alpha_{i^*} = -w_\circ(\alpha_i)$. There is an involution $\eta_{B_\lambda} = \eta_\lambda$ on B_λ specified by the following conditions:

$$(2.6) \quad \eta_\lambda(b_\lambda) = c_\lambda, \quad \tilde{e}_i \eta_\lambda = \eta_\lambda \tilde{f}_{i^*},$$

where $\eta_\lambda(0) = 0$. We can define an involution η_B on any crystal B by applying η_λ to each component of B isomorphic to B_λ . This involution is due to Lusztig, and corresponds to the action of $w_\circ$ on a representation of a quantum group (e.g., see [1, 19]). Thus, we have

$$(2.7) \quad \text{wt}(\eta_B(b)) = w_\circ(\text{wt}(b)),$$

which also follows from the axioms.

2.3. Models for crystals. There are several models for crystals corresponding to semisimple Lie algebras, such as: Kashiwara-Nakashima tableaux [12], Littelmann paths [16, 17], the alcove path model [13, 14, 15], the model in [2] based on Lusztig's parametrization of canonical bases, as well as some models based on geometric constructions. Instead of using a particular model, here we will use the construction of crystals based on their embedding into tensor products of minuscule or quasi-minuscule crystals (i.e., crystals B_λ with λ minuscule or quasi-minuscule weights). We now describe this construction, following [21].

Let us first describe minuscule crystals. If λ is minuscule, then the underlying set of B_λ is the W -orbit $W\lambda$. Naturally, the weight of $\mu \in B_\lambda$ is μ itself, and we

define

$$\varepsilon_i(\mu) := \max(0, -\langle \mu, \alpha_i^\vee \rangle) = \begin{cases} 1 & \text{if } \langle \mu, \alpha_i^\vee \rangle = -1, \\ 0 & \text{otherwise,} \end{cases}$$

$$\phi_i(\mu) := \max(0, \langle \mu, \alpha_i^\vee \rangle) = \begin{cases} 1 & \text{if } \langle \mu, \alpha_i^\vee \rangle = 1, \\ 0 & \text{otherwise.} \end{cases}$$

The root operators are defined as follows, for any $\mu \in B_\lambda$:

$$\begin{aligned} \tilde{f}_i(\mu) &:= s_i(\mu) = \mu - \alpha_i && \text{if } \langle \mu, \alpha_i^\vee \rangle = 1, \\ \tilde{e}_i(\mu) &:= s_i(\mu) = \mu + \alpha_i && \text{if } \langle \mu, \alpha_i^\vee \rangle = -1. \end{aligned}$$

In all other cases, the operators are defined to be 0.

For the description of a quasi-minuscule crystal, we refer to [21].

Now let λ be any dominant weight. By Theorem 2.1, there is a decomposition $\lambda = \lambda_1 + \dots + \lambda_n$ such that λ_i is minuscule or quasi-minuscule, and $\lambda_1 + \dots + \lambda_i$ is dominant for all $i = 1, \dots, n$. Let $\omega_i := \text{dom}_W(\lambda_i)$ denote the dominant representative of the W -orbit $W\lambda_i$. Then, by axiom (A5) and the construction of n -fold tensor products in Section 2.2, we have an embedding

$$(2.8) \quad B_\lambda \hookrightarrow B_{\omega_1} \otimes \dots \otimes B_{\omega_n}.$$

Indeed, by (2.2), the condition that all partial sums $\lambda_1 + \dots + \lambda_i$ are dominant weights guarantees that $(\lambda_1, \dots, \lambda_n)$ is a maximal element in $B_{\omega_1} \otimes \dots \otimes B_{\omega_n}$; thus, we can map $b_\lambda \mapsto (\lambda_1, \dots, \lambda_n)$, and extend this to an embedding (2.8) by using lowering operators.

2.4. A commutor for crystals. This section follows [6, Section 3], and we refer to this paper for more details.

Henriques and Kamnitzer [6], based on an idea of Berenstein, defined a commutor $\sigma_{A,B}$ (i.e., an isomorphism between the crystals $A \otimes B$ and $B \otimes A$) as follows:

$$\sigma_{A,B}(a, b) := \eta_{B \otimes A}(\eta_B(b), \eta_A(a)).$$

In other words, we have

$$(2.9) \quad \sigma_{A,B} = \eta_{B \otimes A} \circ (\eta_B \otimes \eta_A) \circ \text{flip}.$$

It turns out that we also have

$$(2.10) \quad \sigma_{A,B} = \text{flip} \circ \eta_{A \otimes B} \circ (\eta_A \otimes \eta_B).$$

More recently, Kamnitzer and Tingley [9] proved that the action of the above commutor on the highest weight elements (which determines it) is given by Kashiwara's involution on the Verma crystal [11].

Henriques and Kamnitzer proved that the category **g-Crystals** with this commutor is a coboundary category (cf. [4]). This statement amounts to the following three properties:

- (C1) $\sigma_{A,B}$ is an isomorphism of crystals, and is natural in A and B (i.e., it commutes with the maps on $A \otimes B$ and $B \otimes A$ coming from maps of crystals $A \rightarrow C$ and $B \rightarrow D$);
- (C2) $\sigma_{A,B} \circ \sigma_{B,A} = 1$;

(C3) the following diagram commutes:

$$(2.11) \quad \begin{array}{ccc} A \otimes B \otimes C & \xrightarrow{1 \otimes \sigma_{B,C}} & A \otimes C \otimes B \\ \sigma_{A,B} \otimes 1 \downarrow & & \downarrow \sigma_{A,C \otimes B} \\ B \otimes A \otimes C & \xrightarrow{\sigma_{B \otimes A,C}} & C \otimes B \otimes A \end{array}$$

Note that the third condition is an analog of the hexagon axiom for braided categories. By analogy with braid groups acting on multiple tensor products in braided categories, there is a group, denoted by J_n and called the *n-fruit cactus group*, which acts on n -fold tensor products in a coboundary category. This group is generated by $s_{p,q}$, for $1 \leq p < q \leq n$, subject to the following relations:

$$(R1) \quad s_{p,q}^2 = 1;$$

$$(R2) \quad s_{p,q}s_{k,l} = s_{k,l}s_{p,q} \text{ if } p < q, k < l, \text{ and either } q < k \text{ or } l < p;$$

$$(R3) \quad s_{p,q}s_{k,l} = s_{i,j}s_{p,q} \text{ if } p \leq k < l \leq q, \text{ where } i = \widehat{s}_{p,q}(l), j = \widehat{s}_{p,q}(k),$$

and $\widehat{s}_{p,q}$ denotes the following involution in the symmetric group S_n :

$$\widehat{s}_{p,q} = \begin{pmatrix} 1 & \cdots & p-1 & p & \cdots & q & q+1 & \cdots & n \\ 1 & \cdots & p-1 & q & \cdots & p & q+1 & \cdots & n \end{pmatrix}.$$

In the general context of coboundary categories, it is shown in [6, Lemmas 3-4] that the group J_n acts on n -fold tensor products $A_1 \dots A_n := A_1 \otimes \dots \otimes A_n$ by letting its generators “reverse intervals”:

$$\begin{aligned} s_{p,q} : A_1 \cdots A_{p-1} A_p A_{p+1} \cdots A_{q-1} A_q A_{q+1} \cdots A_n \\ \longrightarrow A_1 \cdots A_{p-1} A_q A_{q+1} \cdots A_{p+1} A_p A_{p+1} \cdots A_n. \end{aligned}$$

More precisely, based on the commutor, we first define natural isomorphisms denoted $\sigma_{p,r,q}$ for $1 \leq p \leq r < q \leq n$ by

$$\begin{aligned} (\sigma_{p,r,q})_{A_1, \dots, A_n} &:= 1 \otimes \sigma_{A_p \cdots A_r, A_{r+1} \cdots A_q} \otimes 1 : \\ A_1 \cdots A_{p-1} A_p \cdots A_r A_{r+1} \cdots A_q A_{q+1} \cdots A_n \\ &\longrightarrow A_1 \cdots A_{p-1} A_{r+1} \cdots A_q A_p \cdots A_r A_{q+1} \cdots A_n. \end{aligned}$$

We then define the action of $s_{p,q}$ recursively by $s_{p,p+1} := \sigma_{p,p,p+1}$, and $s_{p,q} := \sigma_{p,p,q} \circ s_{p+1,q}$ for $q - p > 1$. The following expression of $\sigma_{p,r,q}$ in terms of the generators of J_n will be needed:

$$(2.12) \quad \sigma_{p,r,q} = s_{p,q} \circ s_{r+1,q} \circ s_{p,r}.$$

3. VAN LEEUWEN’S JEU DE TAQUIN

In this section, we translate the construction of van Leeuwen in [22] into the language of crystal graphs. We start with the following important remark related to commutators.

Remark 3.1. In order to define a commutor, i.e., an isomorphism between the crystals $B_{\pi'} \otimes B_{\pi}$ and $B_{\pi} \otimes B_{\pi'}$ (for $\pi', \pi \in \Lambda^+$), it suffices to construct a weight-preserving bijection between $\max B_{\pi'} \otimes B_{\pi}$ and $\max B_{\pi} \otimes B_{\pi'}$. Clearly, such a bijection can be uniquely extended to a commutor via the action of the lowering operators.

A bijection of this type was constructed by van Leeuwen in [22], and is described below using the above setup.

We start by embedding the crystals $B_{\pi'}$ and B_{π} into tensor products of minuscule or quasi-minuscule crystals, as in (2.8):

$$(3.1) \quad B_{\pi'} \hookrightarrow B_{\omega'_1} \otimes \dots \otimes B_{\omega'_l}, \quad B_{\pi} \hookrightarrow B_{\omega_1} \otimes \dots \otimes B_{\omega_k};$$

these embeddings are based on decompositions $\pi' = \pi'_1 + \dots + \pi'_l$ and $\pi = \pi_1 + \dots + \pi_k$, i.e., we have $\omega'_j := \text{dom}_W(\pi'_j)$ and $\omega_i := \text{dom}_W(\pi_i)$. From now on, we will identify the crystals $B_{\pi'}$ and B_{π} with their embeddings in the corresponding tensor products.

Remark 3.2. The above embedding is compatible with the commutor of Henriques and Kamnitzer due to the naturality of the latter; cf. its property (C1).

Consider an element (cf. (2.5))

$$(3.2) \quad (b_{\pi'}, p) = ((\pi'_1, \dots, \pi'_l), (p_1, \dots, p_k)) \in \max B_{\pi'} \otimes B_{\pi},$$

which means that $p_i \in B_{\omega_i}$ for $i = 1, \dots, k$. It will be mapped to an element

$$(3.3) \quad (b_{\pi}, p') = ((\pi_1, \dots, \pi_k), (p'_1, \dots, p'_l)) \in \max B_{\pi} \otimes B_{\pi'},$$

which means that $p'_j \in B_{\omega'_j}$ for $j = 1, \dots, l$. This map is weight-preserving, so we have

$$(3.4) \quad \pi' + \text{wt}(p_1) + \dots + \text{wt}(p_k) = \pi + \text{wt}(p'_1) + \dots + \text{wt}(p'_l).$$

The idea is to use certain local moves which generalize the realization of Schützenberger's *jeu de taquin* for Young tableaux (e.g., see [5]) via Fomin's growth diagrams [20, Appendix 1].

We will define two matrices of elements in minuscule or quasi-minuscule crystals:

$$\begin{aligned} h_{i,j} &\in B_{\omega'_{j+1}}, \quad \text{for } i = 0, \dots, k \text{ and } j = 0, \dots, l-1, \\ v_{i,j} &\in B_{\omega_{i+1}}, \quad \text{for } i = 0, \dots, k-1 \text{ and } j = 0, \dots, l. \end{aligned}$$

They are related by

$$(3.5) \quad \text{wt}(h_{i,j}) + \text{wt}(v_{i,j+1}) = \text{wt}(v_{i,j}) + \text{wt}(h_{i+1,j}),$$

for $i = 0, \dots, k-1$ and $j = 0, \dots, l-1$.

These elements should be thought of as labels for the horizontal and vertical segments joining successive points in a square lattice which lie in the interior or on the rectangle with vertices $(0,0)$ and (k,l) . Here we use the matrix notation, with k indicating the row, and l the column. More precisely, $h_{i,j}$ and $v_{i,j}$ label the horizontal and vertical segments for which (i,j) is the left endpoint, respectively, the bottom endpoint. Condition (3.5) guarantees that the sum of weights of elements on each minimum length path between $(0,0)$ and (i,j) is the same, where $0 \leq i \leq k$ and $0 \leq j \leq l$. Let $\lambda^{[i,j]}$ be this sum (by default, we set $\lambda^{[0,0]} := 0$). We also require the following conditions:

$$(3.6) \quad (b_{\lambda^{[i,j]}}, h_{i,j}) \in \max B_{\lambda^{[i,j]}} \otimes B_{\omega'_{j+1}}, \quad \text{for } i = 0, \dots, k \text{ and } j = 0, \dots, l-1,$$

$$(3.7) \quad (b_{\lambda^{[i,j]}}, v_{i,j}) \in \max B_{\lambda^{[i,j]}} \otimes B_{\omega_{i+1}}, \quad \text{for } i = 0, \dots, k-1 \text{ and } j = 0, \dots, l.$$

We start by setting

$$(3.8) \quad h_{0,j} := \pi'_{j+1}, \quad \text{for } j = 0, \dots, l-1, \quad \text{and} \quad v_{i,l} := p_{i+1}, \quad \text{for } i = 0, \dots, k-1.$$

The other elements $h_{i,j}$ and $v_{i,j}$ are defined by local moves, which are bijections

$$\max B_{\lambda^{[i,j]}} \otimes B_{\omega'_{j+1}} \otimes B_{\omega_{i+1}} \longleftrightarrow \max B_{\lambda^{[i,j]}} \otimes B_{\omega_{i+1}} \otimes B_{\omega'_{j+1}},$$

for $i = 0, \dots, k-1$ and $j = 0, \dots, l-1$. More precisely, we have

$$(3.9) \quad (b_{\lambda^{[i,j]}}, h_{i,j}, v_{i,j+1}) \mapsto (b_{\lambda^{[i,j]}}, v_{i,j}, h_{i+1,j}).$$

The procedure ends by reading off the elements in (3.3) as follows:

$$(3.10) \quad p'_{j+1} := h_{k,j}, \quad \text{for } j = 0, \dots, l-1.$$

We now describe the local moves (3.9) as weight-preserving bijections

$$(3.11) \quad \max B_{\kappa} \otimes B_{\omega'} \otimes B_{\omega} \longleftrightarrow \max B_{\kappa} \otimes B_{\omega} \otimes B_{\omega'},$$

for $\kappa, \omega', \omega \in \Lambda^+$ and ω', ω minuscule or quasi-minuscule. Such a bijection is given by

$$(b_{\kappa}, h, v) \mapsto (b_{\kappa}, v', h').$$

Let

$$\lambda := \kappa + \text{wt}(h), \quad \nu := \lambda + \text{wt}(v), \quad \mu := \kappa + \text{wt}(v').$$

When ω' and ω are both dominant minuscule weights, the local moves are easily described by the condition

$$(3.12) \quad \mu = \text{dom}_W(\kappa + \nu - \lambda),$$

where $\text{dom}_W(\cdot)$ denotes, as above, the dominant representative in the corresponding W -orbit. In other words, after computing μ based on h and v , we set

$$v' := \mu - \kappa \quad \text{and} \quad h' := \nu - \mu.$$

It is not hard to check (see [22]) that these moves reduce to Fomin's local moves for his growth diagrams [20, Appendix 1] in the case of the root system A_{n-1} , as long as one uses embeddings of crystals into tensor powers of B_{ε_1} – the crystal of the vector representation of $\mathfrak{sl}_n$. For the definition of the local moves in the case when at least one of the weights ω and ω' is quasi-minuscule, we refer to [22].

In [22], it is proved that these moves are weight-preserving bijections between the sets in (3.11). Hence, conditions (3.5), (3.6), and (3.7) are satisfied. It is also proved that the above moves are reversible, that is, the inverse bijection in (3.11) is given by the same moves. Finally, the following result is proved, based on the initial conditions (3.8) (cf. also (3.2)):

$$(3.13) \quad (h_{i,0}, \dots, h_{i,l-1}) \succeq (h_{i+1,0}, \dots, h_{i+1,l-1}) \quad \text{in } B_{\pi'}, \text{ for } i = 0, \dots, k-1,$$

$$(3.14) \quad (v_{0,j+1}, \dots, v_{k-1,j+1}) \preceq (v_{0,j}, \dots, v_{k-1,j}) \quad \text{in } B_{\pi}, \text{ for } j = 0, \dots, l-1;$$

here we use the partial order on crystals defined in Section 2.2. The above results imply (cf. also (2.5)):

$$\begin{aligned} & ((v_{0,0}, \dots, v_{k-1,0}), (h_{k,0}, \dots, h_{k,l-1})) \\ & = ((\pi_1, \dots, \pi_k), (h_{k,0}, \dots, h_{k,l-1})) \in \max B_{\pi} \otimes B_{\pi'}. \end{aligned}$$

So (3.3) is true based on (3.10), while (3.4) follows from (3.5). The reversibility of the local moves implies that the inverse map, from $\max B_{\pi} \otimes B_{\pi'}$ to $\max B_{\pi'} \otimes B_{\pi}$, is given by the same local moves.

4. THE EXPLICIT REALIZATION OF THE COMMUTOR

In this section, we use the setup in Section 3 without further comment. Assume that we have $(b_{\pi'}, p) \mapsto (b_{\pi}, p')$ by van Leeuwen's *jeu de taquin* (where $p \in B_{\pi}$ and $p' \in B_{\pi'}$). As discussed above, we also have $(b_{\pi}, p') \mapsto (b_{\pi'}, p)$.

Proposition 4.1. *We have $\sigma_{B_{\pi'}, B_{\pi}}(b_{\pi'}, p) = (b_{\pi}, p')$ and $\sigma_{B_{\pi}, B_{\pi'}}(b_{\pi}, p') = (b_{\pi'}, p)$ in the following two cases, which correspond to the special case $l = 1$: (1) π' is a minuscule weight; (2) π' is a quasi-minuscule weight and $p' \in W\pi'$.*

Proof. This is immediate by Remark 2.2. Indeed, in both cases there is a unique choice for the commutor. $\square$

Proposition 4.1 allows us to express the local moves of van Leeuwen (3.11) in terms of the commutor of Henriques and Kamnitzer. We also need to recall the naturality of the latter, namely its property (C1).

Corollary 4.2. *If ω' and ω are minuscule weights, the local move of van Leeuwen from $\max B_{\kappa} \otimes B_{\omega'} \otimes B_{\omega}$ to $\max B_{\kappa} \otimes B_{\omega} \otimes B_{\omega'}$ is given by the following composite:*

$$\sigma_{B_{\omega'}, B_{\kappa} \otimes B_{\omega}} \circ (\sigma_{B_{\kappa}, B_{\omega'}} \otimes 1).$$

The following lemma is based on the notation related to coboundary categories that was introduced in Section 2.4.

Lemma 4.3. *The following two composites in a coboundary category coincide:*

$$(4.1) \quad \begin{aligned} A_1 A_2 A_3 A_4 &\xrightarrow{\sigma_{1,1,2}} A_2 A_1 A_3 A_4 \xrightarrow{\sigma_{1,1,3}} A_1 A_3 A_2 A_4 \\ &\xrightarrow{\sigma_{1,2,4}} A_2 A_4 A_1 A_3 \xrightarrow{\sigma_{1,2,3}} A_1 A_2 A_4 A_3, \end{aligned}$$

$$(4.2) \quad A_1 A_2 A_3 A_4 \xrightarrow{\sigma_{1,2,3}} A_3 A_1 A_2 A_4 \xrightarrow{\sigma_{1,1,4}} A_1 A_2 A_4 A_3.$$

Proof. By (2.12), we have

$$\sigma_{1,1,3} \circ \sigma_{1,1,2} = s_{1,3} \circ s_{2,3} \circ s_{1,2}, \quad \sigma_{1,1,2} \circ \sigma_{1,2,3} = s_{1,2} \circ s_{1,3} \circ s_{1,2}.$$

But, by relation (R3) in the cactus group (see Section 2.4), we have $s_{1,3} \circ s_{2,3} = s_{1,2} \circ s_{1,3}$, which implies

$$(4.3) \quad \sigma_{1,1,3} \circ \sigma_{1,1,2} = \sigma_{1,1,2} \circ \sigma_{1,2,3}.$$

Upon viewing the tensor product $A_3 A_4$ as a single object of the category, the above relation also implies the following one (both sides being defined on $A_1 A_2 A_3 A_4$):

$$(4.4) \quad \sigma_{1,1,4} \circ \sigma_{1,1,2} = \sigma_{1,2,3} \circ \sigma_{1,2,4}.$$

Finally, by (4.3) and (4.4), we have

$$\sigma_{1,2,3} \circ \sigma_{1,2,4} \circ \sigma_{1,1,3} \circ \sigma_{1,1,2} = (\sigma_{1,1,4} \circ \sigma_{1,1,2}) \circ (\sigma_{1,1,2} \circ \sigma_{1,2,3}) = \sigma_{1,1,4} \circ \sigma_{1,2,3}.$$

$\square$

We now arrive at our main result, which we state and prove in the setup of Section 3. At this point, let us recall Remark 3.1 related to the fact that van Leeuwen's *jeu de taquin* can be viewed as a commutor. Let us also recall Remark 3.2 related to the embedding of a crystal into a tensor product of minuscule or quasi-minuscule crystals, which will be used without further comment in the proof below.

Theorem 4.4. *Assume that all weights ω'_j , for $j = 1, \dots, l$, and ω_i , for $i = 1, \dots, k$, are minuscule. We have*

$$\sigma_{B_{\pi'}, B_{\pi}}(b_{\pi'}, p) = (b_{\pi}, p') \quad \text{and} \quad \sigma_{B_{\pi}, B_{\pi'}}(b_{\pi}, p') = (b_{\pi'}, p).$$

Proof. We proceed by induction on $k + l$, also assuming that the result holds for $l = 1$ or $k = 1$, by Proposition 4.1.

Let

$$\bar{\pi}' := \pi'_1 + \dots + \pi'_{l-1}, \quad \bar{\pi} := \pi_1 + \dots + \pi_{k-1}.$$

Let us realize the maxima in the crystals $B_{\bar{\pi}'}$ and $B_{\bar{\pi}}$ as

$$b_{\bar{\pi}'} = (\pi'_1, \dots, \pi'_{l-1}), \quad b_{\bar{\pi}} = (\pi_1, \dots, \pi_{k-1}).$$

Furthermore, let us define

$$\bar{p} := (p_1, \dots, p_{k-1}), \quad \bar{p}' := (p'_1, \dots, p'_{l-1}).$$

Let us also consider the elements $q \in B_{\pi}$ and $q' \in B_{\pi'}$ (cf. (3.13) and (3.14)) defined as follows:

$$\begin{aligned} q &= (q_1, \dots, q_k) \quad \text{for } q_{i+1} := v_{i, l-1}, \quad \text{where } i = 0, \dots, k-1, \text{ and} \\ q' &:= (q'_1, \dots, q'_l) \quad \text{for } q'_{j+1} := h_{k-1, j}, \quad \text{where } j = 0, \dots, l-1. \end{aligned}$$

In a way similar to the definition of $\bar{p}$ and $\bar{p}'$, we define $\bar{q}$ and $\bar{q}'$. Let us now express the van Leeuwen map $(b_{\pi'}, p) \mapsto (b_{\pi}, p')$ as the following composite:

$$\begin{aligned} (4.5) \quad (b_{\pi'}, p) &= (b_{\bar{\pi}'}, \pi'_l, \bar{p}, p_k) \mapsto (b_{\bar{\pi}}, \bar{q}', q'_l, p_k) \mapsto (b_{\bar{\pi}'}, \bar{q}, q'_l, p_k) \\ &\mapsto (b_{\bar{\pi}'}, \bar{q}, q_k, p'_l) \mapsto (b_{\bar{\pi}}, \pi_k, \bar{p}', p'_l) = (b_{\pi}, p'). \end{aligned}$$

Now let us denote

$$B_1 := B_{\omega'_1} \otimes \dots \otimes B_{\omega'_{l-1}}, \quad B_2 := B_{\omega'_l}, \quad B_3 := B_{\omega_1} \otimes \dots \otimes B_{\omega_{k-1}}, \quad B_4 := B_{\omega_k}.$$

By induction, we know that the composition of the first two maps in (4.5) is given by the commutor of Henriques and Kamnitzer, as follows (see the notation in Section 2.4):

$$(4.6) \quad B_1 B_2 B_3 B_4 \xrightarrow{\sigma_{1,2,3}} B_3 B_1 B_2 B_4 \xrightarrow{\sigma_{1,1,2}} B_1 B_3 B_2 B_4.$$

Similarly, the last map in (4.5) is given by

$$(4.7) \quad B_1 B_3 B_4 B_2 \xrightarrow{\sigma_{1,1,3}} B_3 B_4 B_1 B_2.$$

Finally, the third map in (4.5) is just van Leeuwen's local move (3.11); by Corollary 4.2, it can be expressed as

$$(4.8) \quad B_1 B_3 B_2 B_4 \xrightarrow{\sigma_{1,2,3}} B_2 B_1 B_3 B_4 \xrightarrow{\sigma_{1,1,4}} B_1 B_3 B_4 B_2.$$

Combining (4.6), (4.7), and (4.8), we can express the composite (4.5) as the following composition in terms of the commutor of Henriques and Kamnitzer:

$$\begin{aligned} &\sigma_{1,1,3} \circ (\sigma_{1,1,4} \circ \sigma_{1,2,3}) \circ \sigma_{1,1,2} \circ \sigma_{1,2,3} \\ &= \sigma_{1,1,3} \circ (\sigma_{1,2,3} \circ \sigma_{1,2,4} \circ \sigma_{1,1,3} \circ \sigma_{1,1,2}) \circ \sigma_{1,1,2} \circ \sigma_{1,2,3} = \sigma_{1,2,4}. \end{aligned}$$

The first equality follows by rewriting the bracket based on Lemma 4.3, while the second one uses the fact that $\sigma_{1,1,3} = \sigma_{1,2,3}^{-1}$ (cf. property (C2) of the commutor, stated in Section 2.4). $\square$

Remarks 4.5. (1) Any crystal can be embedded into a tensor product of minuscule crystals in the case of an irreducible root system different from the ones of type E_8 , F_4 , and G_2 (cf. Theorem 2.1). Hence, the combinatorial realization (in Theorem 4.4) of the commutor due to Henriques and Kamnitzer based on the local moves of van Leeuwen holds in all the Lie types previously mentioned.

(2) Our construction is more efficient than the original definition of the commutor (2.9). Indeed, the latter uses Lusztig's involution three times. This involution on irreducible crystals B_λ can be realized combinatorially (via *Schützenberger's evacuation* on Young tableaux in type A [1, 5], and via a similar procedure based on the alcove path model in arbitrary type [13]). However, for a tensor product $B_{\pi'} \otimes B_\pi$ of two irreducible crystals, there is no nice realization of this involution other than the definition (2.6); see Example 4.7 for the concrete way to realize it on a minimal element of $B_\pi \otimes B_{\pi'}$ (this is always the case if we compute the action of $\sigma_{B_{\pi'} \otimes B_\pi}$ on $\max B_{\pi'} \otimes B_\pi$). Also note that the realization of the commutor mentioned in [9] involves the Mirković-Vilonen polytopes, which are more complex objects.

(3) We can combine our main result with both realizations of the commutor of Henriques and Kamnitzer (in [6] and [9]) in order to obtain explicit constructions related to: (i) Lusztig's involution on highest weight elements in a tensor product of two irreducible crystals; (ii) Kashiwara's involution.

Remark 4.5 (1) leads us to the following conjecture, which was checked experimentally.

Conjecture 4.6. *Theorem 4.4 holds without the restriction on the embeddings (3.1) of the crystals $B_{\pi'}$ and B_π , i.e., it holds for embeddings into tensor products of minuscule and quasi-minuscule crystals.*

Example 4.7. Consider the root system A_2 , $\pi' = 2\varepsilon_1 + \varepsilon_2$, and $\pi = 2\varepsilon_1$. We use the standard embeddings $B_{\pi'} \hookrightarrow B_{\varepsilon_1}^{\otimes 3}$ and $B_\pi \hookrightarrow B_{\varepsilon_1}^{\otimes 2}$; to be more precise, $b_{\pi'} \mapsto (\varepsilon_1, \varepsilon_1, \varepsilon_2)$. Let

$$(b_{\pi'}, p) = ((\varepsilon_1, \varepsilon_1, \varepsilon_2), (\varepsilon_3, \varepsilon_2)) \in \max B_{\pi'} \otimes B_\pi.$$

We claim that

$$\sigma_{B_{\pi'}, B_\pi}(b_{\pi'}, p) = ((\varepsilon_1, \varepsilon_1), (\varepsilon_2, \varepsilon_2, \varepsilon_3)) \in \max B_\pi \otimes B_{\pi'}.$$

Indeed, by Schützenberger's evacuation, we have $\eta_{B_{\pi'}}(b_{\pi'}) = (\varepsilon_3, \varepsilon_2, \varepsilon_3)$, and $\eta_{B_\pi}(p) = (\varepsilon_2, \varepsilon_1)$. Since the element $((\varepsilon_2, \varepsilon_1), (\varepsilon_3, \varepsilon_2, \varepsilon_3))$ is a minimal element of $B_\pi \otimes B_{\pi'}$, by applying raising operators to it as long as possible, we obtain $((\varepsilon_1, \varepsilon_1), (\varepsilon_2, \varepsilon_2, \varepsilon_3))$. Alternatively, we can use the algorithm based on van Leeuwen's moves, which is shown in Figure 1 below. For simplicity, ε_k is represented here by k ; the labels on the horizontal segments are h_i , while those on the vertical ones are v_j , as indicated. Note that the algorithm inputs the labels on the bottom and right edges of the rectangle; it outputs those on the left and top edges, after labeling the segments in the interior.

FIGURE 1

REFERENCES

1. A. Berenstein and A. Zelevinsky. Canonical bases for the quantum group of type A_r and piecewise-linear combinatorics. *Duke Math. J.*, 82:473–502, 1996. MR1387682 (97g:17007)
2. A. Berenstein and A. Zelevinsky. Tensor product multiplicities, canonical bases and totally positive varieties. *Invent. Math.*, 143:77–128, 2001. MR1802793 (2002c:17005)
3. N. Bourbaki. *Groupes et Algèbres de Lie*. Chp. IV–VI. Masson, Paris, 1981.
4. V. G. Drinfel'd. Quasi-Hopf algebras. *Leningrad Math. J.*, 1:1419–1457, 1990. MR1047964 (91b:17016)
5. W. Fulton. *Young Tableaux*, volume 35 of *London Math. Soc. Student Texts*. Cambridge Univ. Press, Cambridge and New York, 1997. MR1464693 (99f:05119)
6. A. Henriques and J. Kamnitzer. Crystals and coboundary categories. *Duke Math. J.*, 132:191–216, 2006. MR2219257
7. A. Joseph. *Quantum Groups and Their Primitive Ideals*, volume 29 of *Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)]*. Springer-Verlag, Berlin, 1995. MR1315966 (96d:17015)
8. J. Kamnitzer. The crystal structure on the set of Mirković-Vilonen polytopes. *Adv. Math.*, 215:66–93, 2007.
9. J. Kamnitzer and P. Tingley. A definition of the crystal commutor using Kashiwara's involution. [arXiv:math.QA/0610952](#).
10. M. Kashiwara. Crystalizing the q -analogue of universal enveloping algebras. *Commun. Math. Phys.*, 133:249–260, 1990. MR1090425 (92b:17018)
11. M. Kashiwara. On crystal bases. In *Representations of groups (Banff, AB, 1994)*, volume 16 of *CMS Conf. Proc.*, pages 155–197. Amer. Math. Soc., Providence, RI, 1995. MR1357199 (97a:17016)
12. M. Kashiwara and T. Nakashima. Crystal graphs for representations of the q -analogue of classical Lie algebras. *J. Algebra*, 165:295–345, 1994. MR1273277 (95c:17025)
13. C. Lenart. On the combinatorics of crystal graphs, I. Lusztig's involution. *Adv. Math.*, 211:204–243, 2007. MR2313533
14. C. Lenart and A. Postnikov. Affine Weyl groups in K -theory and representation theory. *Int. Math. Res. Not.*, Article ID rnm038, 65pp.
15. C. Lenart and A. Postnikov. A combinatorial model for crystals of Kac-Moody algebras. [arXiv:math.RT/0502147](#). To appear in *Trans. Amer. Math. Soc.*
16. P. Littelmann. Paths and root operators in representation theory. *Ann. of Math. (2)*, 142:499–525, 1995. MR1356780 (96m:17011)
17. P. Littelmann. Characters of representations and paths in $\mathfrak{h}_{\mathbb{R}}^*$. In *Representation theory and automorphic forms (Edinburgh, 1996)*, volume 61 of *Proc. Sympos. Pure Math.*, pages 29–49. Amer. Math. Soc., Providence, RI, 1997. MR1476490 (98j:17024)
18. G. Lusztig. Canonical bases arising from quantized enveloping algebras. II. *Progr. Theoret. Phys. Suppl.*, 102:175–201, 1991. MR1182165 (93g:17019)

19. G. Lusztig. *Introduction to Quantum Groups*, volume 110 of *Progress in Mathematics*. Birkhäuser Boston Inc., Boston, MA, 1993. MR1227098 (94m:17016)
20. R. P. Stanley. *Enumerative Combinatorics. Vol. 2*, volume 62 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 1999. MR1676282 (2000k:05026)
21. J. R. Stembridge. Combinatorial models for Weyl characters. *Adv. Math.*, 168:96–131, 2002. MR1907320 (2003j:17007)
22. M. A. A. van Leeuwen. An analogue of jeu de taquin for Littelmann’s crystal paths. *Sém. Lothar. Combin.*, 41:Art. B41b, 23 pp. (electronic), 1998.

DEPARTMENT OF MATHEMATICS AND STATISTICS, STATE UNIVERSITY OF NEW YORK AT ALBANY,
ALBANY, NEW YORK 12222

E-mail address: `lenart@albany.edu`