MONOTONICITY OF THE PRINCIPAL EIGENVALUE OF THE p -LAPLACIAN IN AN ANNULUS

B. EMAMIZADEH AND M. ZIVARI-REZAPOUR

(Communicated by Walter Craig)

ABSTRACT. In this note we prove a monotonicity result related to the principal eigenvalue of the p -Laplacian in an annulus in $\mathbb{R}^N$.

1. INTRODUCTION

This note is motivated by [3], where the authors consider the principal eigenvalue for $-\Delta$, with respect to the Dirichlet boundary conditions, in $D(h) := B \setminus B_h$. Here B stands for the unit ball in $\mathbb{R}^N$ centered at the origin, and B_h is the ball centered at $(h, 0) \in \mathbb{R} \times \mathbb{R}^{N-1}$ with radius $a < 1$. They show the principal eigenvalue as a function of h decreasing over the interval $[0, 1 - a]$.

In this work we show a similar result, but for $-\Delta_p$. For $p \in (1, \infty)$, Δ_p denotes the usual p -Laplacian operator; that is $\Delta_p u = \nabla \cdot (|\nabla u|^{p-2} \nabla u)$. Let us describe the problem. We fix $-1 < a_0 < 0$, and consider $f \in L^\infty(B)$. We assume $f^+(x) := \max(f(x), 0)$ is supported in $S := \{x \in B : x_1 < a_0\}$ and does not vanish identically. Also, we assume $f^- := f^+ - f$ is non-increasing in the x_1 -variable. Let us consider the following eigenvalue problem:

$$(BVP) \quad \begin{cases} -\Delta_p u = \lambda f(x) |u|^{p-2} u & \text{in } D(h), \\ u = 0 & \text{on } \partial D(h). \end{cases}$$

By $\lambda(h)$ we denote the principal eigenvalue of (BVP), and by $u(h)$ the corresponding positive (principal) eigenfunction normalized by

$$(1) \quad \int_{D(h)} f(x) u^p(h) \, dx = 1.$$

It is well known that $\lambda(h)$ is simple, $u(h) \in C^1(\overline{D(h)})$, and $\partial u(h)/\partial \nu < 0$ on $\partial D(h)$, where ν stands for the unit outward normal on $\partial D(h)$; see for example [2]. The main result of the paper is the following

Theorem 1. *The principal eigenvalue $\lambda(h)$ is decreasing on $[-a_0, 1 - a]$.*

The idea of the proof of Theorem 1 is straightforward; we show the derivative of $\lambda(h)$ is non-positive on $[-a_0, 1 - a]$. This is the same idea employed in [3], however

Received by the editors December 13, 2006 and, in revised form, February 19, 2007 and February 28, 2007.

2000 *Mathematics Subject Classification.* Primary 35P30.

Key words and phrases. p -Laplacian, principal eigenvalue, domain derivative.

©2007 American Mathematical Society
 Reverts to public domain 28 years from publication

in our case the technicalities are very different. In particular we use the concept of domain derivative to a great extent, and the reference we use for this is [4].

2. DOMAIN DERIVATIVE

Throughout this section Ω denotes a bounded smooth domain in $\mathbb{R}^N$. For a vector field $V \in C^2(\mathbb{R}^N, \mathbb{R}^N)$, and non-negative t , we denote by Ω_t the image of Ω under the map $I + tV$. It is well known that when t is small the map $I + tV$ is a diffeomorphism, hence Ω_t would be as “good” as Ω , roughly speaking.

Suppose $\mathcal{L}$ is a differential operator that maps $W^{m,p}(\Omega')$ into $\mathcal{D}'(\Omega')$, the space of distributions on Ω' , where Ω' is any open set inside Ω . Suppose w is the solution of the following boundary value problem:

$$(2) \quad \begin{cases} \mathcal{L}u = 0 & \text{in } \Omega', \\ u = 0 & \text{on } \partial\Omega'. \end{cases}$$

Also suppose w^t is the solution of (2) with Ω' replaced by Ω'_t . Then the function w' defined on Ω' by

$$w'(x) = \lim_{t \rightarrow 0^+} \frac{w^t(x^t) - w(x)}{t}$$

where $x^t := x + tV(x)$ is called the domain derivative of w at Ω' in the direction of V . In practice w' usually satisfies

$$\begin{cases} \frac{\partial \mathcal{L}}{\partial u} w' = 0 & \text{in } \Omega', \\ w' = -\frac{\partial w}{\partial \nu} V \cdot \nu & \text{on } \partial\Omega', \end{cases}$$

where ν stands for the unit outward normal to $\partial\Omega'$. The reader is referred to [4] for a thorough discussion on domain derivatives. As an example we consider the (BVP) introduced in the previous section. For simplicity we write u and λ in place of $u(h)$ and $\lambda(h)$, respectively. Then we have the following result

Lemma 1. *With the notation introduced above, u satisfies*

$$(3) \quad \begin{cases} -\nabla \cdot ((p-2)|\nabla u|^{p-1} \frac{\nabla u \cdot \nabla u'}{|\nabla u|^3} \nabla u + |\nabla u|^{p-2} \nabla u') \\ \quad = \lambda' f(x) u^{p-1} + (p-1) \lambda f(x) u^{p-2} u' & \text{in } D(h), \\ u' = -\frac{\partial u}{\partial \nu} V \cdot \nu & \text{on } \partial D(h). \end{cases}$$

Proof. The boundary equation in (3) follows directly from [4, Theorem 3.2, p. 664], and the right hand side of the differential equation follows from the product rule for domain derivatives, noting that $f(x)$ does not depend on the domain. To derive the left hand side of the differential equation we apply [4, Theorem 3.1, p. 663]. Indeed since the domain differential operator commutes with the divergence operator we only need to show

$$(|\nabla u|^{p-2} \nabla u)' = (p-2)|\nabla u|^{p-1} \frac{\nabla u \cdot \nabla u'}{|\nabla u|^3} \nabla u + |\nabla u|^{p-2} \nabla u',$$

where the “prime” denotes the domain derivative. The identity

$$|\nabla u|^{p-2} \nabla u = (\nabla u \cdot \nabla u)^{\frac{p-2}{2}} \nabla u$$

is clear, so when we differentiate both sides, applying the product rule to the right hand side, we obtain

$$\begin{aligned} (|\nabla u|^{p-2} \nabla u)' &= \frac{p-2}{2} (\nabla u \cdot \nabla u)^{\frac{p-2}{2}-1} (\nabla u' \cdot \nabla u + \nabla u \cdot \nabla u') \nabla u \\ &+ (\nabla u \cdot \nabla u)^{\frac{p-2}{2}} \nabla u' \\ &= (p-2) |\nabla u|^{p-1} \frac{\nabla u \cdot \nabla u'}{|\nabla u|^3} \nabla u + |\nabla u|^{p-2} \nabla u', \end{aligned}$$

as desired. $\square$

3. PROOF OF THEOREM 1

In this section we prove Theorem 1, for which we need the following “weak comparison principle”.

Theorem 2. *Let $\Omega \subset \mathbb{R}^N$ be a bounded domain with piecewise smooth boundary $\partial\Omega$. Suppose ω is an open smooth subset of $\partial\Omega$. Suppose $g : \Omega \times [0, \infty) \rightarrow [0, \infty)$ is a Carathéodory function such that $g(x, \cdot)$ is non-decreasing for every $x \in \Omega$. Let $v, u \in C^1(\overline{\Omega})$ be non-negative weak solutions of the following differential inequalities:*

$$(4) \quad \begin{cases} -\Delta_p v + g(x, v) \leq 0, \\ -\Delta_p u + g(x, u) \geq 0, \end{cases}$$

in Ω , respectively. Suppose $u = v$ on ω and $v \leq u$ on $\partial\Omega \setminus \omega$. Also, suppose $\partial u / \partial \nu < 0$ on ω . Then there is a neighborhood of ω , denoted $\mathcal{N}(\omega) \subset \Omega$, such that $u \geq v$ in $\mathcal{N}(\omega)$.

Proof. It is clear that if $\{x \in \Omega : v(x) > u(x)\}$ is empty, then the assertion of the theorem follows trivially; so we assume otherwise. It suffices to show that the boundary of the open set $\{x \in \Omega : v(x) > u(x)\}$ does not intersect ω . Note that since $u \geq v$ on $\partial\Omega$, $(v - u)^+ \in W_0^{1,p}(\Omega)$; thus it can be used as a test function in both inequalities in (4). After multiplying each of the inequalities in (4) by $(v - u)^+$ and integrating over Ω we obtain

$$\int_{\Omega} |\nabla v|^{p-2} \nabla v \cdot \nabla (v - u)^+ dx + \int_{\Omega} g(x, v) (v - u)^+ dx \leq 0$$

and

$$\int_{\Omega} |\nabla u|^{p-2} \nabla u \cdot \nabla (v - u)^+ dx + \int_{\Omega} g(x, u) (v - u)^+ dx \geq 0.$$

Thus

$$\begin{aligned} &\int_{\Omega} (|\nabla v|^{p-2} \nabla v - |\nabla u|^{p-2} \nabla u) \cdot \nabla (v - u)^+ dx \\ &\leq \int_{\Omega} (g(x, u) - g(x, v)) (v - u)^+ dx \\ &= \int_{\{v > u\}} (g(x, u) - g(x, v)) (v - u)^+ dx. \end{aligned}$$

The last integral is non-positive since $g(x, \cdot)$ is non-decreasing. This in conjunction with

$$\nabla (v - u)^+ = \begin{cases} \nabla (v - u) & \text{if } v > u, \\ 0 & \text{if } v \leq u, \end{cases}$$

yields

$$(5) \quad \int_{\{v>u\}} (|\nabla v|^{p-2} \nabla v - |\nabla u|^{p-2} \nabla u) \cdot \nabla(v-u) \, dx \leq 0.$$

At this stage we recall the following result, whose proof can be found in [1, Lemma 2.1]: There exists $c > 0$ such that for any $\eta, \eta' \in \mathbb{R}^N$ with $|\eta| + |\eta'| > 0$, the following holds:

$$(6) \quad (|\eta|^{p-2} \eta - |\eta'|^{p-2} \eta') \cdot (\eta - \eta') \geq c (|\eta| + |\eta'|)^{p-2} |\eta - \eta'|^2.$$

Applying (6) to (5) we find

$$(7) \quad c \int_{\{v>u\}} (|\nabla u| + |\nabla v|)^{p-2} |\nabla u - \nabla v|^2 \, dx \leq 0.$$

In order to derive a contradiction let us assume $\partial\{x \in \Omega : v(x) > u(x)\}$ intersects ω , and consider $x_0 \in \partial\{x \in \Omega : v(x) > u(x)\} \cap \omega$. From the hypotheses we infer $\frac{\partial u}{\partial \nu}(x_0) < 0$. Since $u \in C^1(\bar{\Omega})$, it follows that there exists a positive constant γ and an open set, relative to Ω , denoted $V(x_0)$, containing x_0 such that $|\nabla u(x)| \geq \gamma$ for every $x \in V(x_0)$. Note that $x_0 \in \partial W(x_0)$, where $W(x_0) := \{x \in \Omega : v(x) > u(x)\} \cap V(x_0)$. From (7) we deduce that

$$c \int_{W(x_0)} (|\nabla u| + |\nabla v|)^{p-2} |\nabla u - \nabla v|^2 \, dx \leq 0.$$

Hence

$$c \inf_{x \in W(x_0)} \{|\nabla u(x)| + |\nabla v(x)|\}^{p-2} \int_{W(x_0)} |\nabla(u-v)|^2 \, dx \leq 0.$$

Since $|\nabla u| > \gamma$ on $W(x_0)$, it follows from the above inequality that

$$c\gamma^{p-2} \int_{W(x_0)} |\nabla(u-v)|^2 \, dx \leq 0,$$

so $\int_{W(x_0)} |\nabla(u-v)|^2 \, dx = 0$. This, in turn, implies that $u-v = \text{const.}$, in $W(x_0)$. Since $x_0 \in \partial W(x_0)$, and $u(x_0) = v(x_0)$, we infer $u \equiv v$, in $W(x_0)$. However, this is not possible since $W(x_0) \subseteq \{x \in \Omega : v(x) > u(x)\}$. So, as desired, $\partial\{x \in \Omega : v(x) > u(x)\}$ does not intersect ω . $\square$

Remark. It is worth noting that if $p \geq 2$, the condition $\frac{\partial u}{\partial \nu} < 0$, on ω , in Theorem 2 can be dropped. Indeed, in order to show that $\partial\{x \in \Omega : v(x) > u(x)\}$ does not intersect ω , we proceed along the same lines as in the proof of the theorem until we reach (5). Then we use the following inequality whose proof like (6) can be found in [1, Lemma 2.1]: If $p \geq 2$, then there exists $\hat{c} > 0$ such that for every $\eta, \eta' \in \mathbb{R}$,

$$(8) \quad (|\eta|^{p-2} \eta - |\eta'|^{p-2} \eta') \cdot (\eta - \eta') \geq \hat{c} |\eta - \eta'|^p.$$

Applying (8) to (5) yields

$$\hat{c} \int_{\{v>u\}} |\nabla u - \nabla v|^p \, dx \leq 0,$$

so $\int_{\{v>u\}} |\nabla u - \nabla v|^p \, dx = 0$. Whence $u-v = \text{const.}$, on $\{x \in \Omega : v(x) > u(x)\}$. So if $x_0 \in \partial\{x \in \Omega : v(x) > u(x)\} \cap \omega$, then $u \equiv v$ in $\{x \in \Omega : v(x) > u(x)\}$, which is false, and we have a contradiction.

We are now ready to prove Theorem 1. As in the previous section we replace $u(h)$ by u .

Proof of Theorem 2. We multiply the differential equation in (3) by u and integrate the result over $D(h)$. Thus we obtain

$$(9) \quad (p-1) \int_{D(h)} |\nabla u|^{p-2} \nabla u \cdot \nabla u' \, dx = \lambda'(h) \int_{D(h)} f u^p \, dx + (p-1) \lambda(h) \int_{D(h)} f u^{p-1} u' \, dx.$$

The right-hand side of (9) simplifies a great deal. Indeed from (1) we infer that the (domain) derivative of $\int_{D(h)} f u^p \, dx$ must be zero. Hence using [4, Theorem 3.3] we obtain

$$\left(\int_{D(h)} f u^p \, dx \right)' = p \int_{D(h)} f u^{p-1} u' \, dx + \int_{\partial D(h)} f u^p V \cdot \nu \, d\sigma = 0,$$

where $d\sigma$ stands for the surface measure, and the “prime” denotes the domain derivative with respect to some vector field V . Since u is zero on $\partial D(h)$ we deduce that $\int_{D(h)} f u^{p-1} u' \, dx = 0$, hence from (9) we find the following formula:

$$(10) \quad (p-1)^{-1} \lambda'(h) = \int_{D(h)} |\nabla u|^{p-2} \nabla u \cdot \nabla u' \, dx,$$

again thanks to (1). Let us recall the following characterization of the principal eigenvalue:

$$\lambda(h) = \int_{D(h)} |\nabla u|^p \, dx.$$

Hence, another application of [4, Theorem 3.3] yields

$$(11) \quad \lambda'(h) = p \int_{D(h)} |\nabla u|^{p-2} \nabla u \cdot \nabla u' \, dx + \int_{\partial D(h)} |\nabla u|^p V \cdot \nu \, d\sigma.$$

From (10) and (11) we find

$$(12) \quad \lambda'(h) = (1-p) \int_{\partial D(h)} |\nabla u|^p V \cdot \nu \, d\sigma.$$

Let us choose a particular vector field as follows. Fix $\tilde{\phi} \in C_0^\infty(B)$ such that $0 \leq \tilde{\phi} \leq 1$ and $\tilde{\phi} \equiv 1$ on a bounded neighborhood of B_h inside B . The zero extension of $\tilde{\phi}$ to the whole of $\mathbb{R}^N$ is denoted by ϕ . Finally we let $V := \phi \vec{i}$, where $\vec{i}$ is the standard unit vector in $\mathbb{R}^N$. Obviously $V \in C^2(\mathbb{R}^N; \mathbb{R}^N)$, so it can be inserted in (12). Thus we obtain

$$(13) \quad \lambda'(h) = (1-p) \int_{\partial B_h} |\nabla u|^p \nu_1 \, d\sigma,$$

where ν_1 is the first component of ν . Note that since $p > 1$ the assertion of the theorem follows once we establish that the integral in (13) is non-negative. To this end we shift the coordinate axes so that the hyperplane $l = \{x \in \mathbb{R}^N : x_1 = 0\}$ is positioned in such a way that it passes through the center of B_h , hence for $(x_1, y) \in B_h$ we have $(-x_1, y) \in B_h$, as well; that is, B_h is symmetric with respect to l . Let us introduce some more notation. By A_l we denote the part of $D(h)$ for which $x_1 \geq 0$, and the image of A_l , with respect to l , is denoted A_l^* . Also B_h^+ will stand for the part of B_h for which $x_1 \geq 0$, and B_h^- , the part of B_h for which $x_1 \leq 0$, in the new coordinate system. Next we introduce the function

$$v(x) = \begin{cases} u(x) & \text{if } x \in A_l, \\ u(x_l) & \text{if } x \in A_l^*, \end{cases}$$

where x_l stands for the reflection of x about l . Observe that since $h \in [a_0, 1 - a]$, $f(x) = -f^-(x)$, for $x \in A_l^*$. Therefore

$$(14) \quad -\Delta_p u + \lambda f^-(x) u^{p-1} = 0, \quad x \in A_l^*.$$

On the other hand, using the definition of v , we have

$$\begin{aligned} -\Delta_p v(x) + \lambda f^-(x) v^{p-1}(x) &= -\Delta_p u(x_l) + \lambda f^-(x) u^{p-1}(x_l) \\ &\leq -\Delta_p u(x_l) + \lambda f^-(x_l) u^{p-1}(x_l) \\ &= -\Delta_p u(x_l) + \lambda f(x_l) u^{p-1}(x_l) \\ &= 0, \end{aligned}$$

for $x \in A_l^*$. In the inequality above we used the fact that f^- is decreasing in the x_1 -variable. So we have

$$(15) \quad -\Delta_p v + \lambda f^-(x) v^{p-1} \leq 0, \quad x \in A_l^*.$$

Thus we can apply Theorem 2, with $g(x, z) := \lambda f^-(x) z^{p-1}$, $\omega := \partial B_h^-$, to (14) and (15). Hence we infer existence of a neighborhood of ∂B_h^- , denoted $\mathcal{N}(\partial B_h^-)$, such that $u \geq v$ in $\mathcal{N}(\partial B_h^-)$. So, since u and v both vanish on ∂B_h^- , we obtain

$$\frac{\partial}{\partial \nu}(u - v)(z) = \lim_{t \rightarrow 0^-} \frac{(u - v)(z + t\nu) - (u - v)(z)}{t} \leq 0, \quad z \in \partial B_h^-,$$

provided t is small enough to ensure $z + t\nu \in \mathcal{N}(\partial B_h^-)$. From the above inequality we infer $\frac{\partial u}{\partial \nu}(z) \leq \frac{\partial v}{\partial \nu}(z)$, for $z \in \partial B_h^-$. However, since $\frac{\partial v}{\partial \nu}(z) \leq 0$, for $z \in \partial B_h^-$, it follows that $|\frac{\partial u}{\partial \nu}(z)| \geq |\frac{\partial v}{\partial \nu}(z)|$, for $z \in \partial B_h^-$. Note that for any $z \in \partial B_h^-$, we have $\nabla u(z) = -|\nabla u(z)|\nu(z)$. This, in turn, implies $\frac{\partial u}{\partial \nu}(z) = -|\nabla u(z)|$, so $|\frac{\partial u}{\partial \nu}(z)| = |\nabla u(z)|$, for $z \in \partial B_h^-$. Similarly one can show $|\frac{\partial v}{\partial \nu}(z)| = |\nabla v(z)|$, for $z \in \partial B_h^-$. Whence we derive $|\nabla u(z)| \geq |\nabla v(z)|$, for $z \in \partial B_h^-$. Thus

$$\begin{aligned} \int_{\partial B_h^-} |\nabla u(z)|^p \nu_1(z) \, d\sigma(z) &\geq \int_{\partial B_h^-} |\nabla v(z)|^p \nu_1(z) \, d\sigma(z) \\ &= \int_{\partial B_h^-} |\nabla u(z_l)|^p \nu_1(z) \, d\sigma(z) \\ &= \int_{\partial B_h^+} |\nabla u(z)|^p \nu_1(z_l) \, d\sigma(z) \\ &= - \int_{\partial B_h^+} |\nabla u(z)|^p \nu_1(z) \, d\sigma(z). \end{aligned}$$

From this we get $\int_{\partial B_h^-} |\nabla u|^p \nu_1 \, d\sigma \geq - \int_{\partial B_h^+} |\nabla u|^p \nu_1 \, d\sigma$, hence $\int_{\partial B_h} |\nabla u|^p \nu_1 \, d\sigma \geq 0$, as desired. $\square$

Remark. From Theorem 1, we infer that if the support of f^+ is concentrated at $(-1, 0) \in \mathbb{R} \times \mathbb{R}^{N-1}$, then $\lambda(h)$ is maximized when the two balls B and B_h are almost concentric.

REFERENCES

- [1] L. Damascelli, *Comparison theorems for some quasilinear degenerate elliptic operators and applications to symmetry and monotonicity results*, Ann. Inst. Henri Poincaré, vol. 15, n. 4, 1998, p. 493-516. MR1632933 (99e:35081)
- [2] M. Otani and T. Teshima, *On the first eigenvalue of some quasilinear elliptic equations*, Proc. Japan Acad. Ser. A Math. Sci., Vol. 64, no. 1, 1988, p. 8-10. MR953752 (89h:35257)

- [3] A.G. Ramm and P.N. Shivakumar, *Inequalities for the minimal eigenvalue of the Laplacian in an annulus*, Math. Inequal. Appl., Vol. 1, n. 4, 1998, p. 559–563. MR1646670 (99f:35149)
- [4] J. Simon, *Differentiation with respect to the domain in boundary value problems*, Numer. Funct. Anal. and Optimiz., Vol. 2(7&8), 1980, p. 649–687. MR619172 (83m:49032)

DEPARTMENT OF MATHEMATICS, THE PETROLEUM INSTITUTE, P.O. Box 2533, ABU DHABI,
UNITED ARAB EMIRATES

E-mail address: `bemamizadeh@pi.ac.ae`

DEPARTMENT OF MATHEMATICS, IRAN UNIVERSITY OF SCIENCE AND TECHNOLOGY, NARMAK,
TEHRAN, IRAN

E-mail address: `m_zivari@iust.ac.ir`