AN EXTREMAL PROPERTY OF JACOBI POLYNOMIALS IN TWO-SIDED CHERNOFF-TYPE INEQUALITIES FOR HIGHER ORDER DERIVATIVES

VLADIMIR D. STEPANOV

(Communicated by Jonathan M. Borwein)

ABSTRACT. For a weight function generating the classical Jacobi polynomials, the sharp double estimate of the distance from the subspace of all polynomials of an arbitrary fixed order is established.

1. INTRODUCTION

Let $\Delta := (-1, 1)$ and let $\omega(x) > 0$ be a Lebesgue integrable weight function. Denote by $L^2(\omega)$ the real Hilbert space with the inner product $(f, g)_{L^2(\omega)} := \int_{\Delta} fg\omega$ and the norm $\|f\|_{L^2(\omega)} := (f, f)_{L^2(\omega)}^{1/2}$. Let $\mathbf{P}$ be the set of all polynomials and let $\mathbf{P}_k \subset \mathbf{P}$ be the set of all polynomials of degree $\leq k$.

We study the problem on a sharp two-sided estimate of the distance

$$D_{k,\omega}(G) := [\text{dist}_{L^2(\omega)}(G, \mathbf{P}_{k-1})]^2 := \inf_{c_0, \dots, c_{k-1}} \int_{\Delta} \left| G(x) - \sum_{j=0}^{k-1} c_j x^j \right|^2 \omega(x) dx$$

by an analog with the well-known two-sided inequality from the probability theory

$$(1.1) \quad [\mathbf{E}G'(X)]^2 \leq \mathbf{D}[G(X)] \leq \mathbf{E}[G'(X)]^2,$$

which is valid for any absolutely continuous function G with a finite right-hand side (1.1) and the Gaussian normal random variable X . Moreover, both inequalities become the equalities for $G(x) = \text{const} \cdot x$. The right-hand side of (1.1), proved in [1], is often cited as the *Chernoff inequality*; however, in a form even more general than (1.1), it can be derived from the earlier work [2]. An accurate piece of the history of the problem and the extension of (1.1) for the L^p -deviation, $p \geq 1$, can be found in [3]. Unlike [3] and many other works, we extend (1.1) in a different way, increasing the order of derivative, but saving the L^2 -metric. In this case the role of the extremal functions, when both inequalities become the equalities, is to

Received by the editors March 4, 2006.

2000 *Mathematics Subject Classification.* Primary 26D10; Secondary 33C45, 60E15.

Key words and phrases. Jacobi polynomials, Chernoff inequality.

The work of the author was financially supported by the Russian Foundation for Basic Researches (Projects 05-01-00422, 06-01-00341, 06-01-04006 and 07-01-00054) and by the INTAS grant 05-1000008-8157.

play the orthogonal polynomials. A natural extension of (1.1) for the k th derivative found in [4] has a form

$$(1.2) \quad \frac{1}{k!} \left[\int_{\mathbb{R}} G^{(k)} dF \right]^2 \leq D_{k,F}(G) \leq \frac{1}{k!} \int_{\mathbb{R}} |G^{(k)}|^2 dF,$$

where $dF(x) := \frac{1}{\sqrt{2\pi}} e^{-x^2/2} dx$. Moreover, the double equality in (1.2) takes place for the Chebyshev-Hermite polynomials of degree k .

In the present paper we consider an analog of (1.1) for the case of a finite interval Δ and the weight

$$(1.3) \quad \omega(x) = (1-x)^\alpha(1+x)^\beta, \quad \alpha > -1, \beta > -1,$$

when the extremal functions are the classical Jacobi polynomials. Similar to [4], we obtain the result with the help of weighted differential operators for which the Jacobi polynomials are the eigenfunctions (Section 3). Section 2 is devoted to analysis of the case with a general weight ω , and we show that the lower bound is always true with the extremal functions as the orthogonal polynomials with weight ω and the upper bound with the same extremal functions is valid, roughly speaking, if a polynomial of the second order is the eigenfunction of a related differential operator of the second order, which in turn implies that the weight function ω has the form (1.3).

We use the symbols $:=$ and $=$: for a definition of new variables, $\mathbb{N}$ is the set of all natural integers, $\mathbb{N}_0 := \{0\} \cup \mathbb{N}$, $c_i, i = 1, 2, \dots$ are positive constants possibly different from place to place.

2. GENERAL CASE

Let $\omega(x) > 0$, $x \in \Delta$, be a weight function such that $\int_{\Delta} \omega(x) dx = 1$. Then the orthonormal system of polynomials $\{\varphi_k\}$, $k \in \mathbb{N}_0$, such that $\varphi_0(x) = 1$,

$$\varphi_k(x) = (-1)^k a_k x^k + \dots + a_0, \quad k \geq 1,$$

with $a_k > 0$ and

$$\int_{\Delta} \varphi_k^2 \omega = 1$$

is complete in the real Hilbert space $\mathbb{H} := L^2(\omega)$ (see [6], § 1.3.2).

We shall need the Riemann-Liouville transform

$$R_k g(x) := \frac{1}{(k-1)!} \int_{\mathbb{R}} r_k(x, t) g(t) dt,$$

where $\mathbb{R} := (-\infty, \infty)$ and

$$r_k(x, t) := \begin{cases} (x-t)^{k-1}, & 0 \leq t \leq x < \infty, \\ (-1)^k (t-x)^{k-1}, & -\infty < x \leq t < 0, \\ 0, & \text{otherwise.} \end{cases}$$

Let $AC^{(k)}(\Delta)$ denote the class of all functions on Δ with absolutely continuous $(k-1)$ -th derivative. It easy to see that for any $G \in AC^{(k)}(\Delta)$ the following is

true:

$$(2.1) \quad G(x) = R_k G^{(k)}(x) + \sum_{j=0}^{k-1} \frac{x^j}{j!} G^{(j)}(0),$$

$$(2.2) \quad G - \sum_{j=0}^{k-1} (G, p_j)_H p_j = R_k G^{(k)} - \sum_{j=0}^{k-1} (R_k G^{(k)}, p_j)_H p_j.$$

Define

$$\rho_k(x) := \frac{1}{(k-1)!} \int_{-1}^x (x-t)^{k-1} \varphi_k(t) \omega(t) dt, \quad k \in \mathbb{N}.$$

Then

$$(2.3) \quad \rho_k^{(n)}(\pm 1) = 0, \quad n = 0, 1, \dots, k-1.$$

Moreover, $\rho_k(x)$ can have alternating signs, but

$$\lambda_k^{1/2} := \int_{\Delta} \rho_k(x) dx = \frac{1}{a_k k!} > 0, \quad k \in \mathbb{N}.$$

If we put

$$\mu_k(x) := \lambda_k^{-1/2} \rho_k(x), \quad k \in \mathbb{N},$$

then for

$$D_{k,\omega}(G) := [\text{dist}_{\mathbb{H}}(G, P_{k-1})]^2, \quad k \in \mathbb{N}$$

the lower bound

$$(2.4) \quad \lambda_k \left[\int_{\Delta} G^{(k)}(x) \mu_k(x) dx \right]^2 \leq D_{k,\omega}(G)$$

holds (see [4], Lemma 1) provided $G \in AC^{(k)}(\Delta)$ and $G^{(k)} \in L^2(\bar{\rho}_k)$, where

$$\bar{\rho}_k(t) := \int_{\Delta} |r_k(x, t) \varphi_k(x)| \omega(x) dx.$$

Moreover, (2.4) turns into an equality for $G = \text{const} \cdot \varphi_k$.

Now, suppose that $\rho_k(x) \geq 0$ and for all $2k$ times differentiable functions $y(x)$ we define differential operators

$$L_k y := (-1)^k D^k [\rho_k D^k y].$$

Then

$$L_k \varphi_k = (-1)^k D^k [\rho_k D^k \varphi_k] = a_k k! \rho_k^{(k)} = \lambda_k^{-1/2} \varphi_k \omega,$$

so φ_k is an eigenfunction of the weighted differential operator L_k .

Suppose $\varphi_n, n > k$, are the eigenfunctions of L_k too, that is,

$$(2.5) \quad L_k \varphi_n = \sigma_{k,n} \varphi_n \omega, \quad n > k \geq 1,$$

and, moreover,

$$(2.6) \quad \lambda_k^{-1/2} = \sigma_{k,k} \leq \sigma_{k,n}, \quad n > k \geq 1.$$

Then for all polynomials $y \in \mathbf{P}$ such that

$$(y, \varphi_n)_{\mathbb{H}} = 0, \quad n = 0, 1, \dots, k-1,$$

the inequality

$$\int_{\Delta} y L_k y \geq \lambda_k^{-1/2} \int_{\Delta} y^2 \omega$$

holds and becomes the equality for $y = \text{const} \cdot \varphi_k$. By the density $\mathbf{P} \subset L^2(\mu_k)$ it implies the inequality

$$(2.7) \quad D_{k,\omega}(G) \leq \lambda_k \int_{\Delta} |G^{(k)}|^2 \mu_k$$

for all $G \in AC^{(k)}(\Delta)$ such that $G^{(k)} \in L^2(\mu_k)$ (see the end of the proof of Theorem 3.1 for details).

Thus, we obtain the following.

Theorem 2.1. *Let $k \in \mathbb{N}$ and let a weight function $\omega(x) > 0$, $x \in \Delta$, be such that $\int_{\Delta} \omega = 1$. Let $\{\varphi_n\}$, $n \in \mathbb{N}$, be the above constructed orthonormal sequence of polynomials in $\mathbb{H} = L^2(\omega)$. Then for any function $G \in AC^{(k)}(\Delta)$ such that $G^{(k)} \in L^2(\bar{\rho}_k)$, the inequality (2.4) holds. Moreover, if the sequence $\{\varphi_n\}$ satisfies (2.5), (2.6), and $\rho_k(x) \geq 0$, then for any function $G \in AC^{(k)}(\Delta)$ such that $G^{(k)} \in L^2(\mu_k)$, the inequality (2.7) holds, and both inequalities (2.4) and (2.7) simultaneously become the equalities, when $G = \text{const} \cdot \varphi_k$.*

Naturally, the sequence of conditions (2.5), (2.6) looks much too restrictive. To this end we consider the first of the equations, (2.5):

$$(2.8) \quad L_1 \varphi_2 = \sigma_{1,2} \varphi_2 \omega =: \sigma \varphi_2 \omega,$$

where $\sigma > 0$ is some number and $\varphi_2(x) = b_2 x^2 + b_1 x + b_0$, $b_2 > 0$. Taking into account that $\varphi_1 \in \mathbf{P}_1$, $\varphi_2 \in \mathbf{P}_2$, and

$$\rho_1(x) = \int_{-1}^x \varphi_1 \omega,$$

we obtain

$$L_1 \varphi_2 = -(\rho_1 \varphi_2')' = -\varphi_1 \varphi_2' \omega - b_2 \rho_1 = \sigma \varphi_2 \omega.$$

Consequently,

$$(2.9) \quad \rho_1 b_2 = [-\varphi_1 \varphi_2' - \sigma \varphi_2] \omega =: q_2 \omega,$$

where $q_2 \in \mathbf{P}_2$. It follows from (2.3) and (2.9) that

$$(2.10) \quad \lim_{x \rightarrow \pm 1} q_2(x) \omega(x) = 0.$$

Moreover, it follows from (2.9) that ω is differentiable on the interval Δ and satisfies the Pirson equation

$$(2.11) \quad \frac{\omega'}{\omega} = \frac{b_2 \varphi_1 - q_2'}{q_2} =: \frac{q_1}{q_2},$$

where $q_1 \in \mathbf{P}_1$. It is known (see [5], Chapter II, § 1) that, under the conditions $\omega(x) > 0$, $x \in \Delta$, $\int_{\Delta} \omega < \infty$ and (2.10), the equation (2.11) has the only solutions of the form

$$\omega(x) = \text{const}(1-x)^{\alpha}(1+x)^{\beta},$$

where $\alpha > -1$, $\beta > -1$. To close the circle, we show in the next section that polynomials generated by this weight (Jacobi polynomials) satisfy (2.5) and (2.6) as well as $\rho_k(x) \geq 0$ for this case, and the condition $G^{(k)} \in L^2(\bar{\rho}_k)$ of Theorem 2.1 can be weakened up to $G^{(k)} \in L^2(\mu_k)$.

3. CHERNOFF-TYPE INEQUALITY

Again let $H := L^2(\omega)$ be the Hilbert space with the inner product $(f, g)_{L^2(\omega)} := \int_{\Delta} fg\omega$ and the norm $\|f\|_{L^2(\omega)} := (f, f)_{L^2(\omega)}^{1/2}$, where the weight function $\omega(x)$ has the form (1.3). Denote by $P_n(x)$, $n \in \mathbb{N}_0$, the orthogonal sequence of Jacobi polynomials. It is known that $\{P_n(x)\}$ is complete in H . By Rodrigues formula (see [5], Chapter VII, § 1), we have

$$(3.1) \quad P_n(x) = \frac{(-1)^n}{2^n n!} (1-x)^{-\alpha} (1+x)^{-\beta} D^n [(1-x)^{\alpha+n} (1+x)^{\beta+n}],$$

where $D := \frac{d}{dx}$. It is known (see [5], Chapter VII, § 1) that

$$(3.2) \quad \eta_n^2 := \|P_n\|_H^2 = \frac{2^{\alpha+\beta+1} \Gamma(\alpha+n+1) \Gamma(\beta+n+1)}{n! (\alpha+\beta+2n+1) \Gamma(\alpha+\beta+n+1)},$$

where $\Gamma(\cdot)$ denotes the gamma function. Let

$$(3.3) \quad p_n := P_n / \|P_n\|_H$$

be the orthogonal sequence of Jacobi polynomials. Then

$$(3.4) \quad D_{k,\omega}(G) = \|G\|_H^2 - \sum_{j=0}^{k-1} (G, p_j)_H^2.$$

For $n \in \mathbb{N}$ we put

$$(3.5) \quad \beta_n := \int_{\Delta} (1-t)^{\alpha+n} (1+t)^{\beta+n} dt = 2^{\alpha+\beta+2n+1} \frac{\Gamma(\alpha+n+1) \Gamma(\beta+n+1)}{\Gamma(\alpha+\beta+2n+2)},$$

$$(3.6) \quad \nu_n(x) := \frac{(1-x)^{\alpha+n} (1+x)^{\beta+n}}{\beta_n},$$

$$(3.7) \quad \gamma_n := \frac{\beta_n^2}{\eta_n^2 2^{2n} [n!]^2}.$$

Theorem 3.1. *Let $k \in \mathbb{N}$. Then for any function $G \in AC^{(k)}(\Delta)$ such that $G^{(k)} \in L^2(\nu_k)$, the inequalities*

$$(3.8) \quad \gamma_k \left[\int_{\Delta} G^{(k)} \nu_k \right]^2 \leq D_{k,\omega}(G) \leq \gamma_k \int_{\Delta} |G^{(k)}|^2 \nu_k$$

hold. Moreover, both inequalities in (3.8) become the equalities for $G = \text{const} \cdot P_k$.

Proof. Observe that if $G \in AC^{(k)}(\Delta)$ and $G^{(k)} \in L^2(\nu_k)$, then $G \in H$. Indeed, it follows from (2.1) that

$$\int_{\Delta} G^2 \omega \leq c_1 \int_{\Delta} |R_k G^{(k)}|^2 \omega + c_2 \sum_{j=0}^{k-1} |G^{(j)}(0)|^2 \int_{\Delta} x^{2j} \omega(x) dx.$$

Applying Schwarz's inequality, we obtain

$$|R_k G^{(k)}(x)|^2 \leq \int_{\Delta} |r_k(x, t)| dt \int_{\Delta} |r_k(x, t)| |G^{(k)}(t)|^2 dt \leq c_3 \int_{\Delta} |r_k(x, t)| |G^{(k)}(t)|^2 dt.$$

Thus,

$$\begin{aligned} \int_{\Delta} G^2 \omega &\leq c_4 + c_5 \int_{\Delta} \omega(x) dx \int_{\Delta} |r_k(x, t)| |G^{(k)}(t)|^2 dt \\ &= c_4 + c_5 \int_{\Delta} |G^{(k)}(t)|^2 dt \int_{\Delta} |r_k(x, t)| \omega(x) dx \leq c_4 + c_6 \int_{\Delta} |G^{(k)}(t)|^2 \nu_k(t) dt < \infty. \end{aligned}$$

First we prove the left-hand side of inequality (3.8). It follows from (3.4) and the Bessel inequality that

$$D_{k,\omega}(G) \geq (G, p_k)_H^2$$

(applying (3.1), (3.2) and (3.3))

$$= \frac{1}{\eta_k^2 2^{2k} [k!]^2} \left[\int_{\Delta} G(x) D^k [(1-x)^{\alpha+k} (1+x)^{\beta+k}] dx \right]^2$$

(using (2.1) and the orthogonality of p_k to all polynomials of degree $\leq k-1$)

$$(3.9) \quad = \frac{1}{\eta_k^2 2^{2k} [k!]^2} \left[\int_{\Delta} R_k G^{(k)}(x) D^k [(1-x)^{\alpha+k} (1+x)^{\beta+k}] dx \right]^2.$$

From the elementary upper bound

$$|D^k [(1-x)^{\alpha+k} (1+x)^{\beta+k}]| \leq c_7 \omega(x), \quad x \in \Delta,$$

applying Schwarz's inequality twice, we obtain

$$\begin{aligned} &\int_{\Delta} \int_{\Delta} |r_k(x, t) G^{(k)}(t)| dt |D^k [(1-x)^{\alpha+k} (1+x)^{\beta+k}]| dx \\ &\leq c_8 \left(\int_{\Delta} |G^{(k)}(t)|^2 dt \int_{\Delta} |r_k(x, t)| \omega(x) dx \right)^{1/2} \leq c_9 \left(\int_{\Delta} |G^{(k)}(t)|^2 \nu_k(t) dt \right)^{1/2} < \infty. \end{aligned}$$

Therefore, by Fubini's theorem, we find from (3.9)

$$\begin{aligned} D_{k,\omega}(G) &\geq \frac{1}{\eta_k^2 2^{2k} [k!]^2} \left[\int_{\Delta} G^{(k)}(t) dt \int_{\Delta} r_k(x, t) D^k [(1-x)^{\alpha+k} (1+x)^{\beta+k}] dx \right]^2 \\ &= \frac{1}{\eta_k^2 2^{2k} [k!]^2} \left[\int_{\Delta} G^{(k)}(t) (1-t)^{\alpha+k} (1+t)^{\beta+k} dt \right]^2 \end{aligned}$$

(using the definitions (3.5), (3.6) and (3.7))

$$= \gamma_k \left[\int_{\Delta} G^{(k)} \nu_k \right]^2.$$

Consequently, the left-hand side of inequality (3.8) is proved. If $G = P_k$, then

$$\int_{\Delta} G^{(k)} \nu_k = \int_{\Delta} P_k^{(k)} \nu_k = \frac{1}{\beta_k} \int_{\Delta} P^{(k)}(x) (1-x)^{\alpha+k} (1+x)^{\beta+k} dx$$

(integrating by parts)

$$= \frac{(-1)^k}{\beta_k} \int_{\Delta} P_k(x) D^k [(1-x)^{\alpha+k} (1+x)^{\beta+k}] dx$$

(applying (3.1) and then (3.7))

$$= \frac{2^k k!}{\beta_k} \int_{\Delta} P_k^2 \omega = \frac{2^k k! \eta_k^2}{\beta_k} = \frac{\eta_k}{\gamma_k^{1/2}}.$$

Thus, we have

$$(3.10) \quad \gamma_k \left[\int_{\Delta} P_k^{(k)} \nu_k \right]^2 = \eta_k^2 = D_{k,\omega}(P_k),$$

that is, the left-hand side of inequality (3.8) becomes the equality for $G = \text{const} \cdot P_k$.

Now we show the required upper bound. To this end we observe first that the Jacobi polynomials $P_n(x)$ satisfy the differential equations

$$(3.11) \quad \begin{aligned} (1-x^2)P_n^{(k+1)}(x) + (\beta - \alpha - (\alpha + \beta + 2k)x)P_n^{(k)}(x) \\ + (\alpha + \beta + n + k)(n - k + 1)P_n^{(k-1)}(x) = 0, \quad k \in \mathbb{N}. \end{aligned}$$

It is well known (see [5], Chapter VII, § 3) that every Jacobi polynomial $P_n(x)$ satisfies the differential equation of the second order

$$(3.12) \quad (1-x^2)P_n''(x) + (\beta - \alpha - (\alpha + \beta + 2)x)P_n'(x) + n(\alpha + \beta + n + 1)P_n(x) = 0.$$

Also we have

$$\begin{aligned} D^{k-1}[(1-x^2)P_n''(x)] &= (1-x^2)P_n^{(k+1)}(x) - 2(k-1)xP_n^{(k)}(x) \\ &\quad - (k-1)(k-2)P_n^{(k-1)}(x), \\ D^{k-1}[(\beta - \alpha - (\alpha + \beta + 2)x)P_n'(x)] &= (\beta - \alpha - (\alpha + \beta + 2)x)P_n^{(k)}(x) \\ &\quad - (\alpha + \beta + 2)(k-1)P_n^{(k-1)}(x). \end{aligned}$$

Differentiating (3.12) $k-1$ times and applying these equations, we obtain (3.11). For any natural integer k we define the differential operation

$$(3.13) \quad L_k y := (-1)^k D^k [(1-x)^{\alpha+k} (1+x)^{\beta+k} D^k y].$$

Denote $\tilde{\nu}_k(x) := (1-x)^{\alpha+k} (1+x)^{\beta+k}$. Let us show that for $n \geq k \geq 1$, the equalities

$$(3.14) \quad \begin{aligned} L_1 P_n(x) &= n(\alpha + \beta + n + 1)P_n(x)\omega(x), \\ L_2 P_n(x) &= (\alpha + \beta + n + 2)(\alpha + \beta + n + 1)n(n-1)P_n(x)\omega(x), \\ &\dots \end{aligned}$$

$$(3.15) \quad L_n P_n(x) = (\alpha + \beta + 2n) \dots (\alpha + \beta + n + 1)n!P_n(x)\omega(x)$$

hold. It is easy to see that (3.14) follows from (3.12). For the proof of the other equalities it is sufficient to establish the recurrent formula

$$(3.16) \quad L_{k+1} P_n = (\alpha + \beta + n + k + 1)(n - k)L_k P_n.$$

Write

$$(3.17) \quad L_{k+1} P_n(x) = (-1)^{k+1} D^{k+1} \cdot D \left[\tilde{\nu}_k(x)(1-x^2)P_n^{(k+1)}(x) \right].$$

We have

$$(3.18) \quad D\tilde{\nu}_k(x) = \tilde{\nu}_k(x)(1-x^2)^{-1}(\beta - \alpha - (\alpha + \beta + 2k)x),$$

and also it follows from (3.11) that

$$\begin{aligned} D \left[(1-x^2)P_n^{(k+1)}(x) \right] &= -(\alpha + \beta + n + k + 1)(n - k)P_n^{(k)}(x) \\ &\quad - (\beta - \alpha - (\alpha + \beta + 2k)x)P_n^{(k+1)}(x). \end{aligned}$$

From this, (3.17) and (3.18) follow (3.16) and (3.15). Note that if the equalities (3.14) and (3.15) are rewritten in the form

$$(3.19) \quad L_k P_n = \sigma_{k,n} P_n \omega, \quad n \geq k \geq 1,$$

then

$$(3.20) \quad \sigma_{k,n} \geq \sigma_{k,k} = (\alpha + \beta + 2k) \dots (\alpha + \beta + k + 1)k!$$

Now we show the upper estimate (3.8). Note that it trivially holds for a polynomial of degree $< k$, and for polynomials $y \in \mathbf{P}$ of degree $\geq k$ it is equivalent to the inequality

$$(3.21) \quad \gamma_k \int_{\Delta} |y^{(k)}|^2 \nu_k \geq \int_{\Delta} y^2 \omega,$$

provided

$$(3.22) \quad (y, p_n)_H = 0, \quad n = 0, 1, \dots, k-1.$$

It follows from (3.22) that there exists $n_y \in \mathbb{N}$ such that

$$y(x) = \sum_{n=k}^{n_y} (y, p_n)_H p_n(x).$$

Hence,

$$\gamma_k \int_{\Delta} |y^{(k)}|^2 \nu_k = \frac{\gamma_k}{\beta_k} \int_{\Delta} y L_k y = \frac{\gamma_k}{\beta_k} \int_{\Delta} \left[\sum_{n=k}^{n_y} (y, p_n)_H p_n \right] \left[\sum_{n=k}^{n_y} (y, p_n)_H L_k p_n \right]$$

(because of (3.19))

$$= \frac{\gamma_k}{\beta_k} \sum_{n=k}^{n_y} (y, p_n)_H^2 \sigma_{k,n} \int_{\Delta} p_n^2 \omega$$

(applying (3.20) and Parseval's equality)

$$\geq \frac{\gamma_k}{\beta_k} \sigma_{k,k} \sum_{n=k}^{n_y} (y, p_n)^2 = \frac{\gamma_k}{\beta_k} (\alpha + \beta + 2k) \dots (\alpha + \beta + k + 1)k! \int_{\Delta} y^2 \omega$$

(inserting the values of γ_k and β_k)

$$= \int_{\Delta} y^2 \omega,$$

and (3.21) is proved.

It is well known that Jacobi's polynomials $\{p_n\}$, $n \in \mathbb{N}_0$, form the complete orthonormal system in H . Therefore, $\mathbf{P}$ is dense in H and, since $\tilde{\nu}_k(x)$ has the form (1.3), then $\mathbf{P}$ is dense in $L^2(\nu_k)$. Let $G \in AC^{(k)}(\Delta)$ be such that $G^{(k)} \in L^2(\nu_k)$. For any $\varepsilon > 0$ there exists such a polynomial $h_\varepsilon \in P$ that

$$\|G^{(k)} - h_\varepsilon\|_{L^2(\nu_k)} \leq \varepsilon.$$

Let $g_\varepsilon := G^{(k)} - h_\varepsilon$, $g := G^{(k)} = h_\varepsilon + g_\varepsilon$. By Schwarz's inequality we find

$$(3.23) \quad \begin{aligned} \int_{\Delta} |R_k g_\varepsilon|^2 \omega &\leq \int_{\Delta} \omega(x) dx \int_{\Delta} |r_k(x, s)| ds \int_{\Delta} |r_k(x, t)| |g_\varepsilon(t)|^2 dt \\ &\leq c_{10} \int_{\Delta} |g_\varepsilon(t)|^2 dt \int_{\Delta} |r_k(x, t)| \omega(x) dx \leq c_{11} \int_{\Delta} |g_\varepsilon|^2 \nu_k \leq c_{11} \varepsilon^2. \end{aligned}$$

Put

$$y_\varepsilon := R_k h_\varepsilon - \sum_{j=0}^{k-1} (R_k h_\varepsilon, p_j)_H p_j.$$

Then $y_\varepsilon \in P$ and satisfies the condition (3.22). Using (2.2) we find

$$G - \sum_{j=0}^{k-1} (G, p_j)_H p_j = R_k g - \sum_{j=0}^{k-1} (R_k g, p_j)_H p_j = y_\varepsilon + R_k g_\varepsilon - \sum_{j=0}^{k-1} (R_k g_\varepsilon, p_j)_H p_j.$$

Hence,

$$[D_{k,\omega}(G)]^{1/2} \leq \|y_\varepsilon\|_H + \|R_k g_\varepsilon\|_H$$

(then (3.21) and (3.23) imply)

$$\begin{aligned} &\leq \gamma_k^{1/2} \|y_\varepsilon^{(k)}\|_{L^2(\nu_k)} + c_{12} \cdot \varepsilon = \gamma_k^{1/2} \|G^{(k)} - g_\varepsilon\|_{L^2(\nu_k)} + c_{12} \cdot \varepsilon \\ &\leq \gamma_k^{1/2} \|G^{(k)}\|_{L^2(\nu_k)} + (\gamma_k^{1/2} + c_{12})\varepsilon \end{aligned}$$

and the right-hand side of (3.8) follows by $\varepsilon \rightarrow 0$. If $G = P_k$, then, integrating by parts, we find

$$\begin{aligned} &\gamma_k \int_{\Delta} |P_k^{(k)}|^2 \nu_k = \frac{\gamma_k}{\beta_k} \int_{\Delta} P_k L_k P_k \\ &= \frac{\gamma_k}{\beta_k} (\alpha + \beta + 2k) \dots (\alpha + \beta + k + 1) k! \int_{\Delta} P_k^2 \omega = D_{k,\omega}(P_k). \quad \square \end{aligned}$$

REFERENCES

1. H. Chernoff, *A note on an inequality involving the normal distribution*, Ann. Probab. **9** (1981), 533-535. MR614640 (82f:60050)
2. H. Brascamp and E. Lieb, *On extensions of the Brunn-Minkowski and Prékopa-Leindler theorem, including inequalities for log concave functions, and with an application to the diffusion equation*. J. Funct. Anal. **22** (1976), 366-389. MR0450480 (56:8774)
3. W. Bischoff and M. Fichter, *Optimal lower and upper bounds for the L_p -mean deviation of functions of a random variable*, Math. Methods in Statistics. **9** (2000), 237-269. MR1807094 (2001k:60022)
4. V. D. Stepanov, *An extremal property of Chebyshev polynomials*, Proc. Steklov Inst. Math. **248** (2005), 230-242. MR2165931 (2006h:41006)
5. P. K. Suetin, *Classical orthogonal polynomials* (in Russian), Nauka, Moscow, 1979. MR548727 (80h:33001)
6. G. Aleksich, *The convergence problems of orthogonal series* (in Russian), Izd. Inostr. Lit., Moscow, 1963. MR0218828 (36:1912)

PEOPLE FRIENDSHIP UNIVERSITY, MIKLUKHO-MAKLAI 6, MOSCOW, 117198, RUSSIA

E-mail address: vstepanov@sci.pfu.edu.ru