SIGNED q -ANALOGS OF TORNHEIM'S DOUBLE SERIES

XIA ZHOU, TIANXIN CAI, AND DAVID M. BRADLEY

(Communicated by Ken Ono)

ABSTRACT. We introduce signed q -analogs of Tornheim's double series and evaluate them in terms of double q -Euler sums. As a consequence, we provide explicit evaluations of signed and unsigned Tornheim double series and correct some mistakes in the literature.

1. INTRODUCTION

Let k be a positive integer. Sums of the form
 (1)

$$\zeta(s_1, s_2, \dots, s_k) := \sum_{n_1 > n_2 > \dots > n_k > 0} \prod_{j=1}^k n_j^{-s_j}, \quad \sum_{j=1}^m \Re(s_j) > m, \quad m = 1, 2, \dots, k,$$

have attracted increasing attention in recent years; see e.g. [2, 3, 4, 5, 6, 8, 9, 10, 11, 13, 16, 24, 26, 27, 29, 30, 31]. The survey articles [7, 18, 19, 36, 37, 38] provide an extensive list of references. In (1) the sum is over all positive integers $n_1, n_2, \dots, n_k$ satisfying the indicated inequalities. Of course (1) reduces to the familiar Riemann zeta function when $k = 1$. When the arguments are all positive integers, we refer to (1) as a multiple zeta value and note that in this case, $s_1 > 1$ is necessary and sufficient for convergence.

The problem of evaluating $\zeta(s_1, s_2)$ with integers $s_1 > 1$ and $s_2 > 0$ seems to have been first proposed in a letter from Goldbach to Euler [21] in 1742. (See also [20, 22] and [1, p. 253].) Calculating several examples led Euler to infer a closed form evaluation in terms of values of the Riemann zeta function in the case when $s_1 + s_2$ is odd. In [4], Borwein, Bradley and Broadhurst considered the more general Euler sum

$$(2) \quad \zeta(s_1, s_2, \dots, s_k; \sigma_1, \sigma_2, \dots, \sigma_k) := \sum_{n_1 > n_2 > \dots > n_k > 0} \prod_{j=1}^k \sigma_j^{n_j} n_j^{-s_j}$$

with each $\sigma_j \in \{-1, 1\}$. Among the many other results for (2) listed therein is an explicit formula for the case $k = 2$ that reduces to Euler's evaluation when $\sigma_1 = \sigma_2 = 1$. For the case of arbitrary k , several infinite classes of closed form evaluations

Received by the editors January 19, 2007, and in revised form April 25, 2007.

2000 *Mathematics Subject Classification.* Primary 11M41; Secondary 11M06, 05A30, 33E20, 30B50.

Key words and phrases. Tornheim's double series, alternating Euler sums, multiple harmonic series, multiple zeta values, q -analog, q -series.

©2008 American Mathematical Society
 Reverts to public domain 28 years from publication

for (1) and (2) are proved in [5, 6, 8, 9, 10]. Subsequently, Bradley [12, 13, 14, 15] found q -analogs of (1) and (2) and thoroughly investigated their properties.

In [32] Tornheim introduced the double series

$$T(r, s, t) := \sum_{u, v=1}^{\infty} \frac{1}{u^r v^s (u+v)^t},$$

with nonnegative integers r, s and t satisfying $r+t > 1$, $s+t > 1$ and $r+s+t > 2$. Huard, Williams and Zhang [25] evaluated Tornheim's series in terms of sums of products of Riemann zeta values when $r+s+t$ is odd. Subbarao and Sitaramachandra Rao [28] considered the alternating variants

$$R(r, s, t) := \sum_{u, v=1}^{\infty} \frac{(-1)^v}{u^r v^s (u+v)^t}, \quad S(r, s, t) := \sum_{u, v=1}^{\infty} \frac{(-1)^{u+v}}{u^r v^s (u+v)^t},$$

and posed the problem to evaluate $R(r, r, r)$ and $S(r, r, r)$ for any positive integer r . Tsumura [33, Cor. 3] and [34, Theorem 3.6] tackled this problem and later [35] evaluated $S(r, s, t)$ for any positive integers r, s, t such that $r+s+t$ is odd. Tsumura's method is elementary but complicated, and also has some mistakes. In this paper, we give a simple formula that expresses R, S and T in terms of double Euler sums. In light of the aforementioned formula of Borwein, Bradley and Broadhurst for the double Euler sums, our results yield a closed form evaluation for the series R, S and T whenever the arguments r, s and t are positive integers with $r+s+t$ odd. More generally, we consider q -analogs of R, S, T and show how they may all be evaluated in terms of double q -Euler sums.

2. q -ANALOGS

Henceforth assume q is real and $q > 1$. The q -analog of a positive integer n is

$$[n]_q := \sum_{j=0}^{n-1} q^j = \frac{q^n - 1}{q - 1}.$$

Let k be a positive integer, let $s_1, s_2, \dots, s_k$ be real numbers, and let $\sigma_1, \sigma_2, \dots, \sigma_k \in \{-1, 1\}$. Define the q -Euler sum

$$(3) \quad \zeta_q[s_1, s_2, \dots, s_k; \sigma_1, \sigma_2, \dots, \sigma_k] := \sum_{n_1 > n_2 > \dots > n_k > 0} \prod_{j=1}^k \frac{\sigma_j^{n_j} q^{(s_j-1)n_j}}{[n_j]_q^{s_j}}$$

and note that this coincides with the special case $\text{Li}_{s_1, \dots, s_k}[\sigma_1 q^{s_1-1}, \dots, \sigma_k q^{s_k-1}]$ of the multiple q -polylogarithm [12, eq. (6.2)]. If $\sigma_j = 1$ for each $j = 1, 2, \dots, k$, then we recover the multiple q -zeta value $\zeta[s_1, s_2, \dots, s_k]$ of [12, 14, 15].

Let $\sigma, \tau \in \{-1, 1\}$. Define q -analogs of the signed and unsigned Tornheim double series R, S and T by

$$T[r, s, t; \sigma, \tau] := \sum_{u, v=1}^{\infty} \frac{\sigma^u \tau^v q^{(r+t-1)u + (s+t-1)v}}{[u]_q^r [v]_q^s [u+v]_q^t}.$$

The sum

$$\varphi[s; \sigma] := \sum_{n=1}^{\infty} \frac{(n-1)\sigma^n q^{(s-1)n}}{[n]_q^s} = \sum_{n=1}^{\infty} \frac{n\sigma^n q^{(s-1)n}}{[n]_q^s} - \zeta_q[s; \sigma],$$

the case $\sigma = 1$ of which was defined in [15], will also be needed. As in [4], it is convenient to combine signs and exponents in (2) and (3) into a single list by writing s_j if $\sigma_j = 1$ and $\overline{s_j}$ if $\sigma_j = -1$. For consistency one may do this also for T and φ ; thus for example, $\varphi[s; 1] = \varphi[s]$, $\varphi[s; -1] = \varphi[\overline{s}]$, $R(r, s, t) = T(r, \overline{s}, t)$ and $S(r, s, t) = T(\overline{r}, \overline{s}, t)$. We also employ the notation

$$\binom{a+b+c}{a, b} := \frac{(a+b+c)!}{a!b!c!}$$

for the multinomial coefficient, in which a , b and c are all nonnegative integers. The following theorem shows how the q -analogs of R , S and T are related to the q -Euler sums.

Theorem 1. *Let r and s be positive integers, and let t be a real number. Then*

$$\begin{aligned} T[r, s, t] &= \sum_{a=0}^{r-1} \sum_{b=0}^{r-1-a} \binom{a+s-1}{a, b} (1-q)^b \zeta_q[s+t+a, r-a-b] \\ &\quad + \sum_{a=0}^{s-1} \sum_{b=0}^{s-1-a} \binom{a+r-1}{a, b} (1-q)^b \zeta_q[r+t+a, s-a-b] \\ &\quad - \sum_{j=1}^{\min(r,s)} \binom{r+s-j-1}{r-j, s-j} (1-q)^j \varphi[r+s+t-j], \\ T[\overline{r}, \overline{s}, t] &= \sum_{a=0}^{r-1} \sum_{b=0}^{r-1-a} \binom{a+s-1}{a, b} (1-q)^b \zeta_q[\overline{s+t+a}, r-a-b] \\ &\quad + \sum_{a=0}^{s-1} \sum_{b=0}^{s-1-a} \binom{a+r-1}{a, b} (1-q)^b \zeta_q[\overline{r+t+a}, s-a-b] \\ &\quad - \sum_{j=1}^{\min(r,s)} \binom{r+s-j-1}{r-j, s-j} (1-q)^j \varphi[\overline{r+s+t-j}], \\ T[r, \overline{s}, t] &= \sum_{a=0}^{r-1} \sum_{b=0}^{r-1-a} \binom{a+s-1}{a, b} (1-q)^b \zeta_q[\overline{s+t+a}, \overline{r-a-b}] \\ &\quad + \sum_{a=0}^{s-1} \sum_{b=0}^{s-1-a} \binom{a+r-1}{a, b} (1-q)^b \zeta_q[r+t+a, \overline{s-a-b}] \\ &\quad - \sum_{j=1}^{\min(r,s)} \binom{r+s-j-1}{r-j, s-j} (1-q)^j (1+q)^{j-r-s-t} \zeta_{q^2}[r+s+t-j]. \end{aligned}$$

Taking the limit as $q \rightarrow 1+$ in Theorem 1 and noting the restrictions on r , s and t now needed for convergence yield the following.

Corollary 1. *Let r and s be positive integers and let t be a real number. If $r+t > 1$ and $s+t > 1$, then*

$$T(r, s, t) = \sum_{a=0}^{r-1} \binom{a+s-1}{s-1} \zeta(s+t+a, r-a) + \sum_{a=0}^{s-1} \binom{a+r-1}{r-1} \zeta(r+t+a, s-a);$$

if $r + t > 0$ and $s + t > 0$, then

$$S(r, s, t) = \sum_{a=0}^{r-1} \binom{a+s-1}{s-1} \zeta(\overline{s+t+a}, r-a) + \sum_{a=0}^{s-1} \binom{a+r-1}{r-1} \zeta(\overline{r+t+a}, s-a);$$

if $r + t > 1$ and $s + t > 0$, then

$$R(r, s, t) = \sum_{a=0}^{r-1} \binom{a+s-1}{s-1} \zeta(\overline{s+t+a}, \overline{r-a}) + \sum_{a=0}^{s-1} \binom{a+r-1}{r-1} \zeta(r+t+a, \overline{s-a}).$$

Putting $t = 0$ in Theorem 1 yields the following decomposition formulas, the first of which was given under slightly more restrictive hypotheses in [15, Theorem 2.1].

Corollary 2. *If r and s are positive integers, then*

$$\begin{aligned} \zeta_q[r] \zeta_q[s] &= \sum_{a=0}^{r-1} \sum_{b=0}^{r-1-a} \binom{a+s-1}{s-1} \binom{s-1}{b} (1-q)^b \zeta_q[s+a, r-a-b] \\ &\quad + \sum_{a=0}^{s-1} \sum_{b=0}^{s-1-a} \binom{a+r-1}{r-1} \binom{r-1}{b} (1-q)^b \zeta_q[r+a, s-a-b] \\ &\quad - \sum_{j=1}^{\min(r,s)} \binom{r+s-j-1}{r-j, s-j} (1-q)^j \varphi[r+s-j]; \\ \zeta_q[\overline{r}] \zeta_q[\overline{s}] &= \sum_{a=0}^{r-1} \sum_{b=0}^{r-1-a} \binom{a+s-1}{s-1} \binom{s-1}{b} (1-q)^b \zeta_q[\overline{s+a}, r-a-b] \\ &\quad + \sum_{a=0}^{s-1} \sum_{b=0}^{s-1-a} \binom{a+r-1}{r-1} \binom{r-1}{b} (1-q)^b \zeta_q[\overline{r+a}, s-a-b] \\ &\quad - \sum_{j=1}^{\min(r,s)} \binom{r+s-j-1}{r-j, s-j} (1-q)^j \varphi[\overline{r+s-j}]; \\ \zeta_q[r] \zeta_q[\overline{s}] &= \sum_{a=0}^{r-1} \sum_{b=0}^{r-1-a} \binom{a+s-1}{s-1} \binom{s-1}{b} (1-q)^b \zeta_q[\overline{s+a}, \overline{r-a-b}] \\ &\quad + \sum_{a=0}^{s-1} \sum_{b=0}^{s-1-a} \binom{a+r-1}{r-1} \binom{r-1}{b} (1-q)^b \zeta_q[r+a, \overline{s-a-b}] \\ &\quad - \sum_{j=1}^{\min(r,s)} \binom{r+s-j-1}{r-j, s-j} (1-q)^j (1+q)^{j-r-s} \zeta_{q^2}[r+s-j]. \end{aligned}$$

Taking the limit as $q \rightarrow 1+$ in Corollary 2 and noting the additional restrictions needed on r and s to guarantee convergence in this case yield the following decomposition formulas, the first of which was known to Euler.

Corollary 3. *If $r-1$ and $s-1$ are positive integers, then*

$$\zeta(r) \zeta(s) = \sum_{a=0}^{r-1} \binom{a+s-1}{s-1} \zeta(s+a, r-a) + \sum_{a=0}^{s-1} \binom{a+r-1}{r-1} \zeta(r+a, s-a);$$

if r and s are positive integers, then

$$\zeta(\overline{r})\zeta(\overline{s}) = \sum_{a=0}^{r-1} \binom{a+s-1}{s-1} \zeta(\overline{s+a}, r-a) + \sum_{a=0}^{s-1} \binom{a+r-1}{r-1} \zeta(\overline{r+a}, s-a);$$

if $r-1$ and s are positive integers, then

$$\zeta(r)\zeta(\overline{s}) = \sum_{a=0}^{r-1} \binom{a+s-1}{s-1} \zeta(\overline{s+a}, \overline{r-a}) + \sum_{a=0}^{s-1} \binom{a+r-1}{r-1} \zeta(r+a, \overline{s-a}).$$

3. PROOF OF THEOREM 1

The key ingredient is the following partial fraction decomposition.

Lemma 1. *If r , s , u , and v are all positive integers, then*

$$\begin{aligned} \frac{1}{[u]_q^r [v]_q^s} &= \sum_{a=0}^{r-1} \sum_{b=0}^{r-1-a} \binom{a+s-1}{a, b} \frac{(1-q)^b q^{(s-1-b)u+av}}{[u]_q^{r-a-b} [u+v]_q^{s+a}} \\ &\quad + \sum_{a=0}^{s-1} \sum_{b=0}^{s-1-a} \binom{a+r-1}{a, b} \frac{(1-q)^b q^{au+(r-1-b)v}}{[v]_q^{s-a-b} [u+v]_q^{r+a}} \\ &\quad - \sum_{j=1}^{\min(r,s)} \binom{r+s-j-1}{r-j, s-j} \frac{(1-q)^j q^{(s-j)u+(r-j)v}}{[u+v]_q^{r+s-j}}. \end{aligned}$$

Proof. Let x and y be nonzero real numbers such that $x+y+(q-1)xy \neq 0$. As in [15], observe that if we apply the partial differential operator

$$\frac{1}{(r-1)!} \left(-\frac{\partial}{\partial x} \right)^{r-1} \frac{1}{(s-1)!} \left(-\frac{\partial}{\partial y} \right)^{s-1}$$

to both sides of the identity

$$\frac{1}{xy} = \frac{1}{x+y+(q-1)xy} \left(\frac{1}{x} + \frac{1}{y} + q-1 \right),$$

then we obtain the identity [15, Lemma 3.1]

$$\begin{aligned} \frac{1}{x^r y^s} &= \sum_{a=0}^{r-1} \sum_{b=0}^{r-1-a} \binom{a+s-1}{a, b} \frac{(1-q)^b (1+(q-1)y)^a (1+(q-1)x)^{s-1-b}}{x^{r-a-b} (x+y+(q-1)xy)^{s+a}} \\ &\quad + \sum_{a=0}^{s-1} \sum_{b=0}^{s-1-a} \binom{a+r-1}{a, b} \frac{(1-q)^b (1+(q-1)x)^a (1+(q-1)y)^{r-1-b}}{y^{s-a-b} (x+y+(q-1)xy)^{r+a}} \\ &\quad - \sum_{j=1}^{\min(r,s)} \binom{r+s-j-1}{r-j, s-j} \frac{(1-q)^j (1+(q-1)y)^{r-j} (1+(q-1)x)^{s-j}}{(x+y+(q-1)xy)^{r+s-j}}. \end{aligned}$$

Now let $x = [u]_q$, $y = [v]_q$ and note that then $1+(q-1)x = q^u$, $1+(q-1)y = q^v$ and $x+y+(q-1)xy = [u+v]_q$. $\square$

To prove Theorem 1, multiply both sides of Lemma 1 by

$$\frac{\sigma^u \tau^v q^{(r+t-1)u+(s+t-1)v}}{[u+v]_q^t}$$

and sum over all ordered pairs of positive integers (u, v) to obtain

$$\begin{aligned} T[r, s, t; \sigma, \tau] &= \sum_{a=0}^{r-1} \sum_{b=0}^{r-1-a} \binom{a+s-1}{a, b} (1-q)^b \sum_{u,v=1}^{\infty} \frac{\sigma^u \tau^v q^{(r-a-b-1)u} q^{(s+t+a-1)(u+v)}}{[u]_q^{r-a-b} [u+v]_q^{s+t+a}} \\ &\quad + \sum_{a=0}^{s-1} \sum_{b=0}^{s-1-a} \binom{a+r-1}{a, b} (1-q)^b \sum_{u,v=1}^{\infty} \frac{\sigma^u \tau^v q^{(s-a-b-1)v} q^{(r+t+a-1)(u+v)}}{[v]_q^{s-a-b} [u+v]_q^{r+t+a}} \\ &\quad - \sum_{j=1}^{\min(r,s)} \binom{r+s-j-1}{r-j, s-j} (1-q)^j \sum_{u,v=1}^{\infty} \frac{\sigma^u \tau^v q^{(r+s+t-j-1)(u+v)}}{[u+v]_q^{r+s+t-j}}. \end{aligned}$$

It follows that

$$\begin{aligned} T[r, s, t; \sigma, \sigma] &= \sum_{a=0}^{r-1} \sum_{b=0}^{r-1-a} \binom{a+s-1}{a, b} (1-q)^b \sum_{m>n>0} \frac{\sigma^m q^{(s+t+a-1)m} q^{(r-a-b-1)n}}{[m]_q^{s+t+a} [n]_q^{r-a-b}} \\ &\quad + \sum_{a=0}^{s-1} \sum_{b=0}^{s-1-a} \binom{a+r-1}{a, b} (1-q)^b \sum_{m>n>0} \frac{\sigma^m q^{(r+t+a-1)m} q^{(s-a-b-1)n}}{[m]_q^{r+t+a} [n]_q^{s-a-b}} \\ &\quad - \sum_{j=1}^{\min(r,s)} \binom{r+s-j-1}{r-j, s-j} (1-q)^j \sum_{m>n>0} \frac{\sigma^m q^{(r+s+t-j-1)m}}{[m]_q^{r+s+t-j}} \\ &= \sum_{a=0}^{r-1} \sum_{b=0}^{r-1-a} \binom{a+s-1}{a, b} (1-q)^b \zeta_q[s+t+a, r-a-b; \sigma, 1] \\ &\quad + \sum_{a=0}^{s-1} \sum_{b=0}^{s-1-a} \binom{a+r-1}{a, b} (1-q)^b \zeta_q[r+t+a, s-a-b; \sigma, 1] \\ &\quad - \sum_{j=1}^{\min(r,s)} \binom{r+s-j-1}{r-j, s-j} (1-q)^j \varphi[r+s+t-j; \sigma], \end{aligned}$$

and also that

$$\begin{aligned} T[r, s, t; 1, -1] &= \sum_{a=0}^{r-1} \sum_{b=0}^{r-1-a} \binom{a+s-1}{a, b} (1-q)^b \\ &\quad \times \sum_{m>n>0} \frac{(-1)^m q^{(s+t+a-1)m} (-1)^n q^{(r-a-b-1)n}}{[m]_q^{s+t+a} [n]_q^{r-a-b}} \\ &\quad + \sum_{a=0}^{s-1} \sum_{b=0}^{s-1-a} \binom{a+r-1}{a, b} (1-q)^b \sum_{m>n>0} \frac{q^{(r+t+a-1)m} (-1)^n q^{(s-a-b-1)n}}{[m]_q^{r+t+a} [n]_q^{s-a-b}} \\ &\quad - \sum_{j=1}^{\min(r,s)} \binom{r+s-j-1}{r-j, s-j} (1-q)^j \sum_{m>n>0} \frac{(-1)^n q^{(r+s+t-j-1)m}}{[m]_q^{r+s+t-j}} \\ &= \sum_{a=0}^{r-1} \sum_{b=0}^{r-1-a} \binom{a+s-1}{a, b} (1-q)^b \zeta_q[s+t+a, r-a-b; \sigma, 1] \end{aligned}$$

$$\begin{aligned}
& + \sum_{a=0}^{s-1} \sum_{b=0}^{s-1-a} \binom{a+r-1}{a, b} (1-q)^b \zeta_q[r+t+a, s-a-b; \sigma, 1] \\
& + \sum_{j=1}^{\min(r,s)} \binom{r+s-j-1}{r-j, s-j} (1-q)^j \sum_{m>0} \frac{q^{(r+s+t-j-1)2m}}{[2m]_q^{r+s+t-j}}.
\end{aligned}$$

Since

$$[2m]_q = \frac{q^{2m} - 1}{q - 1} = \frac{q^2 - 1}{q - 1} \cdot \frac{q^{2m} - 1}{q^2 - 1} = (q + 1)[m]_{q^2},$$

the proof of Theorem 1 is complete. $\square$

4. EXAMPLES

Again, let $\sigma, \tau \in \{-1, 1\}$. It is known [4, 23] that if s and t are positive integers such that $s + t$ is odd and $s > (1 + \sigma)/2$, then $\zeta(s, t; \sigma, \tau)$ lies in the polynomial ring $\mathbf{Q}[\{\zeta(k; \pm 1) : k \in \mathbf{Z}, 2 \leq k \leq s + t\} \cup \{\zeta(1; -1)\}]$. Since

$$\zeta(k; -1) = \zeta(\overline{k}) = \begin{cases} (2^{1-k} - 1)\zeta(k) & \text{if } k > 1, \\ -\log 2 & \text{if } k = 1, \end{cases}$$

it is clear that in fact, $\zeta(s, t; \sigma, \tau) \in \mathbf{Q}[\{\zeta(k) : k \in \mathbf{Z}, 2 \leq k \leq s + t\} \cup \{-\log 2\}]$. It follows that if r, s, t satisfy the conditions of Corollary 1 and if in addition $r + s + t$ is an odd integer, then the signed and unsigned double Tornheim series $R(r, s, t)$, $S(r, s, t)$ and $T(r, s, t)$ also lie in this ring. It is possible to evaluate these series explicitly if we recall the following formula from [4, eq. (75)].

Proposition 1. *Let $\sigma, \tau \in \{-1, 1\}$, and let s and t be positive integers such that $s + t$ is odd, $s > (1 + \sigma)/2$, and $t > (1 + \tau)/2$. Then*

$$\begin{aligned}
(4) \quad \zeta(s, t; \sigma, \tau) &= \frac{1}{2}(1 + (-1)^s)\zeta(s; \sigma)\zeta(t; \tau) - \frac{1}{2}\zeta(s + t; \sigma\tau) \\
&+ (-1)^t \sum_{0 \leq k \leq t/2} \binom{s+t-2k-1}{s-1} \zeta(2k; \sigma\tau)\zeta(s+t-2k; \sigma) \\
&+ (-1)^t \sum_{0 \leq k \leq s/2} \binom{s+t-2k-1}{t-1} \zeta(2k; \sigma\tau)\zeta(s+t-2k; \tau).
\end{aligned}$$

In Proposition 1, it is understood that $\zeta(0; \sigma\tau) = -1/2$ in accordance with the analytic continuation of $s \mapsto \zeta(s; \sigma\tau)$. The restriction $t > (1 + \tau)/2$ can be removed if in (4) we interpret $\zeta(1; 1) = 0$ wherever it occurs. That is, if $\sigma \in \{-1, 1\}$ and s is an even positive integer, then

$$(5) \quad \zeta(s, 1; \sigma, 1) = \frac{1}{2}(s-1)\zeta(s+1; \sigma) + \frac{1}{2}\zeta(s+1) - \sum_{k=1}^{(s/2)-1} \zeta(2k; \sigma)\zeta(s+1-2k).$$

The case $\sigma = 1$ of (5) is subsumed by another formula [4, eq. (31)] of Euler, namely

$$(6) \quad \zeta(s, 1) = \frac{s}{2}\zeta(s+1) - \frac{1}{2} \sum_{k=2}^{s-1} \zeta(k)\zeta(s+1-k),$$

which is valid for *all* integers $s > 1$, not just for even s .

In [35], Tsumura listed evaluation formulas for $S(r, s, t)$ when $r + s + t \leq 9$ is odd. From Corollary 1, Proposition 1 and equation (5), we can deduce explicit

formulas for $R(r, s, t)$, $S(r, s, t)$ and $T(r, s, t)$ when $r + s + t$ is odd. In particular, we have the following new results:

$$R(1, 1, 1) = -\frac{5}{8}\zeta(3),$$

$$R(1, 1, 3) = \frac{1}{16}\pi^2\zeta(3) - \frac{27}{32}\zeta(5),$$

$$R(1, 2, 2) = \frac{5}{48}\pi^2\zeta(3) - \frac{3}{2}\zeta(5),$$

$$R(1, 3, 1) = \frac{1}{12}\pi^2\zeta(3) - \frac{59}{32}\zeta(5),$$

$$R(2, 1, 2) = -\frac{5}{16}\pi^2\zeta(3) + \frac{107}{32}\zeta(5),$$

$$R(2, 2, 1) = -\frac{5}{24}\pi^2\zeta(3) + \frac{59}{32}\zeta(5),$$

$$R(3, 1, 1) = \frac{1}{8}\pi^2\zeta(3) - \frac{59}{32}\zeta(5),$$

$$S(5, 5, 5) = \frac{7}{73728}\pi^4\zeta(11) + \frac{35}{24576}\pi^2\zeta(13) + \frac{63}{8192}\zeta(15),$$

$$S(7, 7, 7) = \frac{31}{35389440}\pi^6\zeta(15) + \frac{49}{1966080}\pi^4\zeta(17) + \frac{77}{262144}\pi^2\zeta(19) + \frac{429}{262144}\zeta(21).$$

The values of $R(5, 5, 5)$, $R(7, 7, 7)$ and $R(9, 9, 9)$ listed in [34] appear to be incorrect. They should be

$$R(5, 5, 5) = \frac{16375}{147456}\pi^4\zeta(11) + \frac{573335}{49152}\pi^2\zeta(13) - \frac{2064195}{16384}\zeta(15),$$

$$R(7, 7, 7) = \frac{1048543}{70778880}\pi^6\zeta(15) + \frac{7339969}{3932160}\pi^4\zeta(17) + \frac{80740121}{524288}\pi^2\zeta(19) \\ - \frac{899676921}{524288}\zeta(21),$$

$$R(9, 9, 9) = \frac{13421747}{7046430720}\pi^8\zeta(19) + \frac{738197141}{2113929216}\pi^6\zeta(21) + \frac{1919313253}{67108864}\pi^4\zeta(23) \\ + \frac{143948506845}{67108864}\pi^2\zeta(25) - \frac{1631416447375}{67108864}\zeta(27).$$

REFERENCES

- [1] B. Berndt, *Ramanujan's Notebooks. Part I*, Springer-Verlag, New York, 1985. MR0781125 (86c:01062)
- [2] D. Borwein, J. M. Borwein, and D. M. Bradley, Parametric Euler sum identities, *J. Math. Anal. Appl.*, **316** (2006), no. 1, 328–338. MR2201764 (2007b:11132)
- [3] J. M. Borwein and D. M. Bradley, Thirty-two Goldbach variations, *Internat. J. Number Theory*, **2** (2006), no. 1, 65–103. MR2217795 (2007e:11109)
- [4] J. M. Borwein, D. M. Bradley, and D. J. Broadhurst, Evaluations of k -fold Euler/Zagier sums: A compendium of results for arbitrary k , *Electron. J. Combin.*, **4** (1997), no. 2, #R5. Wilf Festschrift. MR1444152 (98b:11091)
- [5] J. M. Borwein, D. M. Bradley, D. J. Broadhurst, and P. Lisoněk, Combinatorial aspects of multiple zeta values, *Electron. J. Combin.*, **5** (1998), no. 1, #R38. MR1637378 (99g:11100)
- [6] ———, Special values of multiple polylogarithms, *Trans. Amer. Math. Soc.*, **353** (2001), no. 3, 907–941. MR1709772 (2003i:33003)
- [7] D. Bowman and D. M. Bradley, Multiple polylogarithms: A brief survey, *Proceedings of a Conference on q -Series with Applications to Combinatorics, Number Theory and Physics* (B. C. Berndt and K. Ono, eds., Urbana, IL, 2000), Contemporary Math., **291**, Amer. Math. Soc., Providence, RI, 2001, pp. 71–92. MR1874522 (2003c:33021)

- [8] ———, The algebra and combinatorics of shuffles and multiple zeta values, *J. Combin. Theory Ser. A*, **97** (2002), no. 1, 43–61. MR1879045 (2003j:05010)
- [9] ———, Resolution of some open problems concerning multiple zeta evaluations of arbitrary depth, *Compositio Math.*, **139** (2003), no. 1, 85–100. MR2024966 (2005f:11196)
- [10] D. Bowman, D. M. Bradley, and J. Ryoo, Some multi-set inclusions associated with shuffle convolutions and multiple zeta values, *European J. Combin.*, **24** (2003), no. 1, 121–127. MR1957970 (2004c:11115)
- [11] D. M. Bradley, Partition identities for the multiple zeta function, *Zeta Functions, Topology, and Quantum Physics*, Developments in Mathematics, **14**, T. Aoki, S. Kanemitsu, M. Nakahara, Y. Ohno (eds.), Springer, New York, 2005, pp. 19–29. MR2179270 (2006f:11105)
- [12] ———, Multiple q -zeta values, *J. Algebra*, **283** (2005), no. 2, 752–798. MR2111222 (2006f:11106)
- [13] ———, Duality for finite multiple harmonic q -series, *Discrete Math.*, **300** (2005), no. 1–3, 44–56. MR2170113 (2006m:05019)
- [14] ———, On the sum formula for multiple q -zeta values, *Rocky Mountain J. Math.*, to appear. <http://arxiv.org/abs/math.QA/0411274>
- [15] ———, A q -analog of Euler's decomposition formula for the double zeta function, *Internat. J. Math. Math. Sci.*, **2005**, no. 21, 3453–3458. MR2206867 (2006k:11174)
- [16] D. J. Broadhurst and D. Kreimer, Association of multiple zeta values with positive knots via Feynman diagrams up to 9 loops, *Phys. Lett. B*, **393** (1997), no. 3–4, 403–412. MR1435933 (98g:11101)
- [17] F. C. Brown, Périodes des espaces des modules $\overline{\mathcal{M}}_{0,n}$ et valeurs zêtas multiples [Multiple zeta values and periods of the moduli spaces $\overline{\mathcal{M}}_{0,n}$], *C. R. Math. Acad. Sci. Paris*, **342** (2006), no. 12, 949–954. MR2235616
- [18] P. Cartier, Fonctions polylogarithmes, nombres polyzêtas et groupes pro-unipotents, *Astérisque*, **282** (2002), viii, 137–173, Séminaire Bourbaki, 53ième année, 2000–2001, Exp. No. 885. MR1975178 (2004i:19005)
- [19] ———, Values of the ζ -function, *Surveys in Modern Mathematics*, 260–273, London Math. Soc. Lecture Note Ser., **321**, Cambridge Univ. Press, Cambridge, 2005. MR2166932 (2006g:11182)
- [20] L. Euler, *Meditationes Circa Singulare Serierum Genus*, Novi Comm. Acad. Sci. Petropol., **20** (1775), 140–186. Reprinted in “Opera Omnia,” ser. I, **15**, B. G. Teubner, Berlin (1927), 217–267.
- [21] ———, *Briefwechsel*, vol. 1, Birkhäuser, Basel, 1975. MR0497632
- [22] L. Euler and C. Goldbach, *Briefwechsel 1729–1764*, Akademie-Verlag, Berlin, 1965.
- [23] P. Flajolet and B. Salvy, Euler sums and contour integral representations, *Experiment. Math.*, **7** (1998), no. 1, 15–35. MR1618286 (99c:11110)
- [24] M. E. Hoffman, Periods of mirrors and multiple zeta values, *Proc. Amer. Math. Soc.*, **130** (2002), no. 4, 971–974. MR1873769 (2002k:14068)
- [25] J. G. Huard, K. S. Williams, and N.-Y. Zhang, On Tornheim's double series, *Acta Arith.*, **75** (1996), no. 2, 105–117. MR1379394 (97f:11073)
- [26] K. Ihara and T. Takamuki, The quantum $\mathfrak{g}_2$ invariant and relations of multiple zeta values, *J. Knot Theory Ramifications*, **10** (2001), no. 7, 983–997. MR1867104 (2002m:57016)
- [27] T. Q. T. Le and J. Murakami, Kontsevich's integral for the Homfly polynomial and relations between values of multiple zeta functions, *Topology Appl.*, **62** (1995), no. 2, 193–206. MR1320252 (96c:57017)
- [28] M. V. Subbarao and R. Sitaramachandra Rao, On some infinite series of L. J. Mordell and their analogues, *Pacific J. Math.*, **119** (1985), no. 1, 245–255. MR0797027 (87c:11091)
- [29] T. Terasoma, Selberg integrals and multiple zeta values, *Compositio Math.*, **133** (2002), no. 1, 1–24. MR1918286 (2003k:11142)
- [30] ———, Mixed Tate motives and multiple zeta values, *Invent. Math.*, **149** (2002), no. 2, 339–369. MR1918675 (2003h:11073)
- [31] ———, Period integrals and multiple zeta values, *Sūgaku*, **57** (2005), no. 3, 255–266. MR2163671 (2006h:11110)
- [32] L. Tornheim, Harmonic double series, *Amer. J. Math.*, **72** (1950), 303–314. MR0034860 (11:654a)
- [33] H. Tsumura, On some combinatorial relations for Tornheim's double series, *Acta Arith.*, **105** (2002), no. 3, 239–252. MR1931792 (2003i:11134)

- [34] ———, On alternating analogues of Tornheim's double series, *Proc. Amer. Math. Soc.*, **131** (2003), no. 12, 3633–3641. MR1998168 (2004e:11102)
- [35] ———, Evaluation formulas for Tornheim's type of alternating double series, *Math. Comp.*, **73** (2004), no. 245, 251–258. MR2034120 (2005d:11137)
- [36] M. Waldschmidt, Valeurs zêta multiples: Une introduction, *J. Théor. Nombres Bordeaux*, **12** (2000), no. 2, 581–595. MR1823204 (2002a:11106)
- [37] ———, Multiple polylogarithms: An introduction, in *Number Theory and Discrete Mathematics* (Chandigarh, 2000), Trends Math., Birkhäuser, Basel, 2002, pp. 1–12. MR1952273 (2004d:33003)
- [38] V. V. Zudilin, Algebraic relations for multiple zeta values (Russian), *Uspekhi Mat. Nauk*, **58** (2003), no. 1, 3–32; translation in *Russian Math. Surveys*, **58** (2003), no. 1, 1–29. MR1992130 (2004k:11150)

DEPARTMENT OF MATHEMATICS, ZHEJIANG UNIVERSITY, HANGZHOU, 310027, PEOPLE'S REPUBLIC OF CHINA

E-mail address: xiazhou0821@hotmail.com

DEPARTMENT OF MATHEMATICS, ZHEJIANG UNIVERSITY, HANGZHOU, 310027, PEOPLE'S REPUBLIC OF CHINA

E-mail address: txcai@mail.hz.zj.cn

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF MAINE, 5752 NEVILLE HALL, ORONO, MAINE 04469-5752

E-mail address: bradley@math.umaine.edu

E-mail address: dbradley@member.ams.org