THE PROJECTIVE π -CHARACTER BOUNDS THE ORDER OF A π -BASE

ISTVÁN JUHÁSZ AND ZOLTÁN SZENTMIKLÓSSY

(Communicated by Alexander N. Dranishnikov)

ABSTRACT. All spaces below are Tychonov. We define the projective π -character $p\pi\chi(X)$ of a space X as the supremum of the values $\pi\chi(Y)$ where Y ranges over all (Tychonov) continuous images of X . Our main result says that every space X has a π -base whose order is $\leq p\pi\chi(X)$; that is, every point in X is contained in at most $p\pi\chi(X)$ -many members of the π -base. Since $p\pi\chi(X) \leq t(X)$ for compact X , this is a significant generalization of a celebrated result of Shapirovskii.

Let us start by recalling a few definitions and basic facts. A π -base $\mathcal{B}$ of a space X (resp. a local π -base at a point $x \in X$) is a family of non-empty open sets such that every non-empty open set (resp. every neighbourhood of x) includes some member of $\mathcal{B}$. The π -weight $\pi(X)$ of X is the smallest infinite cardinal such that X has a π -base of at most that cardinality. The π -character $\pi\chi(x, X)$ of x in X is the smallest cardinality of a local π -base at $x \in X$, and

$$\pi\chi(X) = \sup\{\pi\chi(x, X) : x \in X\}$$

is the π -character of the space X . Finally, the local tightness at $x \in X$ is the smallest cardinal κ such that if x belongs to the closure $\overline{A}$ of a set A , then there is $B \subset A$ with $|B| \leq \kappa$ and $x \in \overline{B}$; moreover

$$t(X) = \sup\{t(x, X) : x \in X\}$$

is the tightness of the space X .

Shapirovskii proved the following two important results concerning these cardinal functions for compacta: If X is compact, then $\pi\chi(X) \leq t(X)$. Moreover X has a π -base $\mathcal{B}$ of order $\leq t(X)$; i.e. every point of X is contained in at most $t(X)$ -many members of $\mathcal{B}$. (A trivial consequence is that if $t(X)^+$ is a caliber of X —i.e. among $t(X)^+$ -many open sets there are always $t(X)^+$ -many with non-empty intersection—then X has a π -base of cardinality at most $t(X)$.) The first result was proved in [5], and alternative proofs were given in [1] and [3]. The second result first appeared in [6] and then in [8]. A very short and elegant new proof (using a variant of Shapirovskii’s “algebraic” approach to free sequences) was presented in [9].

Arhangel’skii has recently introduced in [2] the concept of a space of *countable projective π -character* and noticed that any compact space of countable tightness

Received by the editors March 28, 2007, and, in revised form, June 17, 2007.

2000 *Mathematics Subject Classification*. Primary 54A25, 54C10, 54D70.

Key words and phrases. Projective π -character, order of a π -base, irreducible map.

This research was supported by OTKA grant no. 61600.

©2008 American Mathematical Society
 Reverts to public domain 28 years from publication

has countable projective π -character. Then he showed that a compact space of countable projective π -character that has ω_1 as a caliber is separable (or equivalently, has a countable π -base), thereby strengthening the above consequence of Shapirovskii's result for countably tight compacta.

In this paper we introduce the general concept of projective π -character and give the following significant generalization of Shapirovskii's full result: Any Tychonov space has a π -base of order at most the projective π -character of the space. Not only is this result stronger for compacta, because it replaces tightness with projective π -character that is smaller, but somewhat surprisingly it extends to all Tychonov spaces.

Let φ be any cardinal function defined on a class $\mathcal{C}$ of topological spaces. We define the projective version $p\varphi$ of φ on $\mathcal{C}$ as follows. For any $X \in \mathcal{C}$ we let $p\varphi(X)$ be the supremum of the values $\varphi(Y)$ where Y ranges over all continuous images of X belonging to $\mathcal{C}$. In particular, we shall consider the case in which $\varphi = \pi\chi$, the π -character defined on the class of Tychonov spaces. It is easy to show then that a Tychonov space X has countable projective π -character in the sense of [2] iff $p\pi\chi(X) \leq \omega$.

Also, as was already mentioned, if X is compact Hausdorff, then we have $p\pi\chi(X) \leq t(X)$. In fact, this follows because $t(Y) \leq t(X)$ for any continuous image of X and, by Shapirovskii's first result above, $\pi\chi(Y) \leq t(Y)$ for every compact Y . But are $p\pi\chi(X)$ and $t(X)$ really different? Arhangel'skii asked more specifically if there is a compactum of countable projective π -character that is not countably tight; see [2], problem 7. Our next example yields such a compactum.

Example 1. Let X be a compactification of ω whose remainder is (homeomorphic to) the ordinal $\omega_1 + 1$. Then $p\pi\chi(X) \leq \omega < t(X)$.

Proof. It is obvious that $t(\omega_1, X) = t(X) = \omega_1$. To see $p\pi\chi(X) \leq \omega$, consider any continuous surjection $f : X \rightarrow Y$. If $f(\omega_1) = p$ is an isolated point in Y , then there is an $\alpha < \omega_1$ such that f is constant on the interval $[\alpha, \omega_1]$; hence Y is countable and compact and so, trivially, $\pi\chi(Y) \leq w(Y) = \omega$.

If, however, p is not isolated, then Y has a countable dense subset S with $p \notin S$. So there is a closed G_δ set F such that $p \in F \subset Y \setminus S$, and again we can find an $\alpha < \omega_1$ such that $f[\alpha, \omega_1] \subset F$. But then $G = Y \setminus F$ is countable and dense open in Y ; moreover $w(G) = \omega$ because every countable and locally compact space is second countable. So we have $\pi\chi(Y) \leq \pi(Y) = w(G) = \omega$. $\square$

We recall from [3] that $\pi sw(X)$ denotes the π -separating weight of a space X , that is, the minimum order of a π -base of X ; see p. 74 of [3].

With this we may now formulate our main result as follows.

Theorem 2. *For any Tychonov space X we have $\pi sw(X) \leq p\pi\chi(X)$. In particular, any Tychonov space of countable projective π -character has a point-countable π -base.*

Our proof of Theorem 2 will go along similar lines as Shapirovskii's proof of the weaker result $\pi sw(X) \leq t(X)$ for compact spaces. The main idea of that was to show that the compactum X admits an irreducible map onto a subspace of the $\Sigma_{t(X)}$ -power of the unit interval. The role of irreducible maps in our proof will be played by a new, more general, type of maps that we shall call π -irreducible. So we shall first define and deal with these maps. (The referee has pointed out to us that [8] is an excellent source concerning Shapirovskii's original method.)

Definition 3. Let f be a continuous map of X onto Y . We say that the map f is π -irreducible if for every proper closed subset $F \subset X$ its image $f[F]$ is not dense in Y .

Clearly, an onto map f is π -irreducible iff the f -image of a non-dense set is non-dense. Also, it is obvious that a closed map is π -irreducible iff it is irreducible; consequently the two concepts coincide for maps between compact Hausdorff spaces.

The following proposition will be used in the proof of Theorem 2 and explains our terminology.

Proposition 4. Let f be a continuous map of X onto Y . Then the following five statements are equivalent:

- (1) f is π -irreducible;
- (2) for every π -base $\mathcal{B}$ of X and for every $B \in \mathcal{B}$ the f -image of its complement, $f[X \setminus B]$, is not dense in Y ;
- (3) there is a π -base $\mathcal{B}$ of X such that for every $B \in \mathcal{B}$ the f -image $f[X \setminus B]$ is not dense in Y ;
- (4) for every π -base $\mathcal{C}$ of Y the family $\{f^{-1}(C) : C \in \mathcal{C}\}$ is a π -base of X ;
- (5) there is a π -base $\mathcal{C}$ of Y such that $\{f^{-1}(C) : C \in \mathcal{C}\}$ is a π -base of X .

Proof. We shall show (3) $\Rightarrow$ (4) and (5) $\Rightarrow$ (1) only because the other three implications of the cycle are trivial.

So, let $\mathcal{B}$ be as in (3) and $\mathcal{C}$ be any π -base of Y . For every non-empty open set U in X choose $B \in \mathcal{B}$ with $B \subset U$. Then there is a $C \in \mathcal{C}$ such that $C \cap f[X \setminus B] = \emptyset$, and hence $f^{-1}(C) \subset B \subset U$.

Now, let $\mathcal{C}$ be as in (5) and F be a proper closed subset of X . Then there is a $C \in \mathcal{C}$ with $F \cap f^{-1}(C) = \emptyset$; consequently we have $f[F] \cap C = \emptyset$ and so $f[F]$ is not dense in Y . $\square$

Corollary 5. If $f : X \rightarrow Y$ is π -irreducible, then $\pi(X) = \pi(Y)$.

Proof. $\pi(X) \leq \pi(Y)$ is immediate from part (4) of Proposition 4. To see $\pi(X) \geq \pi(Y)$ first note that for any non-empty open $U \subset X$ the interior of $f[U]$ in Y is non-empty. So for any π -base $\mathcal{B}$ of X the family $\{\text{Int}_Y(f[B]) : B \in \mathcal{B}\}$ is a π -base of Y . Indeed, this is because if V is non-empty open in Y and $B \in \mathcal{B}$ with $B \subset f^{-1}(V)$, then $f[B] \subset V$. $\square$

We now consider another key ingredient of the proof of our main result: certain specially embedded subspaces of Tychonov cubes. As usual, we shall denote the unit interval $[0, 1]$ by I . The members of the Tychonov cube I^κ will be construed as functions from κ to I . So if $x \in I^\kappa$ and $\alpha < \kappa$, then $x \restriction \alpha$ is the projection of x to the subproduct I^α .

Definition 6. We say that $Y \subset I^\kappa$ is 0-embedded in the Tychonov cube I^κ if

$$\{y \restriction \alpha : y \in Y \text{ and } y(\alpha) = 0\}$$

is dense in the projection $Y \restriction \alpha = \{y \restriction \alpha : y \in Y\}$ for every $\alpha < \kappa$.

We now present two results concerning 0-embedded subspaces of Tychonov cubes which will be crucial in the proof of our main theorem and are also interesting in themselves.

Theorem 7. *Assume that Y is 0-embedded in the Tychonov cube I^κ where κ is a regular cardinal and $y \in Y$ is such that $y(\alpha) > 0$ for all $\alpha < \kappa$. Then $\pi\chi(y, Y) = \kappa$.*

Proof. Of course, only $\pi\chi(y, Y) \geq \kappa$ needs to be proven. To see this, let $\mathcal{U}$ be any family of elementary open sets in I^κ such that $|\mathcal{U}| < \kappa$ and $U \cap Y \neq \emptyset$ for all $U \in \mathcal{U}$. Every elementary open set $U \in \mathcal{U}$ is supported by a finite subset of κ ; hence the regularity of κ implies the existence of an ordinal $\alpha < \kappa$ such that the support of each $U \in \mathcal{U}$ is included in α .

Since Y is 0-embedded in I^κ , this implies that for every $U \in \mathcal{U}$ we may pick a point $y_U \in U \cap Y$ such that $y_U(\alpha) = 0$. But then $y(\alpha) > 0$ clearly implies that the point y is not in the closure of the set $\{y_U : U \in \mathcal{U}\}$; consequently $\mathcal{U}$ cannot be a local π -base at y in Y , completing the proof. $\square$

From Theorem 7 we can immediately obtain the following useful corollary about the projective π -character of 0-embedded subspaces of Tychonov cubes.

Corollary 8. *If Y is 0-embedded in the Tychonov cube I^κ , then for every non-isolated point $y \in Y$ we have*

$$p\pi\chi(Y) \geq |\{\alpha : y(\alpha) > 0\}|,$$

and if $y \in Y$ is isolated, then $\{\alpha : y(\alpha) > 0\}$ is finite.

Our next result shows that every Tychonov space admits a π -irreducible map onto a suitable 0-embedded subspace of a Tychonov cube.

Theorem 9. *Let X be any Tychonov space of π -weight $\pi(X) = \kappa$. Then there is a π -irreducible map f of X onto a 0-embedded subspace Y of the Tychonov cube I^κ .*

Proof. To begin with, let us choose a π -base $\mathcal{B}$ of X with $|\mathcal{B}| = \kappa$ and fix a well-ordering $\prec$ of $\mathcal{B}$ of order-type κ .

We shall define by transfinite induction on $\alpha < \kappa$ the co-ordinate maps $g_\alpha = p_\alpha \circ f : X \rightarrow I$, where $p_\alpha(y) = y(\alpha)$ is the α th co-ordinate projection, and sets $B_\alpha \in \mathcal{B}$. So assume that $\alpha < \kappa$ and for all $\beta < \alpha$ the maps $g_\beta : X \rightarrow I$ and the sets $B_\beta \in \mathcal{B}$ have been defined.

Let $f_\alpha : X \rightarrow I^\alpha$ be the map whose β th co-ordinate map is g_β for all $\beta < \alpha$ and set $Y_\alpha = f_\alpha[X]$. Then, in view of Corollary 5, the map $f_\alpha : X \rightarrow Y_\alpha$ cannot be π -irreducible because $\pi(Y_\alpha) < \kappa = \pi(X)$; hence using part (2) of Proposition 4 there is a member $B \in \mathcal{B}$ for which $f_\alpha[X \setminus B]$ is dense in Y_α . Let B_α be the $\prec$ -first such member of $\mathcal{B}$. We then define $g_\alpha : X \rightarrow I$ as any continuous function that is identically 0 on $X \setminus B_\alpha$ and takes the value 1 at some point in B_α . As was intended, with the induction completed we let $f : X \rightarrow I^\kappa$ be the unique map having the g_α for $\alpha < \kappa$ as its co-ordinate functions and we also set $Y = f[X]$.

Note first that if $\beta < \alpha$, then $B_\beta \prec B_\alpha$. Indeed, since we have $Y_\beta = Y_\alpha \upharpoonright \beta$, the density of $f_\alpha[X \setminus B_\alpha]$ in Y_α implies that $f_\beta[X \setminus B_\alpha]$ is dense in Y_β ; hence $B_\alpha \prec B_\beta$ would contradict the choice of B_β . Moreover, by our construction, $f_{\beta+1}[X \setminus B_\beta]$ is not dense in $Y_{\beta+1}$ and consequently $f_\alpha[X \setminus B_\beta]$ is not dense in Y_α , which implies $B_\alpha \neq B_\beta$.

Since $\mathcal{B}$ is of order type κ under $\prec$, it follows from this that for every $B \in \mathcal{B}$ there is an $\alpha < \kappa$ with $B \prec B_\alpha$. But then, by the choice of B_α , we have that $f_\alpha[X \setminus B]$ is not dense in $Y_\alpha = Y \upharpoonright \alpha$ and hence $f[X \setminus B]$ cannot be dense in Y . Using part (3) of Proposition 4 implies that f is indeed a π -irreducible map of X onto Y .

Finally, by our construction, for every $\alpha < \kappa$ the image $f_\alpha[X \setminus B_\alpha]$ is dense in $Y_\alpha = Y \restriction \alpha$; moreover we have

$$f_\alpha[X \setminus B_\alpha] \subset \{y \restriction \alpha : y \in Y \text{ and } y(\alpha) = 0\},$$

and consequently Y is indeed 0-embedded in I^κ . $\square$

Let us now recall that the κ th Σ_λ -power of I , denoted by $\Sigma_\lambda(I, \kappa)$, is the subspace of I^κ consisting of all points whose support is of size at most λ . The support of a point $y \in I^\kappa$ is the set $\{\alpha < \kappa : y(\alpha) > 0\}$. Thus, from Theorem 9 and from Corollary 8—moreover from the trivial fact that $p\pi\chi(Y) \leq p\pi\chi(X)$ if Y is any continuous image of X —we immediately obtain the following result.

Corollary 10. *If X is a Tychonov space such that $\pi(X) = \kappa$ and $p\pi\chi(X) = \lambda$, then some π -irreducible image Y of X embeds into $\Sigma_\lambda(I, \kappa)$.*

This corollary is clearly a strengthening of the following result of Shapirovskii from [6] (see also 3.22 of [3]): If X is compact Hausdorff, then some irreducible image of X embeds into a $\Sigma_{t(X)}$ -power of I .

The proof of our main theorem, Theorem 2, can now be easily established by recalling the following result of Shapirovskii from [6] (see also [3], 3.24).

Theorem (Shapirovskii). *If the space Y embeds into a Σ_λ -power of I , then $\pi sw(Y) \leq \lambda$.*

Proof of Theorem 2. Now, to prove Theorem 2, consider any non-discrete Tychonov space X . By Corollary 10 then X has a π -irreducible image Y that embeds into a Σ_λ -power of I , where $\lambda = p\pi\chi(X)$. By the previous theorem of Shapirovskii the space Y has a π -base $\mathcal{C}$ of order at most λ . But by part (4) of Proposition 4, $\{f^{-1}(C) : C \in \mathcal{C}\}$ is a π -base of X that clearly has the same order as $\mathcal{C}$. $\square$

The following result is then an immediate consequence of Theorem 2.

Corollary 11. *Let X be any Tychonov space and $\kappa > p\pi\chi(X)$ be a cardinal such that κ is a caliber of X . Then $\pi(X) < \kappa$.*

Since $t(X) \geq p\pi\chi(X)$ for a compact Hausdorff space X , this corollary implies Shapirovskii's theorem saying that if $t(X)^+$ is a caliber of such a space X , then $\pi(X) \leq t(X)$. Moreover, it also extends from compacta to all Tychonov spaces Arhangel'skii's result from [2] saying that spaces of countable projective π -character and having ω_1 as a caliber are separable.

Let us conclude by pointing out that neither Theorem 2 nor Corollary 11 remain valid if the projective π -character $p\pi\chi$ is replaced by the simple π -character $\pi\chi$ in them. In fact, it has recently been shown in [4] that there are even first countable spaces whose π -separating weight is as large as you wish. Moreover, in the same paper it was also shown that it is consistent to have first countable spaces with caliber ω_1 which have uncountable π -weight (or equivalently, density). However, since first countability implies countable tightness, none of these examples are (or could be) compact, so the following intriguing questions remain open.

Problem 12. Let X be a compact Hausdorff space of countable π -character. Does X have a point-countable π -base? If, in addition, ω_1 is a caliber of X , is X then separable?

REFERENCES

- [1] A. V. Arhangel'skii, *Structure and classification of topological spaces and cardinal invariants*, Russian Math. Surveys 33 (1978), pp. 29–84. MR526012 (80i:54005)
- [2] A. V. Arhangel'skii, *Precalibers, monolithic spaces, first countability, and homogeneity in the class of compact spaces*, preprint.
- [3] I. Juhász, *Cardinal functions – 10 years later*, Math. Center Tract no. 123, Amsterdam, 1980. MR576927 (82a:54002)
- [4] I. Juhász, L. Soukup, and Z. Szentmiklóssy, *First countable spaces without point-countable π -bases*, Fund. Math., 196 (2007), pp. 139–149. MR2342624
- [5] B. E. Shapirovskii, *On the tightness, π -weight and related notions*, Uch. Zap. Latv. Univ. 257 (1976), pp. 88–99.
- [6] B. E. Shapirovskii, *Special types of embeddings in Tychonoff cubes. Subspaces of Σ -products and cardinal invariants*, in: Topology, Coll. Math. Soc. J. Bolyai 23 (North-Holland, Amsterdam, 1980), pp. 1055–1086. MR588855 (82d:54010)
- [7] B. E. Shapirovskii, *Mappings on Tikhonov cubes*, Russian Math. Surveys 35 (1980), pp. 145–156 (1981). MR580628 (82d:54018)
- [8] B. E. Shapirovskii, *Cardinal invariants in compact Hausdorff spaces*, Amer. Math. Soc. Transl. 134 (1987), pp. 93–118.
- [9] S. Todorčević, *Free sequences*, Top. Appl. 35 (1990), pp. 235–238. MR1058803 (91f:54003)

ALFRÉD RÉNYI INSTITUTE OF MATHEMATICS, HUNGARIAN ACADEMY OF SCIENCES, POB 127,
BUDAPEST, H-1364 HUNGARY
E-mail address: juhász@renyi.hu

DEPARTMENT OF ANALYSIS, EÖTVÖS LORÁNT UNIVERSITY, PÁZMÁNY PÉTER SÉTÁNY 1/A, 1117
BUDAPEST, HUNGARY
E-mail address: zoli@renyi.hu