ON A PROBLEM OF BERNARD CHEVREAU CONCERNING THE ρ -CONTRACTIONS

P. GĂVRUȚA

(Communicated by Michael T. Lacey)

ABSTRACT. We prove new results for the operators of class C_ρ ($\rho > 0$) on Hilbert spaces defined by B. Sz.-Nagy and C. Foiaş. The main result is an answer to a problem posed in 2006 by B. Chevreau: Let $p \geq 2$ be a natural number and $T \in L(\mathcal{H})$; if there exists $\rho_0 > 0$ such that $T^p \in C_{\rho_0}$, then necessarily is $T \in \bigcup_{\rho>0} C_\rho$?

1. INTRODUCTION

We denote by $\mathcal{H}$ a Hilbert space over field $\mathbb{C}$ of all complex numbers. Let $L(\mathcal{H})$ denote the algebra of all bounded operators on $\mathcal{H}$. $I_{\mathcal{H}}$ is the identity operator on $\mathcal{H}$. For $\rho > 0$, we denote by $C_\rho(\mathcal{H})$ the class of operators which admits a unitary ρ -dilation: an operator $T \in L(\mathcal{H})$ admits a unitary ρ -dilation if there exists a Hilbert space $\mathcal{K} \supset \mathcal{H}$ and $U \in L(\mathcal{K})$ unitary such that

$$T^n = \rho P U^n|_{\mathcal{H}}, \quad n \geq 1,$$

where P is the orthogonal projection of $\mathcal{K}$ on $\mathcal{H}$.

If $T \in C_\rho(\mathcal{H})$, then T is also called a ρ -contraction. The classes C_ρ were introduced by B. Sz.-Nagy and C. Foiaş in [15] (see also [16]). The class $C_1(\mathcal{H})$ is precisely the class of all contractions on $\mathcal{H}$. The class $C_2(\mathcal{H})$ is precisely the class of all operators $T \in L(\mathcal{H})$ which have numerical radius equal to at most one ([1]).

We remember that for an operator $T \in L(\mathcal{H})$, the numerical radius is defined by

$$w(T) = \sup\{|(Th, h)|, h \in \mathcal{H}, \|h\| = 1\}.$$

It is known ([16]) that $C_\rho(\mathcal{H}) \subset C_{\rho'}(\mathcal{H})$ if $0 < \rho < \rho' < \infty$ and $C_\rho(\mathcal{H}) \neq C_{\rho'}(\mathcal{H})$ if $\rho \neq \rho'$ and $\dim(\mathcal{H}) \geq 2$.

We denote $C_\infty(\mathcal{H}) = \bigcup_{\rho>0} C_\rho(\mathcal{H})$.

In February 2006, Professor Bernard Chevreau [3] posed the following:

Problem. Let $p \geq 2$ be a natural number and $T \in L(\mathcal{H})$. If there exists $\rho_0 > 0$ such that $T^p \in C_{\rho_0}$, then necessarily is $T \in C_\infty(\mathcal{H})$?

The answer is important for recent development in the theory of operators of class C_ρ ([2], [4], [5]).

Received by the editors June 7, 2007.

2000 *Mathematics Subject Classification.* Primary 47A20; Secondary 47B99.

Key words and phrases. Operator, dilation, ρ -contraction.

©2008 American Mathematical Society
 Reverts to public domain 28 years from publication

We give an answer to this question. In fact, we prove that there exists $T \in L(\mathbb{C}^p)$ such that $T^p = I_{\mathbb{C}^p}$ and $T \notin C_\infty(\mathbb{C}^p)$.

We use the following result given by G. Eckstein [6]:

Theorem 1.1. *Let $0 < \rho < \infty$ and $T \in C_\rho(\mathcal{H})$. Then, for any $h \in \mathcal{H}$, the sequence $\{\|T^n h\|\}_{n=0}^\infty$ converges.*

A short proof of this theorem was given in [11]. An another proof was given in [9].

2. THE RESULTS

We proved before the following result:

Theorem 2.1. *Let $p \geq 2$ be a natural number and $0 < \rho < \infty$. If $T \in C_\rho(\mathcal{H})$ and T^p is a unitary operator, then T is unitary.*

Proof. We denote $V := T^p$. Let $h \in \mathcal{H}$ and n a natural number. We have

$$\|T^{np}h\| = \|V^n h\| = \|h\|$$

and

$$\|T^{np+1}h\| = \|T^{np}Th\| = \|V^n Th\| = \|Th\|.$$

Since $T \in C_\rho(\mathcal{H})$, it follows by Theorem 1.1 that the sequence $\{\|T^n h\|\}_{n=0}^\infty$ converges. It follows that the sequences

$$\{\|T^{np}h\|\}_{n=0}^\infty \quad \text{and} \quad \{\|T^{np+1}h\|\}_{n=0}^\infty$$

converge to the same limit. It results in $\|h\| = \|Th\|$. Hence T is an isometry.

We also have $V^* = T^{*p}$. But $T^* \in C_\rho(\mathcal{H})$ since we can write

$$T^{*n}h = \rho P U^{*n}h, \quad n \geq 1, \quad h \in \mathcal{H}.$$

By the previous argument it follows that T^* is also an isometry. Hence T is a unitary operator. $\square$

We give a generalization of Theorem 2.1, but in the proof we use another technique.

Let $T_1, T_2 \in L(\mathcal{H})$. We say that T_1 and T_2 are *double commuting* if T_1 commutes with T_2 and with T_2^* .

We use the following result given by B. Fuglede [7]:

Theorem 2.2. *Let T_1, T_2 be bounded operators on $\mathcal{H}$, with T_2 being normal. If T_1 and T_2 commute, then T_1 and T_2 are double commuting.*

Our result is the following:

Theorem 2.3. *Let $p \geq 2$ be a natural number and $0 < \rho < \infty$. If $T \in C_\rho(\mathcal{H})$ is such that $N := T^p$ is normal and norm increasing (i.e. $N^*N \geq I_{\mathcal{H}}$), then T is unitary.*

Proof. Since N is norm increasing and normal, we have

$$\|Nh\| \geq \|h\| \quad \text{and} \quad \|N^*h\| = \|Nh\| \geq \|h\|, \quad h \in \mathcal{H}.$$

It follows that N is invertible and $\|N^{-1}h\| \leq \|h\|$, $h \in \mathcal{H}$.

We have $TN = T^{p+1}$ and $NT = T^{p+1}$; hence $N^{-1}T = TN^{-1}$. We obtain

$$T(T^{p-1}N^{-1}) = I_{\mathcal{H}} \quad \text{and} \quad (T^{p-1}N^{-1})T = I_{\mathcal{H}}.$$

It follows that T is invertible and $T^{-1} = T^{p-1}N^{-1}$. Since $T \in C_\rho(\mathcal{H})$, it follows by [16] that $T^{p-1} \in C_\rho(\mathcal{H})$.

Since T^{p-1} and N^{-1} commute and N^{-1} is normal, by Theorem 2.2 it results that T^{p-1} and N^{-1} are double commuting. We use the following known result (see [8], [10], [14]):

If $T_1 \in C_{\rho_1}(\mathcal{H})$, $T_2 \in C_{\rho_2}(\mathcal{H})$ are double commuting, then $T_1T_2 \in C_{\rho_1\rho_2}(\mathcal{H})$ ($0 < \rho_1, \rho_2 < \infty$).

We apply this result for $T_1 = T^{p-1} \in C_\rho(\mathcal{H})$ and $T_2 = N^{-1} \in C_1(\mathcal{H})$. It follows that $T^{-1} = T^{p-1}N^{-1} \in C_\rho(\mathcal{H})$. Since $T, T^{-1} \in C_\rho(\mathcal{H})$, by a well-known result of J. G. Stampfli [13] (see also [12]), it follows that T is unitary. $\square$

We now prove the main result of this paper.

Theorem 2.4. *Let $p \geq 2$ be a natural number. Then there exists $T \in L(\mathbb{C}^p)$ such that $T^p = I_{\mathbb{C}^p}$ and $T \notin C_\infty(\mathbb{C}^p)$.*

Proof. We consider T the operator on $\mathbb{C}^p$ which has the matrix

$$[T] = \begin{pmatrix} 0 & 0 & 0 & \dots & 0 & a_1 \\ a_2 & 0 & 0 & \dots & 0 & 0 \\ 0 & a_3 & 0 & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & a_p & 0 \end{pmatrix}$$

where $a_1 \cdot a_2 \cdot \dots \cdot a_p = 1$ and there exists $j \in \{1, 2, \dots, p\}$ such that $a_j \neq 1$. We have

$$\det[T] = (-1)^{p+1} a_1 \cdot a_2 \cdot \dots \cdot a_p \neq 0;$$

hence T is diagonalizable.

We denote by $\lambda_1, \lambda_2, \dots, \lambda_p$ the roots of the equation

$$(2.1) \quad \det([T] - \lambda I_p) = 0,$$

where I_p is the identity matrix of order p . Then we can write

$$[T] = S^{-1} \Lambda S,$$

where

$$[\Lambda] = \text{diag}(\lambda_1, \lambda_2, \dots, \lambda_p)$$

and S is an invertible matrix.

The equation (2.1) is equivalent to

$$\begin{vmatrix} -\lambda & 0 & 0 & 0 & \dots & 0 & a_1 \\ a_2 & -\lambda & 0 & 0 & \dots & 0 & 0 \\ 0 & a_3 & -\lambda & 0 & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & \dots & a_p & -\lambda \end{vmatrix} = 0;$$

that is,

$$(-\lambda)^p + (-1)^{p+1} a_1 a_2 \dots a_p = 0$$

or

$$\lambda^p = a_1 \dots a_p.$$

We have

$$[T]^p = S^{-1} \Lambda^p S = S^{-1} I_p S = I_p.$$

Hence $T^p = I_{\mathbb{C}^p}$.

But $T \notin C_\infty(\mathbb{C}^p)$. In a contrary case, by Theorem 2.1, it follows that T is unitary. $\square$

REFERENCES

1. C. A. Berger, *A strange dilation theorem*, Notices Amer. Math. Soc. **12** (1965), 590.
2. G. Cassier, N. Suciu, *Mapping theorems and Harnack ordering for ρ -contractions*, Indiana Univ. Math. J. **55** (2006), 483–523. MR2225443
3. B. Chevreau, *Talk given at West University of Timișoara*, February 2006.
4. B. Chevreau, A. Crăciunescu, *On factorization properties for ρ -contractions*, Talk given in The 21st Intern. Conference on Operator Theory, Timișoara, 2006.
5. M. Dritschel, S. McCullough, H. J. Woerdeman, *Model theory for ρ -contractions, $\rho \leq 2$* , J. Operator Theory **41** (1999), 321–350. MR1681577 (2000b:47025)
6. G. Eckstein, *Sur les opérateurs de classe C_ρ* , Acta Sci. Math. (Szeged) **33** (1972), 349–352. MR0333768 (48:12092)
7. B. Fuglede, *A commutativity theorem for normal operators*, Proc. Nat. Acad. Sci. **36** (1950), 35–40. MR0032944 (11:371c)
8. D. Gășpar, *Sur certaines familles d'opérateurs dans l'espace de Hilbert*, Rev. Roum. Math. Pures Appl. **13** (1968), 779–788. MR0239443 (39:800)
9. P. Găvruta, D. Păunescu, *Moment problems for operators*, Anal. Univ. Timișoara, Ser. Sci. Mat. **27** (1989), 49–58. MR1140522 (92m:47029)
10. J. Holbrook, *Multiplicative properties of the numerical radius in operator theory*, J. Reine Angew. Math. **237** (1969), 166–174. MR0247515 (40:780)
11. W. Mlak, *On convergence properties of operators of class C_ρ* , Acta Sci. Math. (Szeged) **33** (1972), 353–354. MR0333769 (48:12093)
12. A. Rácz, *Unitary skew-dilations*, Stud. Cerc. Mat. **26** (1974), 545–621 (Roumanian, English summary). MR0355640 (50:8114)
13. J. G. Stampfli, *A local spectral theory for operators*, J. Funct. Analysis **4** (1969), 1–10. MR0243376 (39:4698)
14. B. Sz.-Nagy, *Product of operators of classes C_ρ* , Rev. Roum. Math. Pures Appl. **13** (1968), 897–899. MR0239455 (39:812)
15. B. Sz.-Nagy, C. Foiaș, *On certain classes of power-bounded operators in Hilbert space*, Acta Sci. Math. (Szeged) **27** (1966), 17–25. MR0198254 (33:6413)
16. B. Sz.-Nagy, C. Foiaș, *Analyse harmonique des opérateurs de l'espace Hilbert*, Masson et Cie, Paris, 1967. MR0225183 (37:778)

DEPARTMENT OF MATHEMATICS, UNIVERSITY “POLITEHNICA” OF TIMIȘOARA, PIAȚA VICTORIEI
No. 2, 300006 TIMIȘOARA, ROMANIA
E-mail address: pgavruta@yahoo.com