CESÀRO FUNCTION SPACES FAIL THE FIXED POINT PROPERTY

SERGEI V. ASTASHKIN AND LECH MALIGRANDA

(Communicated by N. Tomczak-Jaegermann)

ABSTRACT. The Cesàro sequence spaces ces_p , $1 < p < \infty$, are reflexive but they have the fixed point property. In this paper we prove that in contrast to these sequence spaces the corresponding Cesàro function spaces Ces_p on both $[0, 1]$ and $[0, \infty)$ for $1 < p < \infty$ are not reflexive and they fail to have the fixed point property.

1. INTRODUCTION

Let $1 \leq p \leq \infty$. The *Cesàro sequence space* ces_p is defined as the set of all real sequences $x = \{x_k\}$ such that

$$\|x\|_{c(p)} = \left(\sum_{n=1}^{\infty} \left(\frac{1}{n} \sum_{k=1}^n |x_k| \right)^p \right)^{1/p} < \infty \text{ when } 1 \leq p < \infty,$$

and

$$\|x\|_{c(\infty)} = \sup_{n \in \mathbf{N}} \frac{1}{n} \sum_{k=1}^n |x_k| < \infty \text{ when } p = \infty.$$

The *Cesàro function spaces* $Ces_p = Ces_p(I)$ are the classes of Lebesgue measurable real functions f on $I = [0, 1]$ or $I = [0, \infty)$ such that the corresponding norms are finite, where

$$\|f\|_{C(p)} = \left(\int_I \left(\frac{1}{x} \int_0^x |f(t)| dt \right)^p dx \right)^{1/p} \text{ for } 1 \leq p < \infty,$$

and

$$\|f\|_{C(\infty)} = \sup_{x \in I, x > 0} \frac{1}{x} \int_0^x |f(t)| dt < \infty \text{ for } p = \infty.$$

The Cesàro sequence spaces ces_p were investigated in the seventies by Shiue, Leibowitz and Jagers. In particular, they proved that $ces_1 = \{0\}$, ces_p are separable reflexive Banach spaces for $1 < p < \infty$ and the l^p spaces are continuously embedded

Received by the editors October 18, 2007.

2000 *Mathematics Subject Classification*. Primary 46E30, 46B20, 46B42.

Key words and phrases. Cesàro sequence spaces, Cesàro function spaces, fixed point property, asymptotically isometric copy of l^1 , B -convex spaces.

This research was supported by a grant from the Royal Swedish Academy of Sciences for cooperation between Sweden and the former Soviet Union (project 35440). The results were presented by the second author at *The 8th International Conference on Fixed Point Theory and Its Applications*, 16-22 July 2007, Chiang Mai, Thailand.

©2008 American Mathematical Society
 Reverts to public domain 28 years from publication

into ces_p for $1 < p \leq \infty$ with strict embeddings. Also if $1 < p < q \leq \infty$, then $ces_p \subset ces_q$ with continuous strict embeddings. Bennett [3] proved that ces_p for $1 < p < \infty$ are not isomorphic to any l^q space with $1 \leq q \leq \infty$ (see also [15] for another proof). Moreover, Maligranda-Petrot-Suantai [15] proved recently that Cesàro sequence spaces ces_p for $1 < p < \infty$ are not uniformly nonsquare; that is, there are sequences $\{x_n\}$ and $\{y_n\}$ on the unit sphere such that $\lim_{n \rightarrow \infty} \min(\|x_n + y_n\|_{c(p)}, \|x_n - y_n\|_{c(p)}) = 2$. They even proved that these spaces are not B -convex. We refer here to [3], [15] and the references given there.

Several geometric properties of the Cesàro sequence spaces ces_p were studied in recent years by many mathematicians, and in 1999-2000 it was also proved by Cui-Hudzik [5], Cui-Hudzik-Li [6] and Cui-Meng-Pluciennik [7] that Cesàro sequence spaces ces_p for $1 < p < \infty$ have the fixed point property (cf. also [4, Part 9]).

Cesàro function spaces $Ces_p[0, \infty)$ for $1 \leq p \leq \infty$ were considered by Shiue [16], Hassard-Hussein [12] and Sy-Zhang-Lee [17]. They proved that $Ces_1[0, \infty) = \{0\}$ and $Ces_p[0, \infty)$ for $1 < p < \infty$ are separable Banach spaces and that $Ces_\infty[0, \infty)$ is a nonseparable Banach space. The space $Ces_\infty[0, 1]$ is known as the Korenblyum-Krein-Levin space already introduced in 1948.

By the Hardy inequality the $L^p(I)$ spaces are continuously embedded into $Ces_p(I)$ for $1 < p \leq \infty$ with strict embedding, where $I = [0, 1]$ or $I = [0, \infty)$ (cf. [11, Theorem 327] and [13, Theorem 2]). Also if $1 < p < q \leq \infty$, then $Ces_q[0, 1] \subset Ces_p[0, 1]$ with continuous strict embedding. Moreover, $Ces_1[0, 1]$ is a weighted $L_w^1[0, 1]$ space with the weight $w(t) = \ln \frac{1}{t}$ for $0 < t \leq 1$. In fact,

$$(1) \quad \int_0^1 \left(\frac{1}{x} \int_0^x |f(t)| dt \right) dx = \int_0^1 \left(\int_t^1 \frac{1}{x} dx \right) |f(t)| dt = \int_0^1 |f(t)| \ln \frac{1}{t} dt.$$

We will show that, in contrast to Cesàro sequence spaces, the Cesàro function spaces $Ces_p(I)$ on both $I = [0, 1]$ and $I = [0, \infty)$ for $1 < p < \infty$ are not reflexive and that they do not have the fixed point property.

A Banach space X has the *fixed point property* (FPP) [resp. *weak fixed point property* (WFPP)] if every nonexpansive mapping of every closed bounded convex [resp. nonempty weakly compact convex] subset K of X into itself has a fixed point. Recall that T is said to be *nonexpansive* if $\|Tx - Ty\| \leq \|x - y\|$ for all $x, y \in K$.

The spaces c_0 and l^1 both fail to have the FPP with their classical norms, but they have the WFPP. The space $L^1[0, 1]$ does not have the WFPP, as was proved by Alspach [1].

Our proof that the Cesàro function spaces $Ces_p(I)$ on $I = [0, 1]$ with $1 \leq p \leq \infty$ and on $I = [0, \infty)$ with $1 < p \leq \infty$ fail to have the fixed point property will be carried out by showing that these spaces contain an asymptotically isometric copy of l^1 .

A Banach space X contains an *asymptotically isometric copy* of l^1 if there exists a null sequence (ε_n) in $(0, 1)$ and a sequence (x_n) in X such that

$$\sum_{n=1}^{\infty} (1 - \varepsilon_n) |\alpha_n| \leq \left\| \sum_{n=1}^{\infty} \alpha_n x_n \right\|_X \leq \sum_{n=1}^{\infty} |\alpha_n|$$

for all $(\alpha_n) \in l^1$ of scalars. This notion was introduced by Dowling and Lennard in [9], where they proved that such spaces fail to have the FPP.

2. MAIN RESULTS

Cesàro sequence spaces ces_p , $1 < p < \infty$, are reflexive but not B -convex and they have the fixed point property. In contrast to these sequence spaces the corresponding Cesàro function spaces $Ces_p(I)$ on both $I = [0, 1]$ and $I = [0, \infty)$ for $1 < p < \infty$ are not reflexive and they do not have the fixed point property. Our main result reads:

Theorem 1. *Let $1 \leq p \leq \infty$. The Cesàro function space $Ces_p[0, 1]$ contains an asymptotically isometric copy of l^1 ; that is, there exist a sequence $\{\varepsilon_n\} \subset (0, 1)$, $\varepsilon_n \rightarrow 0$ as $n \rightarrow \infty$ and a sequence of functions $\{f_n\} \subset Ces_p[0, 1]$ such that for arbitrary $\{\alpha_n\} \in l^1$ we have*

$$(2) \quad \sum_{n=1}^{\infty} (1 - \varepsilon_n) |\alpha_n| \leq \left\| \sum_{n=1}^{\infty} \alpha_n f_n \right\|_{C(p)} \leq \sum_{n=1}^{\infty} |\alpha_n|.$$

Before the proof of this theorem we prove the following auxiliary result.

Lemma 1. *Let $0 < a < b < 1$, $f \in Ces_p[0, 1]$ and $\text{supp } f := \{t \in [0, 1] : f(t) \neq 0\} \subset [a, b]$. Then*

$$(3) \quad (b^{1-p} - 1)^{1/p} \|f\|_1 \leq (p-1)^{1/p} \|f\|_{C(p)} \leq (a^{1-p} - 1)^{1/p} \|f\|_1,$$

for $1 < p < \infty$ and

$$(4) \quad \ln \frac{1}{b} \|f\|_1 \leq \|f\|_{C(1)} \leq \ln \frac{1}{a} \|f\|_1, \quad \frac{1}{b} \|f\|_1 \leq \|f\|_{C(\infty)} \leq \frac{1}{a} \|f\|_1,$$

where $\|f\|_1 = \int_0^1 |f(t)| dt$.

Proof. It is obvious that for any $0 < x \leq 1$ we have

$$\frac{1}{x} \|f\|_1 \chi_{[b, 1]}(x) \leq F_f(x) := \frac{1}{x} \int_0^x |f(t)| dt \leq \frac{1}{x} \|f\|_1 \chi_{[a, 1]}(x).$$

Since, for every $c \in (0, 1)$, $\int_c^1 t^{-p} dt = \frac{c^{1-p}-1}{p-1}$ and $\int_c^1 t^{-1} dt = \ln \frac{1}{c}$ we obtain (3) and (4) for $p = 1$. In the case of $p = \infty$ we see that

$$\frac{1}{b} \|f\|_1 \leq \|F_f\|_{L^\infty[0, 1]} \leq \frac{1}{a} \|f\|_1,$$

and the lemma is proved. $\square$

Proof of Theorem 1. For $1 < p < \infty$ we set

$$g_n = \chi_{[a_n, a_{n+1}]}, \text{ with } a_n = 2^{1/(1-p)} \left(1 - \frac{1}{2^n}\right), \quad n = 1, 2, \dots$$

Since $\|g_n\|_1 = a_{n+1} - a_n = 2^{1/(1-p)} \cdot 2^{-n-1}$ and $a_n^{1-p} - 1 = \frac{2}{(1-2^{-n})^{p-1}} - 1$, then, by Lemma 1 (see the second estimate in (3)), this yields that

$$2^{-1/(1-p)} 2^{n+1} (p-1)^{1/p} \|g_n\|_{C(p)} \leq \left(\frac{2}{(1-2^{-n})^{p-1}} - 1 \right)^{1/p}.$$

Let $f_n = g_n / \|g_n\|_{C(p)}$ and $\alpha_n \in \mathbb{R}$ for $n = 1, 2, \dots$. Since $\text{supp } g_n \subset [2^{1(1-p)-1}, 2^{1/(1-p)})$ for every $n \in \mathbb{N}$ it follows from Lemma 1 (see the first estimate in (3))

that

$$\begin{aligned} \left\| \sum_{n=1}^{\infty} \alpha_n f_n \right\|_{C(p)} &\geq \frac{\left\| \sum_{n=1}^{\infty} \alpha_n f_n \right\|_1}{(p-1)^{1/p}} \\ &= \sum_{n=1}^{\infty} \frac{|\alpha_n| 2^{1/(1-p)}}{2^{n+1} (p-1)^{1/p} \|g_n\|_{C(p)}} \\ &\geq \sum_{n=1}^{\infty} \left(\frac{2}{(1-2^{-n})^{p-1}} - 1 \right)^{-1/p} |\alpha_n|. \end{aligned}$$

Denote

$$\varepsilon_n = 1 - \left(\frac{2}{(1-2^{-n})^{p-1}} - 1 \right)^{-1/p}.$$

Then $\{\varepsilon_n\} \subset (0, 1)$ and $\varepsilon_n \rightarrow 0$ as $n \rightarrow \infty$. This means that the left-hand side of (2) is proved. The right-hand side of (2) is obvious since $\|f_n\|_{C(p)} = 1$.

In the case $p = \infty$ we take $f_n = g_n / \|g_n\|_{C(\infty)}$, where $g_n = \chi_{[a_n, a_{n+1})}$ with $a_n = 1 - 2^{-n}$, $n = 1, 2, \dots$. Then $\|g_n\|_1 = 2^{-n-1}$ and, by Lemma 1 (see the second estimate in (4)) $\|g_n\|_{C(\infty)} \leq \frac{1}{1-2^{-n}} 2^{-n-1}$ or $2^{n+1} \|g_n\|_{C(\infty)} \leq \frac{1}{1-2^{-n}}$. Since $\text{supp } g_n \subset [1/2, 1)$, for every $n \in \mathbb{N}$, it follows from Lemma 1 (see the first estimate in (4)) that:

$$\begin{aligned} \left\| \sum_{n=1}^{\infty} \alpha_n f_n \right\|_{C(\infty)} &\geq \left\| \sum_{n=1}^{\infty} \alpha_n f_n \right\|_1 \\ &= \sum_{n=1}^{\infty} \frac{|\alpha_n|}{2^{n+1} \|g_n\|_{C(\infty)}} \geq \sum_{n=1}^{\infty} (1 - 2^{-n}) |\alpha_n|, \end{aligned}$$

and $\varepsilon_n = 2^{-n}$ is a required sequence.

In the case $p = 1$ we take $g_n = \chi_{[a_n, a_{n+1})}$, where $a_n = \frac{1}{e}(1 - 2^{-n})$, $n = 1, 2, \dots$ and argue in a similar way. The proof is complete. $\square$

The analogous result holds for Cesàro function spaces on $[0, \infty)$.

Theorem 2. *Let $1 < p \leq \infty$. The Cesàro function space $Ces_p[0, \infty)$ contains an asymptotically isometric copy of l^1 .*

Proof. We consider only the case $1 < p < \infty$ (the case $p = \infty$ can be proved similarly as in Theorem 1). We take $g_n = \chi_{[a_n, a_{n+1})}$ with $a_n = 1 - 2^{-n}$, $n = 1, 2, \dots$ and continue the proof as in Theorem 1, observing that the estimate corresponding to (3) for $Ces_p[0, \infty)$ will be (for $0 < a < b < \infty$)

$$b^{1/p-1} \|f\|_1 \leq (p-1)^{1/p} \|f\|_{C(p)} \leq a^{1/p-1} \|f\|_1, \text{ with } \|f\|_1 = \int_0^\infty |f(t)| dt,$$

and then for $f_n = g_n / \|g_n\|_{C(p)}$ we have that

$$\begin{aligned} \left\| \sum_{n=1}^{\infty} \alpha_n f_n \right\|_{C(p)} &\geq \frac{\left\| \sum_{n=1}^{\infty} \alpha_n f_n \right\|_1}{(p-1)^{1/p}} \\ &= \sum_{n=1}^{\infty} \frac{|\alpha_n|}{2^{n+1}(p-1)^{1/p} \|g_n\|_{C(p)}} \geq \sum_{n=1}^{\infty} a_n^{1-1/p} |\alpha_n| \\ &= \sum_{n=1}^{\infty} (1-2^{-n})^{1-1/p} |\alpha_n| = \sum_{n=1}^{\infty} (1-\varepsilon_n) |\alpha_n|, \end{aligned}$$

and $\varepsilon_n = 1 - (1 - 2^{-n})^{1-1/p}$ is a required sequence. The proof is complete. $\square$

Remark 1. It is obvious from Theorem 1 and Theorem 2 that the Cesàro function spaces $Ces_p(I)$ for $1 < p \leq \infty$ are not reflexive.

Dowling-Lennard [9] proved that if a Banach space X contains an asymptotically isometric copy of l^1 , then there exists a nonexpansive mapping defined on a closed bounded convex subset of X without a fixed point, i.e., that X fails to have the fixed point property (see also [10, Theorem 2.3 and Corollary 2.11]). By the Dilworth-Girardi-Hagler result [8, Theorem 2] the dual space X^* does not have the fixed point property since they proved there that X contains an asymptotically isometric copy of l^1 if and only if the dual space X^* contains an isometric copy of $L^1[0, 1]$. Combining these results with our Theorem 1 and Theorem 2 we obtain immediately our main result on the fixed point property of Cesàro function spaces and their dual spaces.

Theorem 3. *If either $1 \leq p \leq \infty$ and $I = [0, 1]$ or $1 < p \leq \infty$ and $I = [0, \infty)$, then the Cesàro function spaces $Ces_p(I)$ and their dual spaces $Ces_p(I)^*$ fail to have the fixed point property.*

Theorem 3 gives information about the fixed point property, and therefore it is natural to ask what one can say about the weak fixed point property.

Note that the space $Ces_1[0, 1]$ is isometric to $L^1[0, 1]$ by the equality (1), and by the Alspach result [1] $L^1[0, 1]$ fails to have WFPP; therefore $Ces_1[0, 1]$ also fails to have WFPP.

By combining Theorem 1, Theorem 2 and the Dilworth-Girardi-Hagler result [8, Corollary 13] we have the following result:

Proposition 1. *The dual spaces to the Cesàro function spaces $Ces_p(I)^*$ do not have the weak fixed point property.*

Proposition 2. *The Cesàro function spaces $Ces_p(I)$ for $1 < p < \infty$ are not isomorphic to any $L^q(I)$ space for $1 \leq q \leq \infty$. In particular, they are not isomorphic to $L^1(I)$.*

Of course, $Ces_p(I)$ for $1 < p < \infty$, as nonreflexive and separable spaces, cannot be isomorphic to any $L^q(I)$ with $1 < q < \infty$ or $q = \infty$. The statement for $q = 1$ will be proved in the forthcoming paper [2].

REFERENCES

- [1] D. E. Alspach, *A fixed point free nonexpansive map*, Proc. Amer. Math. Soc. **82**(1981), 423–424. MR612733 (82j:47070)
- [2] S. V. Astashkin and L. Maligranda, *Geometry of Cesàro function spaces*, in preparation.
- [3] G. Bennett, *Factorizing the Classical Inequalities*, Mem. Amer. Math. Soc. **120**, Providence, 1996. MR1317938 (96h:26020)
- [4] S. Chen, Y. Cui, H. Hudzik and B. Sims, *Geometric properties related to fixed point theory in some Banach function lattices*, in: Handbook on Metric Fixed Point Theory, Kluwer Acad. Publ., Dordrecht, 2001, 339–389. MR1904283 (2003f:46031)
- [5] Y. Cui and H. Hudzik, *Some geometric properties related to fixed point theory in Cesàro sequence spaces*, Collect. Math. **50**(1999), 277–288. MR1744077 (2001f:46033)
- [6] Y. Cui, H. Hudzik and Y. Li, *On the García-Falset coefficient in some Banach sequence spaces*, in: Function Spaces, Proc. of the 5th International Conf. in Poznan (Aug. 28–Sept. 3, 1998), Marcel Dekker, New York, 2000, 141–148. MR1772119 (2001h:46009)
- [7] Y. Cui, C. Meng and R. Pluciennik, *Banach-Saks property and property (β) in Cesàro sequence spaces*, Southeast Asian Bull. Math. **24**(2000), 201–210. MR1810056 (2001m:46031)
- [8] S. Dilworth, M. Girardi and J. Hagler, *Dual Banach spaces which contain an isometric copy of L_1* , Bull. Polish Acad. Sci. Math. **48**(2000), no. 1, 1–12. MR1751149 (2001e:46016)
- [9] P. N. Dowling and C. J. Lennard, *Every nonreflexive subspace of $L_1[0, 1]$ fails the fixed point property*, Proc. Amer. Math. Soc. **125**(1997), no. 2, 443–446. MR1350940 (97d:46034)
- [10] P. N. Dowling, C. J. Lennard and B. Turett, *Renormings of l^1 and c_0 and fixed point properties*, In: Handbook of Metric Fixed Point Theory, Kluwer Acad. Publ., Dordrecht, 2001, 269–297. MR1904280 (2004f:46021)
- [11] G. H. Hardy, J. E. Littlewood and G. Pólya, *Inequalities*, Cambridge Univ. Press, 1952. MR0046395 (13:727e)
- [12] B. D. Hassard and D. A. Hussein, *On Cesàro function spaces*, Tamkang J. Math. **4**(1973), 19–25. MR0333700 (48:12025)
- [13] A. Kufner, L. Maligranda and L.-E. Persson, *The Hardy Inequality. About Its History and Some Related Results*, Vydavatelski Servis Publishing House, Pilzen, 2007. MR2351524
- [14] Y. Q. Liu, B. E. Wu and P. Y. Lee, *Method of Sequence Spaces*, Guangdong of Science and Technology Press, 1996 (in Chinese).
- [15] L. Maligranda, N. Petrot and S. Suantai, *On the James constant and B -convexity of Cesàro and Cesàro-Orlicz sequence spaces*, J. Math. Anal. Appl. **326**(2007), no. 1, 312–331. MR2277785 (2007m:46023)
- [16] J. S. Shiue, *A note on Cesàro function space*, Tamkang J. Math. **1**(1970), 91–95. MR0276751 (43:2491)
- [17] P. W. Sy, W. Y. Zhang and P. Y. Lee, *The dual of Cesàro function spaces*, Glas. Mat. Ser. III **22**(1987), no. 1, 103–112. MR940098 (89g:46059)

DEPARTMENT OF MATHEMATICS AND MECHANICS, SAMARA STATE UNIVERSITY, ACAD. PAVLOV 1, 443011 SAMARA, RUSSIA

E-mail address: `astashkn@ssu.samara.ru`

DEPARTMENT OF MATHEMATICS, LULEÅ UNIVERSITY OF TECHNOLOGY, SE-971 87 LULEÅ, SWEDEN

E-mail address: `lech@sm.luth.se`