A NOTE ON THE $2k$ -TH POWER MEAN OF CHARACTER SUMS OVER THE QUARTER INTERVAL

ZHEFENG XU

(Communicated by Wen-Ching Winnie Li)

ABSTRACT. We obtain an asymptotic formula for the $2k$ -th power mean of odd primitive character sums over the interval $[1, \frac{q}{4})$.

1. INTRODUCTION

Let $q \geq 3$ be an integer, χ be a Dirichlet character modulo q . The various arithmetical properties of the character sums

$$\sum_{a=N+1}^{N+H} \chi(a)$$

were investigated by many authors; see [1], [2], [3], [4]. D.A. Burgess [5] obtained the mean value estimate

$$\sum_{\chi \bmod q}^* \sum_{n=1}^p \left| \sum_{m=1}^h \chi(n+m) \right|^4 \leq 8\tau^7(q)q^2h^2,$$

where $\sum_{\chi \bmod q}^*$ denotes the sum over all primitive characters modulo q and $\tau(n)$ is the divisor function.

The author and Zhang [6] studied the $2k$ -th power mean of the even primitive character sums over the quarter interval $[1, \frac{q}{4})$ by using mean value theorems of Dirichlet L -functions, and obtained the asymptotic formula as follows:

$$\begin{aligned} & \sum_{\chi(-1)=1}^* \left| \sum_{a < \frac{q}{4}} \chi(a) \right|^{2k} \\ &= \frac{J(q)q^k}{16} \left(\frac{\pi}{8} \right)^{2k-2} \prod_{p|q} \left(1 - \frac{1}{p^2} \right)^{2k-1} \prod_{p \nmid 2q} \left(1 - \frac{1 - C_{2k-2}^{k-1}}{p^2} \right) + O(q^{k+\epsilon}), \end{aligned}$$

Received by the editors November 8, 2007.

2000 *Mathematics Subject Classification*. Primary 11L40, 11L10; Secondary 11M38.

Key words and phrases. Odd primitive character sums, quarter interval, mean value, asymptotic formula.

The author was supported by NSF(10671155) of the People's Republic of China.

©2008 American Mathematical Society
 Reverts to public domain 28 years from publication

where $\sum_{\chi(-1)=1}^*$ denotes the sum over all primitive characters modulo q such that $\chi(-1) = 1$, ϵ is any fixed positive number, $J(q)$ denotes the number of all primitive characters modulo q , $\prod_{p|q}$ denotes the product over all prime divisors p of q , and

$C_m^n = \frac{m!}{n!(m-n)!}$. Because Lemma 3 of reference [8] is not correct, which will be explained in our Lemma 2.6, this result is also incorrect. The last factor in the main term $\left(1 - \frac{1-C_{2k-2}^{k-1}}{p^2}\right)$ should be $A(0, k, p, 2)$, as defined in our theorem below. However, this result is correct in the case $k = 2$.

In this paper we study odd primitive character sums over the interval $[1, \frac{q}{4})$ by transforming them to L -functions, and we obtain an asymptotic formula for the $2k$ -th power mean using the same method as in [6].

Theorem 1.1. *Let $q \geq 5$ be an odd integer. Then we have the asymptotic formula*

$$\sum_{\chi(-1)=-1}^* \left| \sum_{a < \frac{q}{4}} \chi(a) \right|^{2k} = C(k) q^k J(q) \zeta^{2k-1}(2) \prod_{p|q} \left(1 - \frac{1}{p^2}\right)^{2k-1} \prod_{p \nmid 2q} A(0, k, p, 2) + O_k(q^{k+\epsilon}),$$

where

$$C(k) = \frac{1}{\pi^{2k}} \sum_{i=0}^k C_k^i (-2)^{k-i} \sum_{j=0}^i 6^j \sum_{s=0}^{k-i} \sum_{t=0}^{i-j} \frac{A(|3i+4s-j-2t-2k|, k, 2, 2)}{2^{|3i+4s-j-2t-2k|+2k+1}},$$

$$A(m, k, p, s) = \sum_{i=0}^{2k-2} \frac{1}{p^{is}} \sum_{j=0}^i (-1)^j C_{2k-1}^j C_{k+m+i-j-1}^{m+i-j} C_{k+i-j-1}^{i-j}$$

and $\zeta(s)$ denotes the Riemann zeta function.

Taking $k = 2$ in our theorem and noting that

$$\begin{aligned} & \prod_{p|q} \left(1 - \frac{1}{p^2}\right)^3 \prod_{p \nmid 2q} \left(1 + \frac{1}{p^2}\right) \\ &= \prod_p \left(1 + \frac{1}{p^2}\right) \prod_{p|q} \left(1 - \frac{1}{p^2}\right)^3 \prod_{p \nmid 2q} \frac{1}{1 + \frac{1}{p^2}} \\ &= \frac{4}{5} \frac{\zeta(2)}{\zeta(4)} \prod_{p|q} \frac{(p^2 - 1)^3}{p^4(p^2 + 1)}, \end{aligned}$$

$\zeta(2) = \frac{\pi^2}{6}$ and $\zeta(4) = \frac{\pi^4}{90}$, we get

Corollary 1.2. *Let $q \geq 5$ be an odd integer. Then*

$$\sum_{\chi(-1)=-1}^* \left| \sum_{a < \frac{q}{4}} \chi(a) \right|^4 = \frac{9}{128} q^2 J(q) \prod_{p|q} \frac{(p^2 - 1)^3}{p^4(p^2 + 1)} + O(q^{2+\epsilon}).$$

Combining this with Corollary 2 of [6], we have the following

Corollary 1.3. *Let $q \geq 5$ be an odd integer. Then we have the asymptotic formula*

$$\sum_{\chi \bmod q}^* \left| \sum_{a < \frac{q}{4}} \chi(a) \right|^4 = \frac{21}{256} q^2 J(q) \prod_{p|q} \frac{(p^2 - 1)^3}{p^4(p^2 + 1)} + O(q^{2+\epsilon}).$$

The number $n > 1$ is called square-full if in the prime factorization $n = p_1^{\alpha_1} p_2^{\alpha_2} \cdots p_r^{\alpha_r}$ we have $\alpha_i \geq 2$ for all $i = 1, 2, \dots, r$. Noting that $J(q) = \frac{\phi^2(q)}{q}$ if q is a square-full number, we have

Corollary 1.4. *Let $q \geq 5$ be a square-full number with $2 \nmid q$. Then we have the asymptotic formula*

$$\sum_{\chi \bmod q}^* \left| \sum_{a < \frac{q}{4}} \chi(a) \right|^4 = \frac{21}{256} q \phi^2(q) \prod_{p|q} \frac{(p^2 - 1)^3}{p^4(p^2 + 1)} + O(q^{2+\epsilon}).$$

Corollary 1.5. *Let $p \geq 5$ be a prime. Then we have the asymptotic formula*

$$\sum_{\substack{\chi \bmod p \\ \chi \neq \chi_0}} \left| \sum_{a < \frac{p}{4}} \chi(a) \right|^4 = \frac{21}{256} p^3 + O(p^{2+\epsilon}).$$

Remark 1.6. From our theorem and the theorem in [6], we find that odd and even primitive characters have a very different contribution to the higher moment of character sums over the quarter interval. For the case $k = 2$, from Corollaries 1.2 and 1.3, we see that the contribution of odd primitive characters is six times that of the even ones, although (see Remark 2.4 in the next section)

$$\sum_{a=1}^{\lfloor \frac{q}{4} \rfloor} \chi(a) = 0$$

for some odd primitive characters, while

$$\sum_{a=1}^{\lfloor \frac{q}{4} \rfloor} \chi(a) \neq 0$$

for all even primitive characters.

2. SOME LEMMAS

To prove the theorem, we need the following lemmas.

Lemma 2.1. *Let χ be a primitive character modulo m with $\chi(-1) = -1$. Then we have*

$$\frac{1}{m} \sum_{b=1}^m b \chi(b) = \frac{i}{\pi} \tau(\chi) L(1, \bar{\chi}),$$

where $\tau(\chi) = \sum_{a=1}^m \chi(a) e\left(\frac{a}{m}\right)$ is the Gauss sum, $e(y) = e^{2\pi i y}$, and $L(s, \chi)$ denotes the Dirichlet L -function corresponding to χ .

Proof. This can be easily deduced from Theorem 12.11 and Theorem 12.20 of [7]. $\square$

Lemma 2.2. *Let $q \geq 3$ be an odd number. For any nonprincipal character $\chi \bmod q$, we have*

$$\sum_{a=1}^q a\chi(a) = \frac{\chi(2)q}{1-2\chi(2)} \sum_{a=1}^{\frac{q-1}{2}} \chi(a).$$

Proof. From the properties of Dirichlet characters, we have

$$\begin{aligned} \sum_{a=1}^q 2a\chi(2a) &= \sum_{a=1}^{\frac{q-1}{2}} 2a\chi(2a) + \sum_{a=\frac{q+1}{2}}^q 2a\chi(2a) \\ &= \sum_{a=1}^{\frac{q-1}{2}} 2a\chi(2a) + \sum_{a=1}^{\frac{q+1}{2}} (2a-1)\chi(q+2a-1) + q \sum_{a=1}^{\frac{q+1}{2}} \chi(2a-1) \\ &= \sum_{a=1}^q a\chi(a) + q \sum_{a=1}^{\frac{q+1}{2}} \chi(2a-1). \end{aligned}$$

Noting that

$$\sum_{a=1}^{\frac{q+1}{2}} \chi(2a-1) + \sum_{a=1}^{\frac{q-1}{2}} \chi(2a) = \sum_{a=1}^q \chi(a) = 0,$$

we can write

$$(1-2\chi(2)) \sum_{a=1}^q a\chi(a) = \sum_{a=1}^q a\chi(a) - \sum_{a=1}^q 2a\chi(2a) = q \sum_{a=1}^{\frac{q-1}{2}} \chi(2a) = \chi(2)q \sum_{a=1}^{\frac{q-1}{2}} \chi(a).$$

That is,

$$\sum_{a=1}^q a\chi(a) = \frac{\chi(2)q}{1-2\chi(2)} \sum_{a=1}^{\frac{q-1}{2}} \chi(a).$$

This proves Lemma 2.2. $\square$

Lemma 2.3. *Let q be an odd number and χ be a primitive Dirichlet character modulo q such that $\chi(-1) = -1$. Then we have*

$$\sum_{a=1}^{\left[\frac{q}{4}\right]} \chi(a) = \frac{2 + \bar{\chi}(2) - \bar{\chi}(4)}{2i\pi} \tau(\chi) L(1, \bar{\chi}).$$

Remark 2.4. For even primitive characters χ , by Lemma 2 of [6],

$$\sum_{a=1}^{\left[\frac{q}{4}\right]} \chi(a) = -\frac{i\bar{\chi}(4)}{2\pi} \tau(\chi\chi_4) L(1, \bar{\chi}\chi_4),$$

where χ_4 is the primitive character modulo 4. Since $\tau(\chi\chi_4) L(1, \bar{\chi}\chi_4) \neq 0$, we have

$$\sum_{a=1}^{\left[\frac{q}{4}\right]} \chi(a) \neq 0$$

for all even primitive characters. For odd primitive characters, from the identity in Lemma 2.3, we know that

$$\sum_{a=1}^{\left[\frac{q}{4}\right]} \chi(a) = 0$$

if and only if $\chi(2) = -1$. For example, if p is a prime, then the real character sum

$$\sum_{a=1}^{\left[\frac{p-1}{4}\right]} \left(\frac{a}{p}\right) = 0$$

if and only if $p \equiv 3 \pmod{8}$. In this case, the number of quadratic residues and quadratic non-residues are equal in $\left[1, \frac{p-1}{4}\right]$.

Proof of Lemma 2.3. We separate q into two cases: $q \equiv 1 \pmod{4}$ and $q \equiv 3 \pmod{4}$. First, we suppose $q \equiv 1 \pmod{4}$. From the properties of the Dirichlet character modulo q , we can write

$$\begin{aligned} 4\chi(4) \sum_{a=1}^{q-1} a\chi(a) &= \sum_{a=1}^{\frac{q-1}{4}} 4a\chi(4a) + \sum_{a=\frac{q+3}{4}}^{\frac{2q-2}{4}} 4a\chi(4a) + \sum_{a=\frac{2q+2}{4}}^{\frac{3q-3}{4}} 4a\chi(4a) + \sum_{a=\frac{3q+1}{4}}^{q-1} 4a\chi(4a) \\ &= \sum_{a=1}^{\frac{q-1}{4}} 4a\chi(4a) + \sum_{a=1}^{\frac{q-1}{4}} (4a+q-1)\chi(4a-1) \\ &\quad + \sum_{a=1}^{\frac{q-1}{4}} (4a+2q-2)\chi(4a-2) + \sum_{a=1}^{\frac{q-1}{4}} (4a+3q-3)\chi(4a-3) \\ &= \sum_{a=1}^{q-1} a\chi(a) + \chi(4)q \sum_{a=1}^{\frac{q-1}{4}} \chi(a-\bar{4}) \\ &\quad + 2\chi(4)q \sum_{a=1}^{\frac{q-1}{4}} \chi(a-2\cdot\bar{4}) + 3\chi(4)q \sum_{a=1}^{\frac{q-1}{4}} \chi(a-3\cdot\bar{4}). \end{aligned} \tag{2.1}$$

Note that $\bar{4} \equiv \frac{3q+1}{4} \pmod{q}$ if $q \equiv 1 \pmod{4}$. So from (2.1), we have

$$\begin{aligned} 4\chi(4) \sum_{a=1}^{q-1} a\chi(a) &= \sum_{a=1}^{q-1} a\chi(a) - \chi(4)q \sum_{a=\frac{2q+2}{4}}^{\frac{3q-3}{4}} \chi(a) \\ &\quad - 2\chi(4)q \sum_{a=\frac{q+3}{4}}^{\frac{2q-2}{4}} \chi(a) - 3\chi(4)q \sum_{a=1}^{\frac{q-1}{4}} \chi(a) \\ &= \sum_{a=1}^{q-1} a\chi(a) - \chi(4)q \sum_{a=\frac{q+3}{4}}^{\frac{2q-2}{4}} \chi(a) - 3\chi(4)q \sum_{a=1}^{\frac{q-1}{4}} \chi(a) \\ &= \sum_{a=1}^{q-1} a\chi(a) - \chi(4)q \sum_{a=1}^{\frac{q-1}{2}} \chi(a) - 2\chi(4)q \sum_{a=1}^{\frac{q-1}{4}} \chi(a), \end{aligned} \tag{2.2}$$

where we used the fact $\chi(-1) = -1$ and

$$\sum_{a=\frac{q+3}{4}}^{\frac{2q-2}{4}} \chi(a) = - \sum_{a=\frac{2q+2}{4}}^{\frac{3q-3}{4}} \chi(a).$$

Now, from (2.2) and Lemma 2.2, we get

$$4\chi(4) \sum_{a=1}^{q-1} a\chi(a) = \sum_{a=1}^{q-1} a\chi(a) - (\chi(2) - 2\chi(4)) \sum_{a=1}^{q-1} a\chi(a) - 2\chi(4)q \sum_{a=1}^{\frac{q-1}{4}} \chi(a).$$

That is,

$$\sum_{a=1}^{\frac{q-1}{4}} \chi(a) = \frac{\bar{\chi}(4) - \bar{\chi}(2) - 2}{2q} \sum_{a=1}^{q-1} a\chi(a) = \frac{\bar{\chi}(4) - \bar{\chi}(2) - 2}{2q} \sum_{a=1}^q a\chi(a).$$

Then from Lemma 2.1, we have

$$(2.3) \quad \sum_{a=1}^{\frac{q-1}{4}} \chi(a) = \frac{2 + \bar{\chi}(2) - \bar{\chi}(4)}{2i\pi} \tau(\chi) L(1, \bar{\chi}).$$

This proves Lemma 2.3 in the case of $q \equiv 1 \pmod{4}$. By the same method, we can also prove

$$(2.4) \quad \sum_{a=1}^{\frac{q-3}{4}} \chi(a) = \frac{2 + \bar{\chi}(2) - \bar{\chi}(4)}{2i\pi} \tau(\chi) L(1, \bar{\chi})$$

if $q \equiv 3 \pmod{4}$. Combining (2.3) and (2.4), we can immediately get

$$\sum_{a=1}^{\left[\frac{q}{4}\right]} \chi(a) = \frac{2 + \bar{\chi}(2) - \bar{\chi}(4)}{2i\pi} \tau(\chi) L(1, \bar{\chi}).$$

This completes the proof of Lemma 2.3. $\square$

Lemma 2.5. Let $f(x)$ be a polynomial of degree k with leading coefficient a_0 , and define the difference operator Δ by $(\Delta f)(x) = f(x) - f(x-1)$. Then we have

$$\Delta^k f(x) = k!a_0, \quad \Delta^l f(x) = 0 \quad (l \geq k+1).$$

Proof. This can be easily deduced by the definition of the difference operator and mathematical induction. $\square$

Lemma 2.6. Let q be an integer with $q > 2$ and let $\tau_k(n)$ denote the k -th divisor function (i.e., the number of solutions of the equation $n_1 n_2 \cdots n_k = n$ in positive integers $n_1, n_2, \dots, n_k$). Then for any complex variable s with $\operatorname{Re}(s) > 1$, we have the identity

$$\sum_{\substack{n=1 \\ (n,q)=1}}^{\infty} \frac{\tau_k^2(n)}{n^s} = \zeta^{2k-1}(s) \prod_{p|q} \left(1 - \frac{1}{p^s}\right)^{2k-1} \prod_{p \nmid q} A(0, k, p, s).$$

Proof. This is Lemma 3 of [8]. But the result in [8] is not correct. From the Euler product formula and noting that (see formula 6.4.12 of [9])

$$\tau_k(n) = \prod_{j=1}^r C_{k+\alpha_j-1}^{\alpha_j}$$

if $n = p_1^{\alpha_1} p_2^{\alpha_2} \cdots p_r^{\alpha_r}$ is the factorization of n into prime powers, we have

$$\sum_{\substack{n=1 \\ (n,q)=1}}^{\infty} \frac{\tau_k^2(n)}{n^s} = \prod_{p \nmid q} S(k, p, s),$$

where

$$S(k, p, s) = 1 + \frac{(C_k^1)^2}{p^s} + \frac{(C_{k+1}^2)^2}{p^s} + \cdots + \frac{(C_{k+i-1}^i)^2}{p^s} + \cdots.$$

Now we calculate $S(k, p, s)$. For convenience, let $f(i) = (C_{k+i-1}^i)^2$. From the definition of the difference operator Δ in Lemma 2.5, we have

$$\begin{aligned} S(k, p, s) \left(1 - \frac{1}{p^s}\right) &= 1 + \frac{\Delta f(1)}{p^s} + \frac{\Delta f(2)}{p^{2s}} + \cdots + \frac{\Delta f(i)}{p^{is}} + \cdots, \\ S(k, p, s) \left(1 - \frac{1}{p^s}\right)^2 &= 1 + \frac{\Delta f(1) - f(0)}{p^s} + \frac{\Delta^2 f(2)}{p^{2s}} + \cdots + \frac{\Delta^2 f(i)}{p^{is}} + \cdots. \end{aligned}$$

So, from the binomial expansion theorem and comparing the coefficients of p^{is} for $1 \leq i \leq 2k-2$, we have

$$S(k, p, s) \left(1 - \frac{1}{p^s}\right)^{2k-1} = S'(k, p, s) + \frac{\Delta^{2k-1} f(2k-1)}{p^{(2k-1)s}} + \frac{\Delta^{2k} f(2k)}{p^{2ks}} + \cdots;$$

here

$$S'(k, p, s) = \sum_{i=0}^{2k-2} \frac{1}{p^{is}} \sum_{j=0}^i (-1)^j C_{2k-1}^j f(i-j).$$

Noting that $f(i)$ is a polynomial in i of degree $2k-2$, so from Lemma 2.5 we know that $\Delta^{2k-1} f(2k-1) = \Delta^{2k} f(2k) = \cdots = 0$. This yields

$$\begin{aligned} S(k, p, s) &= \left(1 - \frac{1}{p^s}\right)^{1-2k} \sum_{i=0}^{2k-2} \frac{1}{p^{is}} \sum_{j=0}^i (-1)^j C_{2k-1}^j (C_{k+i-j-1}^{i-j})^2 \\ &= \left(1 - \frac{1}{p^s}\right)^{1-2k} A(0, k, p, s). \end{aligned}$$

Hence, Lemma 2.6 follows immediately from the fact $\zeta(s) = \prod_p \left(1 - \frac{1}{p^s}\right)^{-1}$. $\square$

Lemma 2.7. *Let q be any odd integer with $q > 2$ and $m \geq 0$ a fixed integer. Then for any complex variable s with $\operatorname{Re}(s) > 1$, we have the identity*

$$\sum_{\substack{n=1 \\ (n,q)=1}}^{\infty} \frac{\tau_k(2^m n) \tau_k(n)}{n^s} = A(m, k, 2, s) \zeta^{2k-1}(s) \prod_{p|q} \left(1 - \frac{1}{p^s}\right)^{2k-1} \prod_{p \nmid 2q} A(0, k, p, s).$$

This lemma is a generalization of Lemma 2.6, which is just the case $m = 0$.

Proof. Noting that $\tau_k(n)$ is a multiplicative function, we can write

$$\begin{aligned} \sum_{\substack{n=1 \\ (n,q)=1}}^{\infty} \frac{\tau_k(2^m n) \tau_k(n)}{n^s} &= \tau_k(2^m) \sum_{\substack{n=1 \\ (n,q)=1 \\ 2 \nmid n}}^{\infty} \frac{\tau_k^2(n)}{n^s} + \sum_{\substack{n=1 \\ (n,q)=1 \\ 2 \mid n}}^{\infty} \frac{\tau_k(2^m n) \tau_k(n)}{n^s} \\ &= \tau_k(2^m) \sum_{\substack{n=1 \\ (n,q)=1 \\ 2 \nmid n}}^{\infty} \frac{\tau_k^2(n)}{n^s} + \sum_{\substack{r=1 \\ (r,q)=1 \\ 2 \nmid r}}^{\infty} \sum_{j=1}^{\infty} \frac{\tau_k(2^{m+j}) \tau_k(2^j) \tau_k^2(r)}{(r \cdot 2^j)^s} \\ &= \left(\sum_{j=0}^{\infty} \frac{\tau_k(2^{m+j}) \tau_k(2^j)}{2^{sj}} \right) \sum_{\substack{n=1 \\ (n,q)=1 \\ 2 \nmid n}}^{\infty} \frac{\tau_k^2(n)}{n^s}. \end{aligned}$$

For $n > 1$, let $n = p_1^{\alpha_1} p_2^{\alpha_2} \cdots p_r^{\alpha_r}$ be the factorization of n into prime powers. Then we have

$$(2.5) \quad \sum_{\substack{n=1 \\ (n,q)=1}}^{\infty} \frac{\tau_k(2^m n) \tau_k(n)}{n^s} = \left(\sum_{j=0}^{\infty} \frac{C_{k+m+j-1}^{m+j} C_{k+j-1}^j}{2^{sj}} \right) \sum_{\substack{n=1 \\ (n,q)=1 \\ 2 \nmid n}}^{\infty} \frac{\tau_k^2(n)}{n^s}.$$

Let

$$S = \sum_{j=0}^{\infty} \frac{C_{k+m+j-1}^{m+j} C_{k+j-1}^j}{2^{sj}}.$$

Now using the same method as calculating $S(k, p, s)$ in Lemma 2.6, we can also get

$$\begin{aligned} S &= \left(1 - \frac{1}{2^s}\right)^{1-2k} \sum_{i=0}^{2k-2} \frac{1}{2^{is}} \sum_{j=0}^i (-1)^j C_{2k-1}^j C_{k+m+i-j-1}^{m+i-j} C_{k+i-j-1}^{i-j} \\ (2.6) \quad &= \left(1 - \frac{1}{2^s}\right)^{1-2k} A(m, k, 2, s). \end{aligned}$$

Noting that

$$\sum_{\substack{n=1 \\ (n,q)=1 \\ 2 \nmid n}}^{\infty} \frac{\tau_k^2(n)}{n^s} = \sum_{\substack{n=1 \\ (n,2q)=1}}^{\infty} \frac{\tau_k^2(n)}{n^s}$$

if q is odd, we can easily get Lemma 2.7 from (2.5), (2.6) and Lemma 2.6. $\square$

Lemma 2.8. *Let q be any odd integer with $q > 2$, χ be the Dirichlet character modulo q and $m \geq 1$ be a fixed integer. Then we have the following asymptotic formulas:*

$$\sum_{\chi(-1)=-1}^* \chi(2^m) |L(1, \bar{\chi})|^{2k} = \frac{J(q)}{2^{m+1}} \sum_{\substack{n=1 \\ (n,q)=1}}^{\infty} \frac{\tau_k(2^m n) \tau_k(n)}{n^2} + O(q^\epsilon)$$

and

$$\sum_{\chi(-1)=-1}^* \bar{\chi}(2^m) |L(1, \bar{\chi})|^{2k} = \frac{J(q)}{2^{m+1}} \sum_{\substack{n=1 \\ (n,q)=1}}^{\infty} \frac{\tau_k(2^m n) \tau_k(n)}{n^2} + O(q^\epsilon).$$

Proof. By using the same method as in proving Lemma 5 of [6], we can get these formulas. $\square$

3. PROOF OF THE THEOREM

In this section we will complete the proof of the theorem. It is well known that $|\tau(\chi)| = \sqrt{q}$ if χ is a primitive character. So from Lemma 2.3, we can write

$$\begin{aligned} & \sum_{\chi(-1)=-1}^* \left| \sum_{a < \frac{q}{4}} \chi(a) \right|^{2k} \\ &= \frac{q^k}{(2\pi)^{2k}} \sum_{\chi(-1)=-1}^* |2 + \bar{\chi}(2) - \bar{\chi}(4)|^{2k} |L(1, \bar{\chi})|^{2k} \\ &= \frac{q^k}{(2\pi)^{2k}} \sum_{\chi(-1)=-1}^* [6 + \bar{\chi}(2) + \chi(2) - 2(\bar{\chi}(4) + \chi(4))]^k |L(1, \bar{\chi})|^{2k} \\ &= \frac{q^k}{(2\pi)^{2k}} \sum_{\chi(-1)=-1}^* \sum_{i=0}^k C_k^i (6 + \bar{\chi}(2) + \chi(2))^i (-2)^{k-i} (\bar{\chi}(4) + \chi(4))^{k-i} |L(1, \bar{\chi})|^{2k} \\ &= \frac{q^k}{(2\pi)^{2k}} \sum_{i=0}^k C_k^i (-2)^{k-i} \sum_{j=0}^i 6^j \sum_{s=0}^{k-i} \sum_{t=0}^{i-j} \sum_{\chi(-1)=-1}^* \chi^{3i+4s-j-2t-2k}(2) |L(1, \bar{\chi})|^{2k}, \end{aligned}$$

where χ^{-1} has the same meaning as $\bar{\chi}$, the conjugate of the character χ . Then from Lemma 2.8 and Lemma 2.7, we easily get

$$\begin{aligned} & \sum_{\chi(-1)=-1}^* \left| \sum_{a < \frac{q}{4}} \chi(a) \right|^{2k} \\ &= C(k) q^k J(q) \zeta^{2k-1}(2) \prod_{p|q} \left(1 - \frac{1}{p^2}\right)^{2k-1} \prod_{p \nmid 2q} A(0, k, p, 2) + O_k(q^{k+\epsilon}), \end{aligned}$$

where

$$C(k) = \frac{1}{\pi^{2k}} \sum_{i=0}^k C_k^i (-2)^{k-i} \sum_{j=0}^i 6^j \sum_{s=0}^{k-i} \sum_{t=0}^{i-j} \frac{A(|3i+4s-j-2t-2k|, k, 2, 2)}{2^{|3i+4s-j-2t-2k|+2k+1}},$$

$$A(m, k, p, s) = \sum_{i=0}^{2k-2} \frac{1}{p^{is}} \sum_{j=0}^i (-1)^j C_{2k-1}^j C_{k+m+i-j-1}^{m+i-j} C_{k+i-j-1}^{i-j}.$$

This completes the proof of the theorem.

ACKNOWLEDGMENT

The author expresses his gratitude to the referee for very helpful and detailed comments.

REFERENCES

1. G. Pólya, *Über die Verteilung der quadratische Reste und Nichtreste*, Göttingen Nachrichten, 1918, pp. 21–29.
2. I. M. Vinogradov, *On the distribution of residues and non-residues of powers*, Journal of the Physico-Mathematical Society of Perm. **1** (1918), 94–96.
3. A. V. Sokolovskii, *On a theorem of Sárkozy*, Acta Arithmetica **41** (1982), 27–31. MR667707 (84e:10046)
4. M. Z. Garaev and A. A. Karatsuba, *On character sums and the exceptional set of a congruence problem*, Journal of Number Theory **114** (2005), 182–192. MR2163912 (2006f:11095)
5. D. A. Burgess, *Mean value of character sums. II*, Mathematika **33** (1987), 1–7. MR908834 (89g:11072)
6. Zhefeng Xu and Wenpeng Zhang, *On the 2kth power mean of the character sums over short intervals*, Acta Arithmetica **121** (2006), 149–160. MR2216139 (2006m:11122)
7. Tom M. Apostol, *Introduction to Analytic Number Theory*, Springer-Verlag, New York, 1976. MR0434929 (55:7892)
8. Wenpeng Zhang, Yuan Yi and Xiali He, *On the 2k-th power mean of Dirichlet L-functions with the weight of general Kloosterman sums*, Journal of Number Theory **84** (2000), 199–213. MR1795790 (2002e:11115)
9. Chengdong Pan and Chengbiao Pan, *Element of Analytic Number Theory*, Science Press, Beijing, 1999, pp. 95.

DEPARTMENT OF MATHEMATICS, NORTHWEST UNIVERSITY, XI'AN, SHAANXI, PEOPLE'S REPUBLIC OF CHINA

E-mail address: zhefengxu@hotmail.com; zfxu@nwu.edu.cn