RESONANCE OF BASIS-CONJUGATING AUTOMORPHISM GROUPS

DANIEL C. COHEN

(Communicated by Daniel Ruberman)

ABSTRACT. We determine the structure of the first resonance variety of the cohomology ring of the group of automorphisms of a finitely generated free group which act by conjugation on a given basis.

1. RESONANCE OF $P\Sigma_n$

Let F_n be the free group generated by $x_1, \dots, x_n$. The basis-conjugating automorphism group, or pure symmetric automorphism group, is the group $P\Sigma_n$ of all automorphisms of F_n which send each generator x_i to a conjugate of itself. Results of Dahm [4] and Goldsmith [7] imply that this group may also be realized as the “group of loops”, the group of motions of a collection of n unknotted, unlinked oriented circles in 3-space, where each circle returns to its original position. McCool [9] found the following presentation for the basis-conjugating automorphism group:

$$(1.1) \quad P\Sigma_n = \langle \beta_{i,j}, 1 \leq i \neq j \leq n \mid [\beta_{i,j}, \beta_{k,l}], [\beta_{i,k}, \beta_{j,k}], [\beta_{i,j}, (\beta_{i,k} \cdot \beta_{j,k})] \rangle,$$

where $[u, v] = uvu^{-1}v^{-1}$ denotes the commutator, the indices in the relations are distinct, and the generators $\beta_{i,j}$ are the automorphisms of F_n defined by

$$\beta_{i,j}(x_k) = \begin{cases} x_k & \text{if } k \neq j, \\ x_j^{-1}x_i x_j & \text{if } k = i. \end{cases}$$

The purpose of this paper is to determine the structure of the first resonance variety of the cohomology ring of this group.

Let $A = \bigoplus_{k=0}^{\ell} A^k$ be a finite-dimensional, graded, connected algebra over an algebraically closed field $\mathbb{k}$ of characteristic 0. Since $a \cdot a = 0$ for each $a \in A^1$, multiplication by a defines a cochain complex (A, δ_a) :

$$A^0 \xrightarrow{\delta_a} A^1 \xrightarrow{\delta_a} A^2 \xrightarrow{\delta_a} \dots \xrightarrow{\delta_a} A^{\ell},$$

where $\delta_a(x) = ax$. The resonance varieties of A are the jumping loci for the cohomology of these complexes: $R_d^j(A) = \{a \in A^1 \mid \dim_{\mathbb{k}} H^j(A, \delta_a) \geq d\}$. As shown by Falk [6], these algebraic subvarieties of A^1 are isomorphism-type invariants of the algebra A . They have been the subject of considerable recent interest in the context of hyperplane arrangements and related areas; see, for instance, Dimca,

Received by the editors November 17, 2008.

2000 *Mathematics Subject Classification*. Primary 20F28; Secondary 14M12, 20J06.

The author was partially supported by National Security Agency grant No. H98230-05-1-0055.

©2009 American Mathematical Society
 Reverts to public domain 28 years from publication

Papadima, and Suciu [5], Yuzvinsky [14], and the references therein. We will focus on the first resonance variety $R^1(A) = \{a \in A^1 \mid H^1(A, \delta_a) \neq 0\}$.

Since the relations in the presentation (1.1) of $P\Sigma_n$ are all commutators, the first homology group $H_1(P\Sigma_n; \mathbb{k})$ is a vector space of dimension $n(n-1)$ with basis $\{[\beta_{p,q}] \mid 1 \leq p \neq q \leq \ell\}$. Let $\{e_{p,q} \mid 1 \leq p \neq q \leq \ell\}$ be the dual basis of $H^1(P\Sigma_n; \mathbb{k})$. Denote the first resonance variety of $A = H^*(P\Sigma_n; \mathbb{k})$ by $R^1(P\Sigma_n, \mathbb{k})$.

Theorem 1.1. *The first resonance variety of the cohomology ring $H^*(P\Sigma_n; \mathbb{k})$ of the basis-conjugating automorphism group is*

$$R^1(P\Sigma_n, \mathbb{k}) = \bigcup_{1 \leq i < j \leq n} C_{i,j} \cup \bigcup_{1 \leq i < j < k \leq n} C_{i,j,k},$$

where $C_{i,j} = \text{span}\{e_{i,j}, e_{j,i}\}$ and $C_{i,j,k} = \text{span}\{e_{j,i} - e_{k,i}, e_{i,j} - e_{k,j}, e_{i,k} - e_{j,k}\}$.

This result reveals an interesting relationship between the resonance variety and the Bieri-Neumann-Strebel (BNS) invariant of the basis-conjugating automorphism group. For a finitely generated group G , let $\mathcal{C}$ be the Cayley graph corresponding to a finite generating set. Given an additive character $\chi: G \rightarrow \mathbb{R}$, let $\mathcal{C}_+(\chi)$ be the full subgraph of $\mathcal{C}$ on the vertex set $\{g \in G \mid \chi(g) \geq 0\}$. Then the (first) BNS invariant of G is the conical subset $\Sigma(G)$ of $\text{Hom}(G, \mathbb{R}) = H^1(G; \mathbb{R})$ defined by

$$\Sigma(G) = \{\chi \in \text{Hom}(G, \mathbb{R}) \setminus \{0\} \mid \mathcal{C}_+(\chi) \text{ is connected}\}.$$

This invariant of G (which is independent of the choice of generating set) may be used to determine which subgroups above the commutator subgroup $[G, G]$ are finitely generated; see [1].

The BNS invariant of the group $G = P\Sigma_n$ was determined by Orlandi-Korner [11]. Combining her result with the above theorem yields the following:

Theorem 1.2. *The Bieri-Neumann-Strebel invariant of the basis-conjugating automorphism group is given by*

$$\Sigma(P\Sigma_n) = H^1(P\Sigma_n; \mathbb{R}) \setminus R^1(P\Sigma_n, \mathbb{R}).$$

This relationship between the resonance variety and the BNS invariant is known to hold for certain other groups, including right-angled Artin groups; see Meier and VanWyk [10] and Papadima and Suciu [12]. More recently, Papadima and Suciu have shown that the containment $\Sigma(G) \subseteq H^1(G; \mathbb{R}) \setminus R^1(G, \mathbb{R})$ holds for an arbitrary 1-formal group G , where, as above, $R^1(G; \mathbb{R})$ denotes the first resonance variety of $H^*(G; \mathbb{R})$. See [13] for a detailed investigation of the relationship between BNS invariants and (co)homology jumping loci in a number of contexts.

2. PROOF OF THEOREM 1.1

In this section, we recall the structure of the cohomology ring of the basis-conjugating automorphism group and use it to prove Theorem 1.1. The cohomology of $P\Sigma_n$ was computed by Jensen, McCammond, and Meier [8], resolving positively a conjecture of Brownstein and Lee [2].

Theorem 2.1 ([8]). *Let $E_{\mathbb{Z}}$ denote the exterior algebra over $\mathbb{Z}$ generated by degree one elements $e_{p,q}$, $1 \leq p \neq q \leq n$, and let $I_{\mathbb{Z}}$ denote the two-sided ideal in $E_{\mathbb{Z}}$ generated by*

$$(2.1) \quad \begin{aligned} \eta_{i,j} &= e_{i,j}e_{j,i}, \quad 1 \leq i < j \leq n; \\ \tau_{i,j}^k &= (e_{k,i} - e_{j,i})(e_{k,j} - e_{i,j}), \quad 1 \leq k \leq n, \quad 1 \leq i < j \leq n, \quad k \notin \{i, j\}. \end{aligned}$$

Then the integral cohomology algebra of the basis-conjugating automorphism group $P\Sigma_n$ is isomorphic to the quotient of $E_{\mathbb{Z}}$ by $I_{\mathbb{Z}}$, $H^*(P\Sigma_n; \mathbb{Z}) \cong E_{\mathbb{Z}}/I_{\mathbb{Z}}$.

Remark 2.2. The above presentation of the cohomology ring $H^*(P\Sigma_n; \mathbb{Z})$ differs slightly from that given in [2, 8] but is easily seen to be equivalent. For instance, the relation in $H^*(P\Sigma_n; \mathbb{Z})$ arising from the generator $\tau_{i,j}^k$ of the ideal I may be obtained by constructing an appropriate linear combination of the relations labeled 2 and 3 in [8, Thm. 6.7].

Let $\mathbb{k}$ be a field of characteristic zero. From Theorem 2.1 and the Universal Coefficient Theorem, the cohomology algebra $H^*(P\Sigma_n; \mathbb{k})$ is isomorphic to $E_{\mathbb{k}}/I_{\mathbb{k}}$, where $E_{\mathbb{k}}$ is the exterior algebra over $\mathbb{k}$ generated by $e_{p,q}$, $1 \leq p \neq q \leq n$, and $I_{\mathbb{k}}$ is the ideal in $E_{\mathbb{k}}$ generated by the elements $\eta_{i,j}$ and $\tau_{i,j}^k$ above.

Recall from Section 1 that the first resonance variety of $A = H^*(P\Sigma_n; \mathbb{k})$ is

$$R^1(P\Sigma_n, \mathbb{k}) = \{a \in A^1 \mid H^1(A, \delta_a) \neq 0\}.$$

Observe that $A^1 = H^1(P\Sigma_n; \mathbb{k})$ is a vector space of dimension $N = n(n-1)$ over $\mathbb{k}$. Elements of A^1 are of the form $a = \sum_{p \neq q} a_{p,q} e_{p,q}$, where $a_{p,q} \in \mathbb{k}$. Recall that

$$(2.2) \quad C_{i,j} = \text{span}\{e_{i,j}, e_{j,i}\} = \{a \in A^1 \mid a_{p,q} = 0 \text{ if } \{p,q\} \neq \{i,j\}\}$$

for $1 \leq i < j \leq n$ and that

$$(2.3) \quad C_{i,j,k} = \text{span}\{e_{j,i} - e_{k,i}, e_{i,j} - e_{k,j}, e_{i,k} - e_{j,k}\} \\ = \left\{ a \in A^1 \mid \begin{array}{l} a_{j,i} + a_{k,i} = 0, \quad a_{i,j} + a_{k,j} = 0, \quad a_{i,k} + a_{j,k} = 0, \\ a_{p,q} = 0 \text{ if } \{p,q\} \not\subset \{i,j,k\} \end{array} \right\}$$

for $1 \leq i < j < k \leq n$. Note that these are linear subspaces of $A^1 \cong \mathbb{k}^N$ of dimensions 2 and 3 respectively. To prove Theorem 1.1, we must show that $R^1(P\Sigma_n, \mathbb{k})$ is equal to the union of these linear subspaces.

Proof of Theorem 1.1. In the case $n = 2$, the group $P\Sigma_n = \langle \beta_{1,2}, \beta_{2,1} \rangle \cong F_2$ is a free group, and the theorem asserts that $R^1(P\Sigma_2, \mathbb{k}) = C_{1,2} = H^1(P\Sigma_2; \mathbb{k})$, which is clear. So assume that $n \geq 3$.

Write $R = R^1(P\Sigma_n, \mathbb{k})$ and $C = \bigcup_{1 \leq i < j \leq n} C_{i,j} \cup \bigcup_{1 \leq i < j < k \leq n} C_{i,j,k}$. Observe that $0 \in C$ and $0 \in R$. So it is enough to show that $R \setminus \{0\} = C \setminus \{0\}$.

Write $E = E_{\mathbb{k}}$ and $I = I_{\mathbb{k}}$. For $a \in A^1 = E^1$, we have a short exact sequence of chain complexes $0 \rightarrow (I, \delta_a) \xrightarrow{\iota} (E, \delta_a) \xrightarrow{p} (A, \delta_a) \rightarrow 0$:

$$\begin{array}{ccccccc} & & & I^2 & \xrightarrow{\delta_a} & I^3 & \longrightarrow \cdots \\ & & & \downarrow \iota^2 & & \downarrow \iota^3 & \\ E^0 & \xrightarrow{\delta_a} & E^1 & \xrightarrow{\delta_a} & E^2 & \xrightarrow{\delta_a} & E^3 \longrightarrow \cdots \\ \downarrow p^0 & & \downarrow p^1 & & \downarrow p^2 & & \downarrow p^3 \\ A^0 & \xrightarrow{\delta_a} & A^1 & \xrightarrow{\delta_a} & A^2 & \xrightarrow{\delta_a} & A^3 \longrightarrow \cdots \end{array}$$

where $\iota: I \rightarrow E$ is the inclusion, $p: E \rightarrow A$ the projection, and $\delta_a(x) = ax$. Note that since I is generated in degree two, the maps $p^0: E^0 \rightarrow A^0$ and $p^1: E^1 \rightarrow A^1$ are identity maps. If $a \neq 0$, then the complex (E, δ_a) is acyclic. Consequently, the corresponding long exact cohomology sequence yields

$$H^1(A, \delta_a) \cong H^2(I, \delta_a) = \ker(\delta_a: I^2 \rightarrow I^3) = \ker(\iota^3 \circ \delta_a: I^2 \rightarrow E^3).$$

Thus, $a \in R \setminus \{0\}$ if and only if the map $\psi_a := \iota^3 \circ \delta_a$ fails to inject.

Since the elements $\eta_{i,j}$ and $\tau_{i,j}^k$ recorded in (2.1) generate the ideal I and are of degree two (and are linearly independent in E^2), these elements form a basis for I^2 . We record the images of these basis elements under the map ψ_a . For $1 \leq i < j \leq n$,

$$(2.4) \quad \psi_a(\eta_{i,j}) = \sum_{\{p,q\} \neq \{i,j\}} a_{p,q} e_{p,q} \eta_{i,j} = \sum_{\{p,q\} \neq \{i,j\}} a_{p,q} e_{p,q} e_{i,j} e_{j,i}.$$

For $1 \leq k \leq n$, $1 \leq i < j \leq n$, $k \notin \{i, j\}$,

$$(2.5) \quad \begin{aligned} \psi_a(\tau_{i,j}^k) &= (a_{j,i} + a_{k,i}) e_{j,i} e_{k,i} (e_{k,j} - e_{i,j}) - (a_{i,j} + a_{k,j}) e_{i,j} e_{k,j} (e_{k,i} - e_{j,i}) \\ &\quad + (a_{i,k} e_{i,k} + a_{j,k} e_{j,k}) \tau_{i,j}^k + \sum_{\{p,q\} \not\subset \{i,j,k\}} a_{p,q} e_{p,q} \tau_{i,j}^k. \end{aligned}$$

These calculations immediately yield the containment $C \setminus \{0\} \subseteq R \setminus \{0\}$. If $a \in C_{i,j}$, then $a_{p,q} = 0$ for $\{p,q\} \neq \{i,j\}$. For such an a , we have $\psi_a(\eta_{i,j}) = 0$ by (2.4), so $C_{i,j} \subset R$. If $1 \leq i < j < k \leq n$ and $a \in C_{i,j,k}$, then $a_{j,i} + a_{k,i} = 0$, $a_{i,j} + a_{k,j} = 0$, $a_{i,k} + a_{j,k} = 0$, and $a_{p,q} = 0$ for $\{p,q\} \not\subset \{i,j,k\}$. In this instance, (2.5) may be used to check that $a_{j,i} \tau_{i,j}^k - a_{i,k} \tau_{j,k}^i$ and $a_{i,j} \tau_{i,j}^k - a_{i,k} \tau_{i,k}^j$ are elements of $\ker(\psi_a)$. If $a \in C_{i,j,k}$ is nonzero, at least one of $a_{i,j}, a_{i,k}, a_{j,i}$ must be nonzero. Consequently, ψ_a has nontrivial kernel, and $C_{i,j,k} \subset R$.

Establishing the reverse containment, $R \setminus \{0\} \subseteq C \setminus \{0\}$, is more involved. We will show that $a \notin C$ implies that $a \notin R$. If $a \notin C$, then $a \neq 0$. So assume without loss that $a_{2,1} \neq 0$. Since $a \notin C_{1,2} \subset C$, we must also have $a_{p,q} \neq 0$ for some $\{p,q\} \neq \{1,2\}$. We will consider several cases depending on the relationship between the sets $\{1,2\}$ and $\{p,q\}$.

Case 1. $\{1,2\} \cap \{p,q\} = \emptyset$

Assume first that $\{1,2\}$ and $\{p,q\}$ are disjoint. Note that $n \geq 4$ in this instance. Permuting indices if need be, we may assume that $a \in H^1(P\Sigma_n; \mathbb{k})$ satisfies $a_{2,1} \neq 0$ and $a_{3,4} \neq 0$. We will show that this assumption implies that the map $\psi_a: I^2 \rightarrow E^3$ injects; hence $a \notin R$. Specifically, we will exhibit a subspace $V \subset E^3$ and a projection $\pi: E^3 \rightarrow V$ so that the composition $\pi \circ \psi_a: I^2 \rightarrow V$ is an isomorphism.

Let $\mathcal{V}$ be the union of the sets

$$\begin{aligned} &\{e_{1,2} e_{2,1} e_{3,4}\} \cup \{e_{2,1} e_{i,j} e_{j,i} \mid 1 \leq i < j \leq n, \{i,j\} \neq \{1,2\}\}, \\ &\{e_{3,4} e_{1,2} e_{k,1}, e_{3,4} e_{2,1} e_{1,k}, e_{3,4} e_{1,2} e_{1,k} \mid 3 \leq k \leq n\}, \\ &\{e_{2,1} e_{k,i} e_{k,j}, e_{2,1} e_{j,i} e_{j,k}, e_{2,1} e_{i,k} e_{k,j} \mid i \leq 2 < j < k \leq n \text{ or } 3 \leq i < j < k \leq n\}. \end{aligned}$$

There is a bijection between $\mathcal{V}$ and the set of generators of I^2 given by

$$\begin{aligned} \eta_{1,2} &\leftrightarrow e_{1,2} e_{2,1} e_{3,4}, & \eta_{i,j} &\leftrightarrow e_{2,1} e_{i,j} e_{j,i} & (\{i,j\} \neq \{1,2\}), \\ \tau_{1,2}^k &\leftrightarrow e_{3,4} e_{1,2} e_{k,1}, & \tau_{1,k}^2 &\leftrightarrow e_{3,4} e_{2,1} e_{1,k}, & \tau_{2,k}^1 &\leftrightarrow e_{3,4} e_{1,2} e_{1,k} \quad (3 \leq k \leq n), \\ \tau_{i,j}^k &\leftrightarrow e_{2,1} e_{k,i} e_{k,j}, & \tau_{i,k}^j &\leftrightarrow e_{2,1} e_{j,i} e_{j,k}, & \tau_{j,k}^i &\leftrightarrow e_{2,1} e_{i,k} e_{k,j} \quad (i \leq 2 < j < k \leq n). \end{aligned}$$

In particular, the monomials in $\mathcal{V}$ are distinct. Hence, $\mathcal{V}$ is a linearly independent set in E^3 of cardinality $|\mathcal{V}| = \dim_{\mathbb{k}} I^2 = \binom{n}{2}(n-1)$. Let $V = \text{span } \mathcal{V} \subset E^3$.

Define $\pi: E^3 \rightarrow V$ on basis elements by

$$\pi(e_{a,b} e_{c,d} e_{p,q}) = \begin{cases} e_{a,b} e_{c,d} e_{p,q} & \text{if } e_{a,b} e_{c,d} e_{p,q} \in \mathcal{V}, \\ 0 & \text{if } e_{a,b} e_{c,d} e_{p,q} \notin \mathcal{V}. \end{cases}$$

Then, a calculation using (2.4) and (2.5) reveals that $\pi \circ \psi_a: I^2 \rightarrow V$ is an isomorphism. For instance, ordering the bases of I^2 and V appropriately, one can check that the matrix M of $\pi \circ \psi_a$ has determinant $\det M = y_{2,1}^{m_{2,1}} y_{3,4}^{m_{3,4}} \neq 0$, where $m_{3,4} = 3n - 5$ and $m_{2,1} = \binom{n}{2}(n-1) - m_{3,4}$. Thus, if $a \notin C$ satisfies $a_{2,1} \neq 0$ and $a_{p,q} \neq 0$ for some p, q with $\{1, 2\} \cap \{p, q\} = \emptyset$, then $a \notin R$.

Case 2. $\{1, 2\} \cap \{p, q\} \neq \emptyset$

Now assume that $a \notin C$, $a_{2,1} \neq 0$, and $a_{r,s} = 0$ for all r, s with $\{1, 2\} \cap \{r, s\} = \emptyset$. Since $a \notin C_{1,2} \subset C$, we must have $a_{p,q} \neq 0$ for some p, q with $|\{p, q\} \cap \{1, 2\}| = 1$. Permuting indices if necessary, we may assume that $3 \in \{p, q\}$.

In the case $n = 3$, since $a \notin C_{1,2,3} \subset C$, one of the sums $a_{2,1} + a_{3,1}$, $a_{1,2} + a_{3,2}$, $a_{1,3} + a_{2,3}$ must be nonzero; see (2.3). In this instance, ordering bases appropriately, the map $\psi_a: I^2 \rightarrow E^3$ has matrix

$$(2.6) \quad M_3 = \begin{pmatrix} a_{3,2} & 0 & 0 & 0 & -a_{1,2} - a_{3,2} & 0 \\ a_{3,1} & 0 & 0 & 0 & -a_{2,1} - a_{3,1} & 0 \\ a_{2,3} & 0 & 0 & a_{1,2} & -a_{2,3} & a_{2,1} \\ -a_{1,3} & 0 & 0 & a_{1,2} & a_{1,3} & a_{2,1} \\ 0 & a_{3,2} & 0 & -a_{3,2} & a_{1,3} & -a_{3,1} \\ 0 & -a_{2,3} & 0 & a_{1,3} + a_{2,3} & 0 & 0 \\ 0 & -a_{2,1} & 0 & a_{2,1} + a_{3,1} & 0 & 0 \\ 0 & a_{1,2} & 0 & -a_{1,2} & -a_{1,3} & a_{3,1} \\ 0 & 0 & -a_{3,1} & a_{3,2} & a_{2,3} & a_{3,1} \\ 0 & 0 & a_{2,1} & a_{3,2} & a_{2,3} & -a_{2,1} \\ 0 & 0 & a_{1,3} & 0 & 0 & -a_{1,3} - a_{2,3} \\ 0 & 0 & a_{1,2} & 0 & 0 & -a_{1,2} - a_{3,2} \\ 0 & 0 & 0 & a_{3,2} & -a_{1,3} & -a_{2,1} \\ 0 & 0 & 0 & a_{2,1} + a_{3,1} & 0 & 0 \\ 0 & 0 & 0 & a_{1,3} + a_{2,3} & 0 & 0 \\ 0 & 0 & 0 & a_{1,2} & -a_{2,3} & -a_{3,1} \\ 0 & 0 & 0 & 0 & a_{1,2} + a_{3,2} & 0 \\ 0 & 0 & 0 & 0 & a_{2,1} + a_{3,1} & 0 \\ 0 & 0 & 0 & 0 & 0 & a_{1,2} + a_{3,2} \\ 0 & 0 & 0 & 0 & 0 & a_{1,3} + a_{2,3} \end{pmatrix}.$$

Using the assumptions on $a_{2,1}$, the sums $a_{2,1} + a_{3,1}$, $a_{1,2} + a_{3,2}$, $a_{1,3} + a_{2,3}$, and $a_{p,q} \in \{a_{1,3}, a_{2,3}, a_{3,1}, a_{3,2}\}$, one can check that the matrix M_3 has maximal rank. Hence, if $a \notin C$, then $\psi_a: I^2 \rightarrow E^3$ injects in the case $n = 3$.

For general n , the assumption that $a \notin C_{1,2,3}$ implies that the set

$$\{a_{2,1} + a_{3,1}, a_{1,2} + a_{3,2}, a_{1,3} + a_{2,3}\} \cup \{a_{r,s} \mid \{r, s\} \not\subset \{1, 2, 3\}\}$$

contains a nonzero element. Recall that, by Case 1, we may assume that $a_{r,s} = 0$ for all r, s with $\{1, 2\} \cap \{r, s\} = \emptyset$. If $a_{r,s} = 0$ for all $\{r, s\} \not\subset \{1, 2, 3\}$, let W be the subspace of E^3 spanned by the union of the sets

$$\begin{aligned} & \{e_{i_1, j_1} e_{i_2, j_2} e_{i_3, j_3} \mid 1 \leq i_k, j_k \leq 3, i_k \neq j_k\}, \\ & \{e_{1,2} e_{2,1} e_{p,q}\} \cup \{e_{2,1} e_{i,j} e_{j,i} \mid 1 \leq i < j \leq n, \{i, j\} \neq \{1, 2\}\}, \\ & \{e_{2,1} e_{k,i} e_{k,j}, e_{2,1} e_{j,i} e_{j,k}, e_{2,1} e_{i,k} e_{k,j} \mid i \leq 2 < j < k \leq n \text{ or } 3 \leq i < j < k \leq n\}, \\ & \left\{ \begin{array}{ccc} e_{p,q} e_{k,1} e_{k,2}, & e_{p,q} e_{k,1} e_{1,2}, & e_{p,q} e_{2,1} e_{k,2}, \\ e_{p,q} e_{2,1} e_{2,k}, & e_{p,q} e_{2,1} e_{1,k}, & e_{p,q} e_{k,1} e_{2,k}, \\ e_{p,q} e_{1,2} e_{1,k}, & e_{p,q} e_{1,2} e_{2,k}, & e_{p,q} e_{k,2} e_{1,k} \end{array} \mid 3 \leq k \leq n \right\}. \end{aligned}$$

Define $\pi: E^3 \rightarrow W$ on basis elements as before. Ordering bases appropriately, one can use (2.4) and (2.5) to find a submatrix M of the matrix of $\pi \circ \psi_a: I^2 \rightarrow W$ of the form

$$M = \begin{pmatrix} U & * \\ 0 & M_3 \end{pmatrix},$$

where M_3 is given by (2.6) and U is upper triangular, with diagonal entries $a_{2,1} \neq 0$ and $a_{p,q} \neq 0$. (The choice of U depends on which $a_{p,q} \in \{a_{1,3}, a_{2,3}, a_{3,1}, a_{3,2}\}$ is nonzero.) Hence, the matrix M has maximal rank. It follows that $\psi_a: I^2 \rightarrow E^3$ injects.

Finally, consider the case where $a \notin C_{1,2,3} \subset C$, $a_{2,1} \neq 0$, $a_{p,q} \neq 0$ for some $a_{p,q} \in \{a_{1,3}, a_{2,3}, a_{3,1}, a_{3,2}\}$, and $a_{r,s} \neq 0$ for some $\{r, s\} \not\subset \{1, 2, 3\}$. Since we may assume by Case 1 that $a_{r,s} = 0$ if $\{1, 2\} \cap \{r, s\} = \emptyset$, we have $a_{r,s} \neq 0$ for some r, s with $r \in \{1, 2\}$ and $4 \leq s \leq n$. In this instance, let W be the subspace of E^3 spanned by the union of the sets

$$\begin{aligned} & \{e_{1,2}e_{2,1}e_{p,q}\} \cup \{e_{2,1}e_{i,j}e_{j,i} \mid 1 \leq i < j \leq n, \{i, j\} \neq \{1, 2\}\}, \\ & \{e_{2,1}e_{k,i}e_{k,j}, e_{2,1}e_{j,i}e_{j,k}, e_{2,1}e_{i,k}e_{k,j} \mid i \leq 2 < j < k \leq n \text{ or } 3 \leq i < j < k \leq n\}, \\ & \{e_{r,s}e_{3,1}e_{3,2}, e_{r,s}e_{1,3}e_{3,2}, e_{r,s}e_{2,3}e_{3,1}\}, \\ & \{e_{p,q}e_{k,1}e_{k,2}, e_{p,q}e_{2,k}e_{k,1}, e_{p,q}e_{1,k}e_{k,2} \mid 4 \leq k \leq n\}. \end{aligned}$$

Defining $\pi: E^3 \rightarrow W$ on basis elements as above, a calculation using (2.4) and (2.5) shows that $\pi \circ \psi_a: I^2 \rightarrow W$ is an isomorphism. For instance, ordering bases appropriately, one can check that the matrix M of $\pi \circ \psi_a$ has determinant $\det M = a_{2,1}^{m_{2,1}} a_{p,q}^{m_{p,q}} a_{r,s}^3$, where $m_{p,q} = 3n - 8$ and $m_{2,1} = \binom{n}{2}(n-1) - m_{p,q}$. Hence, ψ_a injects in this final case.

Thus, for any $a \notin C$, the map $\psi_a: I^2 \rightarrow E^3$ injects, and $a \notin R$. This completes the proof of Theorem 1.1. $\square$

Remark 2.3. It follows from Theorem 2.1 that the integral cohomology groups of $P\Sigma_n$ are torsion free, with Betti numbers $b_k(P\Sigma_n) = \text{rank } H^k(P\Sigma_n; \mathbb{Z})$ given by the coefficients of the Poincaré polynomial $p(P\Sigma_n, t) = \sum_{k \geq 0} b_k(P\Sigma_n) \cdot t^k = (1+nt)^{n-1}$; see [8, §6]. Thus the cohomology groups cannot distinguish $P\Sigma_n$ from a direct product $F_n \times \cdots \times F_n$ of $n-1$ free groups of rank n . For $n=2$, this is to be expected, since $P\Sigma_2 \cong F_2$.

For $n \geq 3$, the groups $P\Sigma_n$ and $F_n \times \cdots \times F_n$ are not isomorphic and are, in fact, distinguished by their cohomology rings. By Theorem 1.1, the irreducible components of $R^1(P\Sigma_n, \mathbb{K})$ are two- and three-dimensional. On the other hand, the results of [3] or [12] may be used to show that the irreducible components of the first resonance variety of $H^*(F_n \times \cdots \times F_n; \mathbb{K})$ are all n -dimensional.

ACKNOWLEDGMENTS

We thank Graham Denham and Michael Falk for useful conversations.

REFERENCES

1. R. Bieri, W. D. Neumann, R. Strebel, *A geometric invariant of discrete groups*, Invent. Math. **90** (1987), 451–477. MR0914846
2. A. Brownstein, R. Lee, *Cohomology of the group of motions of n strings in 3-space*, in: *Mapping class groups and moduli spaces of Riemann surfaces* (Goettingen, 1991 / Seattle, WA, 1991), pp. 51–61, Contemp. Math. **150**, Amer. Math. Soc., Providence, RI, 1993. MR1234259

3. D. Cohen, A. Suciu, *Characteristic varieties of arrangements*, Math. Proc. Cambridge Phil. Soc. **127** (1999), 33–53. MR1692519
4. D. Dahm, *A generalization of a braid theory*, Ph.D. Thesis, Princeton Univ., 1962.
5. A. Dimca, S. Papadima, A. Suciu, *Topology and geometry of cohomology jumping loci*, Duke Math. J., to appear. arXiv:0902.1250
6. M. Falk, *Arrangements and cohomology*, Ann. Combin. **1** (1997), 135–157. MR1629681
7. D. Goldsmith, *The theory of motion groups*, Michigan Math. J. **28** (1981), 3–17. MR0600411
8. C. Jensen, J. McCammond, J. Meier, *The integral cohomology of the group of loops*, Geom. Topol. **10** (2006), 759–784. MR2240905
9. J. McCool, *On basis-conjugating automorphisms of free groups*, Canad. J. Math. **38** (1986), 1525–1529. MR0873421
10. J. Meier, L. VanWyk, *The Bieri-Neumann-Strebel invariants for graph groups*, Proc. London Math. Soc. (3) **71** (1995), 263–280. MR1337468
11. L. Orlandi-Korner, *The Bieri-Neumann-Strebel invariant for basis-conjugating automorphisms of free groups*, Proc. Amer. Math. Soc. **128** (2000), 1257–1262. MR1712889
12. S. Papadima, A. Suciu, *Algebraic invariants for right-angled Artin groups*, Math. Ann. **334** (2006), 533–555. MR2207874
13. S. Papadima, A. Suciu, *Bieri-Neumann-Strebel-Renz invariants and homology jumping loci*, preprint, 2008. arXiv:0812.2660
14. S. Yuzvinsky, *Orlik-Solomon algebras in algebra and topology*, Russian Math. Surveys **56** (2001), 293–364. MR1859708

DEPARTMENT OF MATHEMATICS, LOUISIANA STATE UNIVERSITY, BATON ROUGE, LOUISIANA 70803

E-mail address: `cohen@math.lsu.edu`

URL: `http://www.math.lsu.edu/~cohen/`