COMMUTING TOEPLITZ OPERATORS ON THE HARDY SPACE OF THE POLYDISK

YOUNG JOO LEE

(Communicated by Marius Junge)

ABSTRACT. We study the commuting problem for Toeplitz operators on the Hardy space of the polydisk. We consider two symbols where one is an arbitrarily bounded symbol and the other is a bounded symbol which has the form of a sum of an analytic function and a co-analytic function. We then give a necessary and sufficient condition on the symbols for which the corresponding Toeplitz operators commute each other.

1. INTRODUCTION

Let T be the boundary of the unit disk D in the complex plane $\mathbb{C}$. The polydisk D^n and torus T^n are the cartesian products of n copies of D and T , respectively. We let $L^p(T^n) = L^p(T^n, \sigma)$ denote the usual Lebesgue space on T^n where $\sigma = \sigma_n$ is the normalized Haar measure on T^n . The Hardy space $H^2(D^n)$ is the closure of the polynomials in $L^2(T^n)$. As is well known, we can identify functions in $H^2(D^n)$ with its holomorphic extension to D^n via the Poisson extension. Thus, we will use the same notation for a function $f \in H^2(D^n)$ and its holomorphic extension f on D^n . Let P denote the orthogonal projection from $L^2(T^n)$ onto $H^2(D^n)$. For a function $u \in L^\infty(T^n)$, the *Toeplitz operator* T_u with symbol u is defined by

$$T_u f = P(uf)$$

for functions $f \in H^2(D^n)$. Then, clearly, T_u is a bounded linear operator on $H^2(D^n)$.

In this paper, we study the commuting problem of when two Toeplitz operators commute each other. On the Hardy space of the unit disk, Brown and Halmos [2] first obtained a complete description of bounded symbols of commuting Toeplitz operators by asserting that two Toeplitz operators with bounded symbols commute on $H^2(D)$ if and only if both symbols are all analytic or both symbols are all co-analytic, or a nontrivial linear combination of the symbols is constant. Here and in what follows, a function in $L^1(T^n)$ is said to be *analytic* if all its Fourier coefficients vanish outside $\mathbb{Z}_+^n$, where $\mathbb{Z}_+$ is the set of all nonnegative integers. Also, the complex conjugate of an analytic function is called *co-analytic*. In [10], Stroethoff gave a new proof to prove the result of Brown and Halmos as well as a generalization.

Received by the editors February 24, 2009.

2000 *Mathematics Subject Classification.* Primary 47B35; Secondary 32A35.

Key words and phrases. Toeplitz operators, Hardy space, polydisk.

©2009 American Mathematical Society
 Reverts to public domain 28 years from publication

Also, Zheng [11] extended the result of Brown and Halmos to the Hardy space of the ball with bounded pluriharmonic symbols.

Also, the same commuting problem has been studied on the polydisk. In a recent paper, the author [8] studied the same problem on the bidisk when one symbol is a general bounded symbol and the other is a certain type of bounded symbol. On the general polydisk, Choe, Koo and the author [3] characterized two pluriharmonic symbols of commuting Toeplitz operators in the course of studying the corresponding problem on the Bergman space. The result in [3] shows that for given functions $f, g, h, k \in H^2(D^n)$, $T_{f+\bar{g}}$ and $T_{h+\bar{k}}$ commute on $H^2(D^n)$ if and only if $f\bar{k} - h\bar{g}$ is n -harmonic on D^n . Recall that a function $u \in C^2(D^n)$ is called n -harmonic, as in [9], if u is harmonic in each variable separately. More explicitly, u is n -harmonic if

$$\partial_j \bar{\partial}_j u = 0, \quad j = 1, 2, \dots, n,$$

where $\partial_j = \partial/\partial z_j$.

In this paper, we continue to study the same commuting problem on the polydisk for Toeplitz operators with two symbols where one is an arbitrarily bounded symbol and the other is a bounded symbol which has the form of $f + \bar{g}$ where f and g are analytic on T^n . We then give a characterization for which the corresponding Toeplitz operators commute.

The following is the our main result.

Main Theorem. *Let $u, v \in L^\infty(T^n)$. Suppose $v = f + \bar{g}$ where f, g are analytic on T^n . Then the following statements are equivalent:*

- (a) $T_u T_v = T_v T_u$ on $H^2(D^n)$.
- (b) $\hat{u}[f - \bar{g}]$ is n -harmonic on D^n .

Here $\hat{u}$ is the Poisson integral of u .

We were not able to obtain a characterization when two symbols are all arbitrarily bounded symbols. Our result generalizes the results in [2] and [3] mentioned above. To prove the Main Theorem, we use an entirely different argument from those of [2] and [3]. In the next section, we collect some basic facts. In Section 3, we prove the Main Theorem and recover the known results as immediate consequences.

2. PRELIMINARIES

Recall that we can identify functions in $H^2(D^n)$ with its holomorphic extension on D^n via the Poisson extension. With this identification, the point evaluation at each point in D^n is a bounded linear functional on $H^2(D^n)$. Hence, for each $a \in D^n$, there exists a unique kernel function $K_a \in H^2(D^n)$ which has reproducing property

$$(1) \quad f(a) = \langle f, K_a \rangle$$

for functions $f \in H^2(D^n)$. Here and in what follows, the notation $\langle \cdot, \cdot \rangle$ denotes the inner product in $L^2(T^n)$ with respect to the measure σ . For $a = (a_1, \dots, a_n) \in D^n$, it is well known that K_a is the Cauchy kernel whose the explicit formula is given by

$$K_a(\zeta) = \prod_{j=1}^n \frac{1}{(1 - \bar{a}_j \zeta_j)}, \quad \zeta = (\zeta_1, \dots, \zeta_n) \in T^n.$$

Thus, by (1), we can write the projection P as the integral operator

$$P\varphi(a) = \langle P\varphi, K_a \rangle = \langle \varphi, K_a \rangle = \int_{T^n} \frac{\varphi(\zeta)}{\prod_{j=1}^n (1 - a_j \bar{\zeta}_j)} d\sigma(\zeta)$$

for $\varphi \in L^2(T^n)$ and $a = (a_1, \dots, a_n) \in T^n$. The above integral formula for P shows that the domain of P can be extended to $L^1(T^n)$. Moreover, it is known that $Pf = f$ for analytic functions $f \in L^1(T^n)$. Also, since P is the projection from $L^2(T^n)$ onto $H^2(D^n)$, we have

$$P[\bar{f}P(\varphi)](a) = \int_{T^n} P(\varphi) \overline{fK_a} d\sigma = \int_{T^n} \varphi \overline{fK_a} d\sigma = P(\bar{f}\varphi)(a)$$

for points $a \in D^n$ and hence

$$(2) \quad P[\bar{f}P(\varphi)] = P(\bar{f}\varphi)$$

for functions $f \in H^2(D^n)$ and $\varphi \in L^\infty(T^n)$. Also, note that

$$P(\bar{f}K_a)(z) = \langle K_a, fK_z \rangle = \overline{f(a)K_z(a)} = \overline{f(a)}K_a(z), \quad z \in D^n,$$

and then we have

$$(3) \quad P(\bar{f}K_a) = \overline{f(a)}K_a$$

for functions $f \in H^2(D^n)$ and points $a \in D^n$. In particular, for $n = 1$, we have

$$(4) \quad \int_T \frac{\overline{\phi}(\xi)}{(1 - \lambda\bar{\xi})(1 - \bar{\mu}\xi)} d\sigma_1(\xi) = \frac{\overline{\phi(\mu)}}{(1 - \bar{\mu}\lambda)}$$

for $\phi \in H^2(D)$ and points $\lambda, \mu \in D$.

For $a = (a_1, \dots, a_n) \in D^n$, we let k_a be the normalized kernel given by

$$k_a(\zeta) = \prod_{j=1}^n \frac{\sqrt{1 - |a_j|^2}}{(1 - \bar{a}_j \zeta_j)}$$

for $\zeta = (\zeta_1, \dots, \zeta_n) \in T^n$. Since $|k_a|^2$ is the Poisson kernel for D^n , the Poisson integral $\widehat{\varphi}$ of $\varphi \in L^1(T^n)$ can be written as

$$\widehat{\varphi}(a) = \int_{T^n} \varphi |k_a|^2 d\sigma, \quad a \in D^n.$$

As is well known, each Poisson integral is n -harmonic on D^n . Given a function ψ on D^n , the radial limit ψ^* is defined by $\psi^*(\zeta) = \lim_{r \rightarrow 1} \psi(r\zeta)$ at every $\zeta \in T^n$ at which the limit exists. For $\varphi \in L^1(T^n)$, it is known that $[\widehat{\varphi}]^* = \varphi$ a.e on T^n , and we note that

$$(5) \quad \widehat{\varphi}(a) = \langle \varphi k_a, k_a \rangle = \frac{1}{K_a(a)} P(\varphi K_a)(a)$$

for points $a \in D^n$. See Chapter 2 of [9] for related facts and details.

3. THE PROOF OF THE MAIN THEOREM

In this section, we prove the Main Theorem and then derive the known results as immediate consequences. Before doing this, we need a couple of preliminary lemmas. We first need the following characterization of harmonicity of functions which is a finite sum of products of holomorphic and antiholomorphic functions on the unit disk D .

Lemma 1. Let $f_1, \dots, f_N$ and $k_1, \dots, k_N$ be holomorphic functions on D for which $\sum_{j=1}^N f_j \overline{k_j}$ is harmonic on D . Then we have

$$\sum_{j=1}^N f_j \overline{k_j} = \sum_{j=1}^N \left[f_j \overline{k_j(0)} + f_j(0) \overline{k_j} - f_j(0) \overline{k_j(0)} \right].$$

Proof. See Lemma 2 of [8]. □

We also need the following lemma which is taken from Lemma 9 of [4].

Lemma 2. Let $f_1, \dots, f_N$ and $k_1, \dots, k_N$ be finitely many holomorphic functions on D . If $\sum_{j=1}^N f_j \overline{k_j} = 0$ on D , then

$$\sum_{j=1}^N f_j(\lambda) \overline{k_j(\mu)} = 0$$

for every $\lambda, \mu \in D$.

We are now ready to prove the Main Theorem.

Proof of the Main Theorem. Since $v \in L^\infty(T^n)$, we note that $f, g \in H^2(D^n)$. Suppose (a) and let $a \in D^n$. Since $T_u T_v = T_v T_u$, we have in particular,

$$(6) \quad \langle T_v T_u k_a, k_a \rangle = \langle T_u T_v k_a, k_a \rangle.$$

By (2), we first note that $P[\overline{g}P(uk_a)] = P(\overline{g}uk_a)$ and hence

$$(7) \quad T_v T_u k_a = P[(f + \overline{g})P(uk_a)] = fP(uk_a) + P(\overline{g}uk_a).$$

Since $f \in H^2(D^n)$, we note by (5)

$$\langle fP(uk_a), k_a \rangle = \frac{1}{K_a(a)} f(a) P(uk_a)(a) = f(a) \widehat{u}(a)$$

and $\langle P[\overline{g}uk_a], k_a \rangle = \widehat{[\overline{g}u]}(a)$. It follows that

$$(8) \quad \begin{aligned} \langle T_v T_u k_a, k_a \rangle &= \langle fP(uk_a), k_a \rangle + \langle P(\overline{g}uk_a), k_a \rangle \\ &= f(a) \widehat{u}(a) + \widehat{[\overline{g}u]}(a). \end{aligned}$$

On the other hand, by (3), we have that $T_{f+\overline{g}} k_a = [f + \overline{g(a)}] k_a$ and hence

$$(9) \quad T_u T_v k_a = T_u T_{f+\overline{g}} k_a = P(ufk_a) + \overline{g(a)} P(uk_a).$$

It follows from (5) that

$$\langle T_u T_v k_a, k_a \rangle = \widehat{[uf]}(a) + \overline{g(a)} \widehat{u}(a).$$

Combining the above with (8) and together with (6), we have

$$\widehat{u}[f - \overline{g}] = \widehat{[uf - \overline{g}u]}$$

on D^n . Since $\widehat{[uf - \overline{g}u]}$ is n -harmonic, we see that the function $\widehat{u}[f - \overline{g}]$ is also n -harmonic on D^n . Thus we have (b).

Now suppose (b) and prove (a). In the proof, we will use the usual multi-index notations. The notation $\mathbb{Z}$ denotes the set of all integers. For $\zeta = (\zeta_1, \dots, \zeta_n) \in T^n$ and $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{Z}^n$, we write $\zeta^\alpha = \zeta_1^{\alpha_1} \zeta_2^{\alpha_2} \dots \zeta_n^{\alpha_n}$ as usual. Also, let $J = \{0, 1\}$.

Let us consider the multiple Fourier series for u given by

$$u(\zeta) = \sum_{\alpha \in \mathbb{Z}^n} \hat{u}(\alpha) \zeta^\alpha, \quad \zeta = (\zeta_1, \dots, \zeta_n) \in T^n,$$

where $\hat{u}(\alpha)$ are Fourier coefficients of u . Given $I = (i_1, \dots, i_n)$, where each $i_k \in J$, we let $[I]$ be the set of all $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{Z}^n$ for which $\alpha_k \leq 0$ if $i_k = 0$, and $\alpha_k > 0$ if $i_k = 1$ for each k . Put

$$u_I(\zeta) = \sum_{\alpha \in [I]} \hat{u}(\alpha) \zeta^\alpha.$$

Note that for each $k = 1, \dots, n$, if $i_k = 1$ (resp. $i_k = 0$), then the function $u_I(\zeta)$ is analytic (resp. co-analytic) in the variable ζ_k . For each $\alpha = (\alpha_1, \dots, \alpha_n) \in [I]$ and $z = (z_1, \dots, z_n) \in D^n$, we define

$$\tilde{z}_k^{\alpha_k} = \begin{cases} \overline{z_k^{-\alpha_k}} & \text{if } i_k = 0, \\ z_k^{\alpha_k} & \text{if } i_k = 1, \end{cases} \quad k = 1, \dots, n,$$

and let $\tilde{z}^\alpha = \tilde{z}_1^{\alpha_1} \dots \tilde{z}_n^{\alpha_n}$. Define a function $\mathbf{U}_I$ on D^n by

$$\mathbf{U}_I(z) = \sum_{\alpha \in [I]} \hat{u}(\alpha) \tilde{z}^\alpha, \quad z = (z_1, \dots, z_n) \in D^n.$$

Then, for each k , if $i_k = 1$ (resp. $i_k = 0$), the function $\mathbf{U}_I(z)$ is holomorphic (resp. antiholomorphic) in the variable z_k . Also, by (4) and an application of Corollary 2.3 of [6], we can easily verify

$$(10) \quad P(u_I K_a)(z) = \mathbf{U}_I\left((1 - i_1)a_1 + i_1 z_1, \dots, (1 - i_n)a_n + i_n z_n\right) K_a(z)$$

for $a = (a_1, \dots, a_n) \in D^n$. Thus, by (5) we have $\widehat{u_I} = \mathbf{U}_I$. Since the multiple Fourier series for u can be rewritten as

$$(11) \quad u = \sum_{i_1, \dots, i_n \in J} u_{i_1, \dots, i_n},$$

we have

$$\widehat{u} = \sum_{i_1, \dots, i_n \in J} \mathbf{U}_{i_1, \dots, i_n}.$$

Put $\Psi = \widehat{u}[f - \bar{g}]$. Since Ψ is n -harmonic on D^n by the assumption, we see that

$$\Psi = f \sum_{i_1, \dots, i_n \in J} \mathbf{U}_{i_1, \dots, i_n} - \bar{g} \sum_{i_1, \dots, i_n \in J} \mathbf{U}_{i_1, \dots, i_n}$$

is n -harmonic on D^n . For each $k = 1, \dots, n$, we recall that if $i_k = 1$ (resp. $i_k = 0$), then the function $\mathbf{U}_{i_1, \dots, i_n}(z)$ is holomorphic (resp. antiholomorphic) in the variable z_k . Thus, the function Ψ is a finite sum of products of a holomorphic function and an antiholomorphic function in each variable separately. Also, since Ψ is n -harmonic on D^n , Ψ is harmonic in each variable separately. Given $i, j \in J$, we put

$$\vartheta(i, j) = \begin{cases} 1 & \text{if } (i, j) = (1, 0), (0, 1) \text{ or } (1, 1), \\ -1 & \text{if } (i, j) = (0, 0). \end{cases}$$

Now, applying Lemma 1 to Ψ with respect to the first variable, we see that

$$\begin{aligned}
\Psi(a) &= f(a) \sum_{i_2, \dots, i_n \in J} \mathbf{U}_{1, i_2, \dots, i_n}(a) - \overline{g(a)} \sum_{i_2, \dots, i_n \in J} \mathbf{U}_{0, i_2, \dots, i_n}(a) \\
&\quad + f(a) \sum_{i_2, \dots, i_n \in J} \mathbf{U}_{0, i_2, \dots, i_n}(a) - \overline{g(a)} \sum_{i_2, \dots, i_n \in J} \mathbf{U}_{1, i_2, \dots, i_n}(a) \\
&= f(a) \sum_{i_2, \dots, i_n \in J} \mathbf{U}_{1, i_2, \dots, i_n}(a) - \overline{g(a)} \sum_{i_2, \dots, i_n \in J} \mathbf{U}_{0, i_2, \dots, i_n}(a) \\
&\quad + \sum_{\substack{l_1, m_1 \in J \\ l_1 m_1 = 0}} \vartheta(l_1, m_1) f(l_1 a_1, a') \sum_{i_2, \dots, i_n \in J} \mathbf{U}_{0, i_2, \dots, i_n}(m_1 a_1, a') \\
&\quad - \sum_{\substack{l_1, m_1 \in J \\ l_1 m_1 = 0}} \vartheta(l_1, m_1) \overline{g(m_1 a_1, a')} \sum_{i_2, \dots, i_n \in J} \mathbf{U}_{1, i_2, \dots, i_n}(l_1 a_1, a') \\
&= \sum_{l_1, m_1 \in J} \vartheta(l_1, m_1) f(l_1 a_1, a') \sum_{i_2, \dots, i_n \in J} \mathbf{U}_{l_1 m_1, i_2, \dots, i_n}(m_1 a_1, a') \\
&\quad - \sum_{l_1, m_1 \in J} \vartheta(l_1, m_1) \overline{g(m_1 a_1, a')} \sum_{i_2, \dots, i_n \in J} \mathbf{U}_{(1-l_1 m_1), i_2, \dots, i_n}(l_1 a_1, a'),
\end{aligned}$$

where $a' = (a_2, \dots, a_n)$ and $a = (a_1, a') \in D^n$. Repeating the same argument as above with respect to variables $a_2, \dots, a_n$ in order, we see that the n -harmonicity of Ψ implies that Ψ admits the following decomposition:

(12)

$$\Psi(a) = \sum_{l_k, m_k \in J} \vartheta_{l, m} \left[f(l \cdot a) \mathbf{U}_{l_1 m_1, \dots, l_n m_n}(m \cdot a) - \overline{g(m \cdot a)} \mathbf{U}_{1-l_1 m_1, \dots, 1-l_n m_n}(l \cdot a) \right],$$

where we use the notation $l \cdot a = (l_1 a_1, \dots, l_n a_n)$, $m \cdot a = (m_1 a_1, \dots, m_n a_n)$ and $\vartheta_{l, m} = \vartheta(l_1, m_1) \cdots \vartheta(l_n, m_n)$ for simplicity. Recall that

$$\Psi = [f - \overline{g}] \sum_{i_1, \dots, i_n \in J} \mathbf{U}_{i_1, \dots, i_n}.$$

Note that Ψ is a finite sum of products of a holomorphic function and an antiholomorphic function in each variable separately. Also, we see that the function on the right-hand side of (12) is also a finite sum of products of a holomorphic function and an antiholomorphic function in each variable separately. Hence, by (12) and an application of Lemma 2, we have

$$\begin{aligned}
&\left[f(z) - \overline{g(a)} \right] \sum_{i_k \in J} \mathbf{U}_{i_1, \dots, i_n} \left((1-i_1)a_1 + i_1 z_1, \dots, (1-i_n)a_n + i_n z_n \right) \\
(13) \quad &= \sum_{l_k, m_k \in J} \vartheta_{l, m} \left[f(lz) \mathbf{U}_{l_1 m_1, \dots, l_n m_n}(\mathbf{P}) - \overline{g(ma)} \mathbf{U}_{1-l_1 m_1, \dots, 1-l_n m_n}(\mathbf{Q}) \right]
\end{aligned}$$

for all $a, z \in D^n$. Here we use the notation

$$\begin{aligned}
\mathbf{P} &= \left([(1-l_1 m_1)a_1 + (l_1 m_1)z_1]m_1, \dots, [(1-l_n m_n)a_n + (l_n m_n)z_n]m_n \right), \\
\mathbf{Q} &= \left([(1-l_1 m_1)z_1 + (l_1 m_1)a_1]l_1, \dots, [(1-l_n m_n)z_n + (l_n m_n)a_n]l_n \right)
\end{aligned}$$

for simplicity.

Note that the set $\{K_a : a \in D^n\}$ spans a dense subset of $H^2(D^n)$. So, in order to prove (a), it is sufficient to show that $[T_u T_v - T_v T_u]K_a = 0$ for all $a \in D^n$. Using (7) and (9), we see that $T_u T_v = T_v T_u$ is equivalent to

$$(14) \quad [f(z) - \overline{g(a)}]P(uK_a)(z) = P[(f - \bar{g})uK_a](z)$$

for all $a, z \in D^n$. Let $a = (a_1, \dots, a_n)$, $z = (z_1, \dots, z_n) \in D^n$ be arbitrary points. By (11) and (10), we first have

$$P(uK_a)(z) = K_a(z) \sum_{i_1, \dots, i_n \in J} \mathbf{U}_{i_1, \dots, i_n}(\mathbf{R})$$

and then

$$(15) \quad [f(z) - \overline{g(a)}]P(uK_a)(z) = [f(z) - \overline{g(a)}]K_a(z) \sum_{i_1, \dots, i_n \in J} \mathbf{U}_{i_1, \dots, i_n}(\mathbf{R})$$

for all $a, z \in D^n$. Here, $\mathbf{R} = ((1-i_1)a_1 + i_1 z_1, \dots, (1-i_n)a_n + i_n z_n)$ for simplicity.

On the other hand, by applications of (4) and Corollary 2.3 of [6] again, we can verify that

$$P[f(l \cdot)u_{l_1 m_1, \dots, l_n m_n}(m \cdot)K_a](z) = f(l \cdot z)\mathbf{U}_{l_1 m_1, \dots, l_n m_n}(\mathbf{P})K_a(z)$$

and

$$P[\bar{g}(m \cdot)u_{1-l_1 m_1, \dots, 1-l_n m_n}(l \cdot)K_a](z) = \bar{g}(m \cdot a)\mathbf{U}_{1-l_1 m_1, \dots, 1-l_n m_n}(\mathbf{Q})K_a(z)$$

for each $l = (l_1, \dots, l_n)$ and $m = (m_1, \dots, m_n)$, where $l_k, m_k \in J$. Here $l \cdot z = (l_1 z_1, \dots, l_n z_n)$ and $m \cdot a = (m_1 a_1, \dots, m_n a_n)$. Also, given functions ψ on T^n and $i = (i_1, \dots, i_n)$ with each $i_k \in J$, we use the notation $\psi(i \cdot)$ to denote the function on T^n defined by $\psi(i \cdot)(\zeta) = \psi(i_1 \zeta_1, \dots, i_n \zeta_n)$ for $\zeta = (\zeta_1, \dots, \zeta_n) \in T^n$.

Now, since $\Psi = [f - \bar{g}]u$, the radial limit Ψ^* of Ψ is equal to $[f - \bar{g}]u$. Also, since $\mathbf{U}_{i_1, \dots, i_n} = \widehat{u_{i_1, \dots, i_n}}$, we have that $\mathbf{U}_{i_1, \dots, i_n}^* = u_{i_1, \dots, i_n}$ for each $i_1, \dots, i_n \in J$. It follows from (12) and observations above that

$$\begin{aligned} & P[(f - \bar{g})uK_a](z) \\ &= P[\Psi^* K_a](z) \\ &= \sum_{l_k, m_k \in J} \vartheta_{l, m} P\left([f(l \cdot)u_{l_1 m_1, \dots, l_n m_n}(m \cdot) - \bar{g}(m \cdot)u_{1-l_1 m_1, \dots, 1-l_n m_n}(l \cdot)]K_a\right)(z) \\ &= \sum_{l_k, m_k \in J} \vartheta_{l, m} [f(lz)\mathbf{U}_{l_1 m_1, \dots, l_n m_n}(\mathbf{P}) - \bar{g}(ma)\mathbf{U}_{1-l_1 m_1, \dots, 1-l_n m_n}(\mathbf{Q})] K_a(z). \end{aligned}$$

Now, combining the above with (15) and together with (13), we have (14). The proof is complete. $\square$

We now close the paper with some immediate consequences of the Main Theorem. First, if one of the symbols of commuting Toeplitz operators is analytic, then we have the following result, which is a special case of Theorem 3.2 of [5] or Corollary 1 of [7].

Corollary 3. *Let $u, v \in L^\infty(T^n)$. Suppose v is analytic. Then $T_u T_v = T_v T_u$ on $H^2(D^n)$ if and only if for each $k = 1, \dots, n$, $v(\zeta)$ is constant or $u(\zeta)$ is analytic in variable ζ_k .*

Proof. By the Main Theorem, we have $T_u T_v = T_v T_u$ if and only if $v\hat{u}$ is n -harmonic on D^n . Using the same notation as in the proof of the Main Theorem, we have

$$\hat{u} = \sum_{i_1, \dots, i_n \in J} \mathbf{U}_{i_1, \dots, i_n}.$$

Note that the n -harmonicity of $v\hat{u}$ is equivalent to that for each $k = 1, \dots, n$,

$$\partial_k v(z) \sum_{i_m \neq i_k} \bar{\partial}_k \mathbf{U}_{i_1, \dots, i_{k-1}, 0, i_{k+1}, \dots, i_n}(z) = 0, \quad z \in D^n,$$

which is equivalent to either v being constant or each $\mathbf{U}_{i_1, \dots, i_{k-1}, 0, i_{k+1}, \dots, i_n}$ being constant in variable z_k . Thus, n -harmonicity of $v\hat{u}$ is equivalent to that for each k , $v(z)$ is constant or $\hat{u}(z)$ is holomorphic in variable z_k , which in turn is equivalent to $v(\zeta)$ being constant or $u(\zeta)$ being analytic in variable ζ_k . The proof is complete. $\square$

As another simple application, we recover the results in [2] and [3] mentioned before.

Corollary 4. *Let $u, v \in L^\infty(T^n)$. Suppose $v = f + \bar{g}$ and $u = h + \bar{k}$, where f, g, h, k are analytic functions on T^n . Then $T_u T_v = T_v T_u$ on $H^2(D^n)$ if and only if $f\bar{k} - h\bar{g}$ is n -harmonic on D^n .*

Proof. Note that $\hat{u} = h + \bar{k}$. Hence, by the Main Theorem, we have that T_u and T_v commute if and only if $[f - \bar{g}][h + \bar{k}]$ is n -harmonic, which is equivalent to the n -harmonicity of $f\bar{k} - h\bar{g}$. The proof is complete. $\square$

Given functions f, g, h, k holomorphic on the unit disk D , we let $v = f + \bar{g}$ and $u = h + \bar{k}$. Then one can verify that $f\bar{k} - h\bar{g}$ is harmonic on D if and only if u, v are all holomorphic, u, v are all antiholomorphic, or a nontrivial linear combination of u, v is constant. For example, see the proof of Theorem 1 of [1]. Also, every $u \in L^2(T)$ can be decomposed as $u = f + \bar{g}$ for some $f, g \in H^2(D)$. Combining these facts with Corollary 4, we can recover the result of Brown and Halmos in [2] on the unit disk.

Corollary 5. *Let $u, v \in L^\infty(T)$. Then $T_u T_v = T_v T_u$ on $H^2(D)$ if and only if u and v are all analytic, u and v are all co-analytic, or a nontrivial linear combination of u and v is constant on T .*

REFERENCES

- [1] S. Axler and Ž. Čučković, *Commuting Toeplitz operators with harmonic symbols*, Integr. Equ. Oper. Theory 14(1991), 1–11. MR1079815 (92f:47018)
- [2] A. Brown and P. Halmos, *Algebraic properties of Toeplitz operators*, J. Reine Angew. Math. 213(1964), 89–102. MR0160136 (28:3350)
- [3] B. R. Choe, H. Koo and Y. J. Lee, *Commuting Toeplitz operators on the polydisk*, Trans. Amer. Math. Soc. 356(2004), 1727–1749. MR2031039 (2004k:47054)
- [4] B. R. Choe and Y. J. Lee, *Pluriharmonic symbols of commuting Toeplitz operators*, Illinois J. Math. 37(1993), 424–436. MR1219648 (94i:47041)
- [5] X. Ding, *Products of Toeplitz operators on the polydisk*, Integr. Equ. Oper. Theory 45(2003), 389–403. MR1971745 (2004b:47051)
- [6] X. Ding, *The finite sum of finite products of Toeplitz operators on the polydisk*, J. Math. Anal. Appl. 320(2006), 464–481. MR2230483 (2007b:47072)
- [7] C. Gu, *Some algebraic properties of Toeplitz and Hankel operators on polydisk*, Arch. Math. 80(2003), 393–405. MR1982839 (2004e:47039)
- [8] Y. J. Lee, *Commuting Toeplitz operators on the Hardy space of the bidisk*, J. Math. Anal. Appl. 341(2008), 738–749. MR2394121 (2009c:47037)

- [9] W. Rudin, *Function theory in polydiscs*, W. A. Benjamin, Inc., 1969. MR0255841 (41:501)
- [10] K. Stroethoff, *Algebraic properties of Toeplitz operators on the Hardy space via the Berezin transform*, Contemp. Math., 232, Amer. Math. Soc., Providence, RI, 1999, 313–319. MR1678343 (2000a:47060)
- [11] D. Zheng, *Commuting Toeplitz operators with pluriharmonic symbols*, Trans. Amer. Math. Soc. 350(1998), 1595–1618. MR1443898 (98i:47027)

DEPARTMENT OF MATHEMATICS, CHONNAM NATIONAL UNIVERSITY, GWANGJU 500-757,
REPUBLIC OF KOREA

E-mail address: leeyj@chonnam.ac.kr