SOLUTION TO FARHADI-GHAHRAMANI'S MULTIPLIER PROBLEM

MATTHIAS NEUFANG

(Communicated by Marius Junge)

ABSTRACT. We answer, in the negative, the main question raised by H. Farhadi and F. Ghahramani in their article *Involutions on the second duals of group algebras and a multiplier problem*, Proc. Edinburgh Math. Soc. (2) **50** (2007), no. 1, 153–161.

Let G be a locally compact group, and consider its group algebra $L_1(G)$ with convolution product. Let $\mathcal{A} = L_1(G)^{**}$, endowed with the first Arens product, which we shall denote by $*$. We write $\cdot$ for the first Arens product on $\mathcal{A}^{**}$, as well as for the canonical left action of $\mathcal{A}^{**}$ on $\mathcal{A}^*$, and of $\mathcal{A}^*$ on $\mathcal{A}$. Recall that these are defined as follows, for $m, n \in \mathcal{A}^{**}$, $f \in \mathcal{A}^*$ and $a, b \in \mathcal{A}$:

$$\begin{aligned}\langle m \cdot n, f \rangle &= \langle m, n \cdot f \rangle, \\ \langle n \cdot f, a \rangle &= \langle n, f \cdot a \rangle, \\ \langle f \cdot a, b \rangle &= \langle f, a * b \rangle.\end{aligned}$$

Analogously, one defines the second Arens product starting from the product $b * a$ (but we shall not need this here).

In [3], noting that for any $N \in L_1(G)^{****}$ the map

$$R_N(f) := N \cdot f \quad (f \in L_1(G)^{***})$$

is a right $L_1(G)$ -module homomorphism on $L_1(G)^{***}$, the authors ask the following question [3, Problem 4.1]:

- (*) *Is every w^* -continuous surjective right $L_1(G)$ -module map*
 $R : L_1(G)^{***} \rightarrow L_1(G)^{***}$ *equal to R_N for some $N \in L_1(G)^{****}$?*

In the present paper we shall show:

Theorem 1. *The answer to the above multiplier problem is negative for all infinite, countable, discrete, abelian groups (e.g., for $G = \mathbb{Z}$).*

To prepare the proof, let us first consider $\mathcal{A} = \ell_1(G)^{**}$ for an arbitrary discrete group G . Denote by $\mathcal{B}_{\mathcal{C},r}(\mathcal{A}^*)$ the algebra of all bounded linear right $\mathcal{C}$ -module maps on $\mathcal{A}^*$, where $\mathcal{C}$ is a (closed) subalgebra of $\mathcal{A}$, and by $\mathcal{B}_{\mathcal{C},r}^c(\mathcal{A}^*)$ the subalgebra of w^* -continuous module maps.

Received by the editors February 11, 2008.

2000 *Mathematics Subject Classification.* Primary 43A20, 43A22.

Key words and phrases. Arens products, group algebras, Stone-Čech compactification, module homomorphisms, topological centres.

The author was partially supported by NSERC. This support is gratefully acknowledged.

Since $\mathcal{A}$ is unital, by [8, Proposition 4.1], the map R_N induces an isometric algebra isomorphism

$$\mathcal{A}^{**} = (\mathcal{A}^* \cdot \mathcal{A})^* \cong \mathcal{B}_{\mathcal{A},r}(\mathcal{A}^*).$$

So we need to find a surjective map

$$(1) \quad \Phi \in \mathcal{B}_{\ell_1(G),r}^\sigma(\ell_1(G)^{***}) \setminus \mathcal{B}_{\ell_1(G)^{**},r}(\ell_1(G)^{***})$$

(which, as we shall see, exists whenever G is infinite, countable, and abelian). Note that, as is easily checked, for any $N \in \ell_1(G)^{****}$ the map R_N is in fact a right $\ell_1(G)^{**}$ -module homomorphism, i.e., belongs to $\mathcal{B}_{\ell_1(G)^{**},r}(\ell_1(G)^{***})$.

For later use we shall give below some background on semigroup compactifications; for a full meal, we refer the reader to [5] and [1]. Recall that the spectrum of the commutative C^* -algebra $\ell_\infty(G)$ is the Stone-Ćech compactification βG , which is a compact right topological semigroup with the first Arens product (inherited from $\ell_1(G)^{**}$). Denoting the closure in βG of a subset S of G by $\overline{S}$, one defines the *growth* or *remainder* of S as

$$S^* := \overline{S} \cap (\beta G \setminus G);$$

then S^* is compact, and G^* is a compact right topological semigroup. An element $s \in \beta G$ is called *left cancellable* if left multiplication by s is injective on βG .

For $m \in \mathcal{A}$, write λ_m for left multiplication by m in $\mathcal{A}$. We note the following.

Lemma 2. *If m is a left cancellable element of βG , then $\lambda_m : \ell_1(G)^{**} \rightarrow \ell_1(G)^{**}$ is an isometry.*

Proof. This can be shown by transferring *verbatim* the argument given in [1, Proposition 4.4] for right multiplication to our situation (the compact right topological semigroup V in [1, Proposition 4.4] is βG in our case). $\square$

We are now ready to derive Theorem 1.

Proof. Let G be an infinite, countable, discrete, abelian group, and recall that we write $\mathcal{A} = \ell_1(G)^{**}$ (endowed with the first Arens product). Then, by [5, Section 8.4], there exists $m \in G^* \subseteq \mathcal{A}$ such that m is left cancellable in βG . In view of Lemma 2, $\Phi := \lambda_m^* \in \mathcal{B}^\sigma(\mathcal{A}^*)$ is surjective. To establish (1), we need to show that

- (i) Φ is a right $\ell_1(G)$ -module map, and
- (ii) Φ is *not* a right $\ell_1(G)^{**}$ -module map.

To see (i), let $H \in \mathcal{A}^*$, $a \in \ell_1(G) \subseteq \mathcal{A}$ and $b \in \mathcal{A}$. Then we have

$$\langle \Phi(H \cdot a), b \rangle = \langle H, a * m * b \rangle.$$

But $a \in \ell_1(G)$, and since G is abelian, $\ell_1(G)$ is contained in the algebraic centre of $\ell_1(G)^{**}$. Hence a commutes with m , and we get

$$\langle \Phi(H \cdot a), b \rangle = \langle H, m * a * b \rangle = \langle \Phi(H) \cdot a, b \rangle,$$

as desired.

As for (ii), assume towards a contradiction that Φ is a right $\mathcal{A}$ -module map. Then we obtain for all $H \in \mathcal{A}^*$ and $a, b \in \mathcal{A}$ that

$$\langle H, a * m * b \rangle = \langle \Phi(H \cdot a), b \rangle = \langle \Phi(H) \cdot a, b \rangle = \langle H, m * a * b \rangle.$$

Thus, we have $a * m * b = m * a * b$ for all $a, b \in \mathcal{A}$, whence (choosing $b = \delta_e$, the unit of $\mathcal{A}$) we see that m lies in the algebraic centre $Z_a(\mathcal{A})$ of $\mathcal{A} = \ell_1(G)^{**}$. But since $\ell_1(G)$ is commutative, $Z_a(\mathcal{A})$ equals the first topological centre $Z_t^1(\mathcal{A})$ of $\mathcal{A}$, i.e., the

set of all elements $n \in \mathcal{A}$ such that λ_n is w^* -continuous on $\mathcal{A}$. By [7, Theorem 1], $Z_t^1(\ell_1(G)^{**}) = \ell_1(G)$. Hence we obtain that $m \in \ell_1(G)$, which contradicts the fact that $m \in G^*$. $\square$

Remark 3. The motivation for question (*), i.e., [3, Problem 4.1], stems from its link to a long-standing open problem raised by Duncan and Hosseiniun [2, p. 323] asking if, for a locally compact group G , the natural involution on $L_1(G)$ extends to an involution on its bidual. Indeed, Farhadi-Ghahramani show in [3, Theorem 4.2] that the latter has a negative answer for all infinite, locally compact groups G satisfying (*). Note that the existence of an extension of the involution forces G to be discrete (cf. [4, Theorem 2] or [3, Proposition 3.1]). Now, on the one hand, our Theorem 1 shows that there are discrete groups which violate (*). On the other hand, [3, Theorem 3.2(a)] answers Duncan-Hosseiniun's question in the negative for all (discrete) groups which admit an infinite, amenable subgroup. It follows from work by Ol'shanskii [9] that there are infinite, countable, discrete groups without infinite, amenable subgroups (cf. [6, Remark 11]). So, the Duncan-Hosseiniun problem remains open for this class of groups, and the approach proposed through [3, Theorem 4.2] may very well prove fruitful since these groups, being of course radically different from the ones appearing in Theorem 1, may satisfy (*).

REFERENCES

1. H.G. Dales, A.T.-M. Lau and D. Strauss, *Banach algebras on semigroups and on their compactifications*, to appear in: *Memoirs of the American Mathematical Society*, <http://www.amsta.leeds.ac.uk/~pmt6hgd/preprints/AMSseconddualsfinal1.pdf>.
2. J. Duncan and S.A.R. Hosseiniun, *The second dual of a Banach algebra*, *Proc. Roy. Soc. Edinburgh Sect. A* **84** (1979), no. 3-4, 309–325. MR559675 (81f:46057)
3. H. Farhadi and F. Ghahramani, *Involutions on the second duals of group algebras and a multiplier problem*, *Proc. Edinburgh Math. Soc. (2)* **50** (2007), no. 1, 153–161. MR2294009 (2008b:43004)
4. M. Grosser, *Algebra involutions on the bidual of a Banach algebra*, *Manuscripta Math.* **48** (1984), no. 1–3, 291–295. MR753735 (85k:46058)
5. N. Hindman and D. Strauss, *Algebra in the Stone-Ćech Compactification. Theory and Applications*, de Gruyter Expositions in Mathematics **27**, Walter de Gruyter & Co., Berlin, 1998. MR1642231 (99j:54001)
6. A.S. Kechris, *On the classification problem for rank 2 torsion-free abelian groups*, *J. London Math. Soc. (2)* **62** (2000), 437–450. MR1783636 (2002h:20078)
7. A.T.M. Lau and V. Losert, *On the second conjugate algebra of $L_1(G)$ of a locally compact group*, *J. London Math. Soc. (2)* **37** (1988), 464–470. MR939122 (89e:43007)
8. M. Neufang, *On a conjecture by Ghahramani-Lau and related problems concerning topological centres*, *J. Funct. Anal.* **224** (2005), no. 1, 217–229. MR2139110 (2006b:46063)
9. A. Yu. Ol'shanskii, *On residualizing homomorphisms and G -subgroups of hyperbolic groups*, *Internat. J. Algebra Comput.* **3** (1993), no. 4, 365–409. MR1250244 (94i:20069)

SCHOOL OF MATHEMATICS AND STATISTICS, CARLETON UNIVERSITY, OTTAWA, ONTARIO, K1S 5B6, CANADA

E-mail address: mneufang@math.carleton.ca