A NOTE ON THE BUCHSBAUM-RIM MULTIPLICITY OF A PARAMETER MODULE

FUTOSHI HAYASAKA AND EERO HYRY

(Communicated by Bernd Ulrich)

ABSTRACT. In this article we prove that the Buchsbaum-Rim multiplicity $e(F/N)$ of a parameter module N in a free module $F = A^r$ is bounded above by the colength $\ell_A(F/N)$. Moreover, we prove that once the equality $\ell_A(F/N) = e(F/N)$ holds true for some parameter module N in F , then the base ring A is Cohen-Macaulay.

1. INTRODUCTION

Let $(A, \mathfrak{m})$ be a Noetherian local ring with the maximal ideal $\mathfrak{m}$ and $d = \dim A > 0$. Let $F = A^r$ be a free module of rank $r > 0$, and let M be a submodule of F such that F/M has finite length and $M \subseteq \mathfrak{m}F$.

In their article [5] from 1964 Buchsbaum and Rim introduced and studied a multiplicity associated to a submodule of finite colength in a free module. This multiplicity, which generalizes the notion of Hilbert–Samuel multiplicity for ideals, is nowadays called the Buchsbaum-Rim multiplicity. In more detail, it first turns out that the function

$$\lambda(n) := \ell_A(\mathcal{S}_n(F)/\mathcal{R}_n(M))$$

is eventually a polynomial of degree $d + r - 1$, where $\mathcal{S}_A(F) = \bigoplus_{n \geq 0} \mathcal{S}_n(F)$ is the symmetric algebra of F and $\mathcal{R}(M) = \bigoplus_{n \geq 0} \mathcal{R}_n(M)$ is the image of the natural homomorphism from $\mathcal{S}_A(M)$ to $\mathcal{S}_A(F)$. The polynomial $P(n)$ corresponding to $\lambda(n)$ can then be written in the form

$$P(n) = \sum_{i=0}^{d+r-1} (-1)^i e_i \binom{n+d+r-2-i}{d+r-1-i}$$

with integer coefficients e_i . The *Buchsbaum-Rim multiplicity of M in F* , denoted by $e(F/M)$, is now defined to be the coefficient e_0 .

Buchsbaum and Rim also introduced in their article the notion of a parameter module (matrix), which generalizes the notion of a parameter ideal (system of parameters). The module N in F is said to be a *parameter module in F* if the following three conditions are satisfied: (i) F/N has finite length, (ii) $N \subseteq \mathfrak{m}F$,

Received by the editors August 17, 2008, and, in revised form, July 14, 2009.

2000 *Mathematics Subject Classification*. Primary 13H15; Secondary 13D25.

Key words and phrases. Buchsbaum-Rim multiplicity, parameter module, Euler-Poincaré characteristic, generalized Koszul complex.

©2009 American Mathematical Society
 Reverts to public domain 28 years from publication

and (iii) $\mu_A(N) = d + r - 1$, where $\mu_A(N)$ is the minimal number of generators of N .

Buchsbaum and Rim utilized in their study the relationship between the Buchsbaum-Rim multiplicity and the Euler-Poincaré characteristic of a certain complex and proved the following:

Theorem 1.1 (Buchsbaum-Rim [5, Corollary 4.5]). *Let $(A, \mathfrak{m})$ be a Noetherian local ring of dimension $d > 0$. Then the following statements are equivalent:*

- (1) *A is a Cohen-Macaulay local ring.*
- (2) *For any rank $r > 0$, the equality $\ell_A(F/N) = e(F/N)$ holds true for every parameter module N in $F = A^r$.*

Then it is natural to ask the following:

- Question 1.2.**
- (1) Does the inequality $\ell_A(F/N) \geq e(F/N)$ hold true for any parameter module N in F ?
 - (2) Does the equality $\ell_A(F/N) = e(F/N)$ for some parameter module N in F imply that the ring A is Cohen-Macaulay?

The purpose of this article is to give a complete answer to Question 1.2. Our results can be summarized as follows:

Theorem 1.3. *Let $(A, \mathfrak{m})$ be a Noetherian local ring of dimension $d > 0$.*

- (1) *For any rank $r > 0$, the two inequalities*

$$\ell_A(F/N) \geq e(F/N) \text{ and } \ell_A(A/I(N)) \geq e(F/N)$$

always hold true for every parameter module N in $F = A^r$, where $I(N)$ is the 0-th Fitting ideal of F/N .

- (2) *The following statements are equivalent:*

- (i) *A is a Cohen-Macaulay local ring.*
- (ii) *For some rank $r > 0$, there exists a parameter module N in $F = A^r$ such that the equality $\ell_A(F/N) = e(F/N)$ holds true.*
- (iii) *For some rank $r > 0$, there exists a parameter module N in $F = A^r$ such that the equality $\ell_A(A/I(N)) = e(F/N)$ holds true.*

When this is the case, the equality $\ell_A(F/N) = \ell_A(A/I(N)) = e(F/N)$ holds true for all parameter modules N in $F = A^r$ of any rank $r > 0$.

Note that the equality $\ell_A(F/N) = \ell_A(A/I(N))$ is known by [1, 2.10].

The proof of our Theorem 1.3 will be completed in section 4. Section 2 is of a preliminary character. In that section we will recall the definition and some basic facts about the generalized Koszul complex. In order to prove Theorem 1.3, we will investigate in section 3 the higher Euler-Poincaré characteristics of the generalized Koszul complex and show that they are non-negative. Finally, in section 4, we will obtain Theorem 1.3 as a corollary of a more general result (Theorem 4.1).

2. PRELIMINARIES

In this section we will recall the definition and some basic facts about the generalized Koszul complex introduced in [3, 8] (for more details, see also [7, Appendix A2.6]).

Let A be a commutative Noetherian ring, and let $n \geq r > 0$ be integers. Let $\mathfrak{a} = (a_{ij})$ be an $r \times n$ matrix over A , and let $I_r(\mathfrak{a})$ denote the ideal generated by

the maximal minors of $\mathbf{a}$. Let F and G be free modules with bases $\{f_1, \dots, f_r\}$ and $\{e_1, \dots, e_n\}$, respectively. Let S be the symmetric algebra of F , and let S_ℓ be the ℓ -th symmetric power of F . Let $\wedge$ be the exterior algebra of G , and let $\wedge^\ell$ be the ℓ -th exterior power of G . Associated with the i -th row $[a_{i1} \cdots a_{in}]$ of $\mathbf{a}$, there is a differentiation homomorphism $\delta_i : \wedge \rightarrow \wedge$ given by

$$\delta_i(f_{j_1} \wedge \cdots \wedge f_{j_p}) = \sum_{k=1}^p (-1)^{k-1} a_{ij_k} f_{j_1} \wedge \cdots \wedge \widehat{f_{j_k}} \wedge \cdots \wedge f_{j_p}.$$

Let $f_i : S \rightarrow S$ and $f_i^{-1} : S \rightarrow S$ denote the multiplication and division maps by f_i , respectively, i.e.,

$$f_i^{-1}(f_1^{\mu_1} \cdots f_i^{\mu_i} \cdots f_r^{\mu_r}) = \begin{cases} f_1^{\mu_1} \cdots f_i^{\mu_i-1} \cdots f_r^{\mu_r} & (\mu_i > 0) \\ 0 & (\mu_i = 0). \end{cases}$$

Then the generalized Koszul complex $K_\bullet(\mathbf{a}; t)$ associated to a matrix $\mathbf{a}$ and an integer t is the complex

$$K_\bullet(\mathbf{a}; t) : \cdots \rightarrow K_{p+1}(\mathbf{a}; t) \xrightarrow{d_{p+1}} K_p(\mathbf{a}; t) \xrightarrow{d_p} K_{p-1}(\mathbf{a}; t) \rightarrow \cdots$$

defined by

$$K_p(\mathbf{a}; t) = \begin{cases} \wedge^{r+p-1} \otimes_A S_{p-t-1} & (p \geq t+1) \\ \wedge^p \otimes_A S_{t-p} & (p \leq t) \end{cases}$$

and

$$d_{p+1} = \begin{cases} \sum_{j=1}^r \delta_j \otimes f_j^{-1} & (p > t) \\ \delta_r \circ \cdots \circ \delta_1 \otimes 1 & (p = t) \\ \sum_{j=1}^r \delta_j \otimes f_j & (p < t). \end{cases}$$

The generalized Koszul complex $K_\bullet(\mathbf{a}; t)$ is a free complex of A -modules. We note that it is of length $n - r + 1$ when $-1 \leq t \leq n - r + 1$. Also recall that $K_\bullet(\mathbf{a}; t)$ coincides with the ordinary Koszul complex for any t in the case $r = 1$, whereas $K_\bullet(\mathbf{a}; 0)$ is the Eagon-Northcott complex and $K_\bullet(\mathbf{a}; 1)$ is the Buchsbaum-Rim complex. Moreover, the generalized Koszul complex has the following important properties (see [8, 10] and [7, Appendix A2.6]):

Proposition 2.1. *Let $\mathbf{a}$ be an $r \times n$ matrix over A with $n \geq r > 0$. Then*

- (1) [8, Theorem 1] *For any $t, p \in \mathbb{Z}$, $I_r(\mathbf{a})H_p(K_\bullet(\mathbf{a}; t)) = (0)$.*
- (2) [7, Theorem A2.10] *If the grade of $I_r(\mathbf{a})$ is at least $n - r + 1$, then $K_\bullet(\mathbf{a}; t)$ is acyclic for all $-1 \leq t \leq n - r + 1$. Furthermore, if $\mathbf{a}$ is a generic matrix, then $K_\bullet(\mathbf{a}; t)$ is acyclic for all $t \geq -1$.*

3. HIGHER EULER-POINCARÉ CHARACTERISTICS

In this section we will investigate higher Euler-Poincaré characteristics of a generalized Koszul complex.

Throughout this section, let $(A, \mathfrak{m})$ be a Noetherian local ring of dimension $d > 0$. Let $F = A^r$ be a free module of rank $r > 0$ with a basis $\{f_1, \dots, f_r\}$. Let M be a submodule of F generated by $c_1, c_2, \dots, c_n$, where $n = \mu_A(M)$ is the minimal number of generators of M . Writing $c_j = c_{1j}f_1 + \cdots + c_{rj}f_r$ for some $c_{ij} \in A$, we have an $r \times n$ matrix (c_{ij}) associated to M . We call this matrix the matrix of M and denote it by $\widetilde{M}$. Let $I(M) = \text{Fitt}_0(F/M)$ be the 0-th Fitting ideal of F/M . We assume that F/M has finite length and $M \subseteq \mathfrak{m}F$. Then $I(M)$ is an $\mathfrak{m}$ -primary ideal, because $\sqrt{I(M)} = \sqrt{\text{Ann}_A(F/M)}$. Hence each homology

module $H_p(K_\bullet(\widetilde{M}; t))$ has finite length by Proposition 2.1(1). So the Euler-Poincaré characteristics of $K_\bullet(\widetilde{M}; t)$ can be defined as follows:

Definition 3.1. For any integer $q \geq 0$, we set

$$\chi_q(K_\bullet(\widetilde{M}; t)) := \sum_{p \geq q} (-1)^{p-q} \ell_A(H_p(K_\bullet(\widetilde{M}; t)))$$

and call it the q -th partial Euler-Poincaré characteristic of $K_\bullet(\widetilde{M}; t)$. When $q = 0$, we simply write $\chi(K_\bullet(\widetilde{M}; t))$ for $\chi_0(K_\bullet(\widetilde{M}; t))$ and call it the Euler-Poincaré characteristic of $K_\bullet(\widetilde{M}; t)$.

Buchsbaum and Rim studied in [5] the Euler-Poincaré characteristic of the Buchsbaum-Rim complex in analogy with the Euler-Poincaré characteristic of the ordinary Koszul complex in the case of usual multiplicities. In 1985 Kirby investigated in [9] Euler-Poincaré characteristics of the complex $K_\bullet(\widetilde{M}; t)$ for all t and proved the following:

Theorem 3.2 (Buchsbaum-Rim, Kirby). *For any integer $t \in \mathbb{Z}$, we have*

$$\chi(K_\bullet(\widetilde{M}; t)) = \begin{cases} e(F/M) & (n = d + r - 1), \\ 0 & (n > d + r - 1), \end{cases}$$

where $n = \mu_A(M)$ is the minimal number of generators of M . In particular, $\chi(K_\bullet(\widetilde{M}; t)) \geq 0$ for all $t \in \mathbb{Z}$.

The last statement holds for the higher Euler-Poincaré characteristics, too:

Theorem 3.3. *For any $q \geq 0$ and any $t \geq -1$, we have*

$$\chi_q(K_\bullet(\widetilde{M}; t)) \geq 0.$$

Proof. We use ideas from [6]. Let $\widetilde{M} = (c_{ij}) \in \text{Mat}_{r \times n}(A)$ be the matrix of M , and let $X = (X_{ij})$ be the generic matrix of the same size $r \times n$. Let $A[X] = A[X_{ij} \mid 1 \leq i \leq r, 1 \leq j \leq n]$ be a polynomial ring over A , and let $B = A[X]_{(\mathfrak{m}, X)}$. We will consider the ring A as a B -algebra via the substitution homomorphism $\phi : B \rightarrow A ; X_{ij} \mapsto c_{ij}$. Let

$$\mathfrak{b} = \text{Ker } \phi = (X_{ij} - c_{ij} \mid 1 \leq i \leq r, 1 \leq j \leq n)B.$$

We note here that $K_\bullet(X; t) \otimes_B A \cong K_\bullet(\widetilde{M}; t)$, because the generalized Koszul complex is compatible with the base change. Let $C_t(X) := H_0(K_\bullet(X; t))$. By Proposition 2.1(2), the complex $K_\bullet(X; t)$ is a B -free resolution of the B -module $C_t(X)$ for any $t \geq -1$. By tensoring with A and taking the homology, we have that

$$\begin{aligned} H_p(K_\bullet(\widetilde{M}; t)) &\cong H_p(K_\bullet(X; t) \otimes_B A) \\ &\cong \text{Tor}_p^B(C_t(X), A) \end{aligned}$$

for all $p \geq 0$. On the other hand, since the ideal $\mathfrak{b}$ in B is generated by a regular sequence of length rn , the ordinary Koszul complex $K_\bullet(\mathfrak{b})$ associated to the sequence $\mathfrak{b}$ is a B -free resolution of A . Hence, by tensoring with $C_t(X)$, we can compute the Tor as follows:

$$\text{Tor}_p^B(C_t(X), A) \cong H_p(K_\bullet(\mathfrak{b}) \otimes_B C_t(X)).$$

Therefore, for any $p \geq 0$, we have

$$H_p(K_\bullet(\widetilde{M}; t)) \cong H_p(K_\bullet(\mathfrak{b}) \otimes_B C_t(X)).$$

It follows that for any $t \geq -1$ and any $q \geq 0$ we have the equality

$$\chi_q(K_\bullet(\widetilde{M}; t)) = \chi_q(K_\bullet(\mathfrak{b}) \otimes_B C_t(X)).$$

Here the right hand side is non-negative by Serre's Theorem ([12, Ch. IV, Appendix II]). Therefore $\chi_q(K_\bullet(\widetilde{M}; t)) \geq 0$. $\square$

4. PROOF OF THEOREM 1.3

Theorem 1.3 will be a consequence of the following more general result:

Theorem 4.1. *Let $(A, \mathfrak{m})$ be a Noetherian local ring of dimension $d > 0$.*

- (1) *For any rank $r > 0$, the inequality $\ell_A(H_0(K_\bullet(\widetilde{N}; t))) \geq e(F/N)$ holds true for any integer $t \geq -1$ and any parameter module N in $F = A^r$.*
- (2) *The following statements are equivalent:*
 - (i) *A is a Cohen-Macaulay local ring.*
 - (ii) *For some rank $r > 0$, there exists an integer $-1 \leq t \leq d$ and a parameter module N in $F = A^r$ such that the equality $\ell_A(H_0(K_\bullet(\widetilde{N}; t))) = e(F/N)$ holds true.*

When this is the case, the equality $\ell_A(H_0(K_\bullet(\widetilde{N}; t))) = e(F/N)$ holds true for any integer $-1 \leq t \leq d$ and any parameter module N in $F = A^r$ of any rank $r > 0$.

Proof. (1): Let N be a parameter module in $F = A^r$, and let $t \geq -1$. By Theorem 3.2 we obtain that

$$\begin{aligned} e(F/N) &= \chi(K_\bullet(\widetilde{N}; t)) \\ &= \ell_A(H_0(K_\bullet(\widetilde{N}; t))) - \chi_1(K_\bullet(\widetilde{N}; t)). \end{aligned}$$

Since $\chi_1(K_\bullet(\widetilde{N}; t)) \geq 0$ by Theorem 3.3, the desired inequality follows.

(2): Assume that A is Cohen-Macaulay. Let N be any parameter module in $F = A^r$ of any rank $r > 0$. Let $n = \mu_A(N) = d + r - 1$. Then $\text{grade } I(N) = \text{ht } I(N) = d = n - r + 1$. Hence, by Proposition 2.1(2), $K_\bullet(\widetilde{N}; t)$ is acyclic for all $-1 \leq t \leq n - r + 1 = d$. Therefore, by Theorem 3.2, we have $e(F/N) = \chi(K_\bullet(\widetilde{N}; t)) = \ell_A(H_0(K_\bullet(\widetilde{N}; t)))$. This proves the implication (i) $\Rightarrow$ (ii) and also the last assertion.

It remains to show the implication (ii) $\Rightarrow$ (i). Assume that there exist integers $r > 0$, $-1 \leq t \leq d$, and a parameter module N in $F = A^r$ such that $\ell_A(H_0(K_\bullet(\widetilde{N}; t))) = e(F/N)$. Arguing as in the proof of Theorem 3.3 and using the same notation, we get

$$\begin{aligned} \chi_1(K_\bullet(\mathfrak{b}) \otimes_B C_t(X)) &= \chi_1(K_\bullet(\widetilde{N}; t)) \\ &= \ell_A(H_0(K_\bullet(\widetilde{N}; t))) - e(F/N) \\ &= 0. \end{aligned}$$

We observe here that $\sqrt{\text{Ann}_B C_t(X)} = \sqrt{I_r(X)}$ (see [11, Lemma 2.7]). Thus $\dim_B C_t(X) = \dim B/I_r(X) = d + (n+1)(r-1) = rn$ (see [2, (5.12), Corollary]). Therefore $\mathfrak{b}$ is a parameter ideal of $C_t(X)$. Hence we have the equality

$$\ell_B(C_t(X)/\mathfrak{b}C_t(X)) - e(\mathfrak{b}; C_t(X)) = \chi_1(K_\bullet(\mathfrak{b}) \otimes_B C_t(X)) = 0,$$

where $e(\mathfrak{b}; C_t(X))$ is the multiplicity of the module $C_t(X)$ with respect to an ideal $\mathfrak{b}$. Since $\ell_B(C_t(X)/\mathfrak{b}C_t(X)) = e(\mathfrak{b}; C_t(X))$, this implies that $C_t(X)$ is a Cohen-Macaulay B -module. On the other hand, $\text{pd}_B C_t(X) = d$, because the complex

$K_{\bullet}(X; t)$ is a minimal B -free resolution of $C_t(X)$ of length $n - r + 1 = d$. Hence, by the Auslander-Buchsbaum formula, we have

$$\begin{aligned} d + rn &= \operatorname{pd}_B C_t(X) + \operatorname{depth}_B C_t(X) \\ &= \operatorname{depth} B \\ &\leq \dim B \\ &= d + rn. \end{aligned}$$

Thus $\operatorname{depth} B = \dim B$ so that B is Cohen-Macaulay. Therefore A is also a Cohen-Macaulay local ring. $\square$

Taking $t = 0, 1$ in Theorem 4.1 now readily gives Theorem 1.3.

We want to close this article with a question. For that, let us first recall the notion of a Buchsbaum local ring, which was introduced by Stückrad and Vogel (for more details on Buchsbaum rings, we refer the reader to [13]). Let A be a Noetherian local ring. Then A is said to be a *Buchsbaum ring* if the difference

$$\ell_A(A/Q) - e(A/Q)$$

between the colength and multiplicity of a parameter ideal Q in A is independent of the choice of Q . This difference, which is an invariant of a Buchsbaum ring A , is denoted by $I(A)$. The ring A is Cohen-Macaulay if and only if it is Buchsbaum and $I(A) = 0$. In this sense, the notion of a Buchsbaum ring is a natural generalization of that of a Cohen-Macaulay ring. In Theorem 1.3, the inequality $\ell_A(F/N) \geq e(F/N)$, for any parameter module N in F , is an analogue of the well-known inequality $\ell_A(A/Q) \geq e(A/Q)$ for any parameter ideal Q in A . Also, the characterization of the Cohen-Macaulay property of A based on the equality $\ell_A(F/N) = e(F/N)$ generalizes the usual one using parameter ideals. With these remarks in mind, it is natural to ask the following question:

Question 4.2. Let F be a fixed free module of rank $r > 0$. Is it then true that the difference

$$\ell_A(F/N) - e(F/N)$$

between the colength and multiplicity of a parameter module N in F is independent of the choice of N if the ring A is Buchsbaum?

REFERENCES

- [1] W. Bruns and U. Vetter, Length formulas for the local cohomology of exterior powers, *Math. Z.* 191 (1986), 145–158. MR812608 (87c:13016)
- [2] W. Bruns and U. Vetter, *Determinantal Rings*, Lecture Notes in Math., 1327, Springer-Verlag, Berlin-Heidelberg, 1988. MR953963 (89i:13001)
- [3] D. A. Buchsbaum and D. Eisenbud, Generic free resolutions and a family of generically perfect ideals, *Adv. in Math.* 18 (1975), 245–301. MR0396528 (53:391)
- [4] D. A. Buchsbaum and D. S. Rim, A generalized Koszul complex, *Bull. Amer. Math. Soc.* 69 (1963), 382–385. MR0148720 (26:6226)
- [5] D. A. Buchsbaum and D. S. Rim, A generalized Koszul complex. II. Depth and multiplicity, *Trans. Amer. Math. Soc.* 111 (1964), 197–224. MR0159860 (28:3076)
- [6] D. A. Buchsbaum and D. S. Rim, A generalized Koszul complex. III. A remark on generic acyclicity, *Proc. Amer. Math. Soc.* 16 (1965), 555–558. MR0177020 (31:1285)
- [7] D. Eisenbud, *Commutative algebra. With a view toward algebraic geometry*, Graduate Texts in Mathematics, 150, Springer-Verlag, New York, 1995. MR1322960 (97a:13001)
- [8] D. Kirby, A sequence of complexes associated with a matrix, *J. London Math. Soc.* 7 (1974), 523–530. MR0337939 (49:2708)

- [9] D. Kirby, On the Buchsbaum-Rim multiplicity associated with a matrix, J. London Math. Soc. (2) 32 (1985), no. 1, 57–61. MR813385 (87d:13025)
- [10] D. Kirby, Generalized Koszul complexes and the extension functor, Comm. Algebra 18 (1990), no. 4, 1229–1244. MR1059948 (91e:13015)
- [11] A. G. Rodicio, On the rigidity of the generalized Koszul complexes with applications to Hochschild homology, J. Algebra 167 (1994), no. 2, 343–347. MR1283291 (95e:13011)
- [12] J-P. Serre, Local Algebra (translated from the French by CheeWhye Chin), Springer Monographs in Mathematics, Springer-Verlag, Berlin-Heidelberg, 2000. MR1771925 (2001b:13001)
- [13] J. Stückrad and W. Vogel, Buchsbaum Rings and Applications, Springer-Verlag, Berlin-Heidelberg-New York-London-Paris-Tokyo, 1986. MR881220 (88h:13011a)

DEPARTMENT OF MATHEMATICS, SCHOOL OF SCIENCE AND TECHNOLOGY, MEIJI UNIVERSITY,
1-1-1 HIGASHIMITA, TAMA-KU, KAWASAKI 214-8571, JAPAN
E-mail address: `hayasaka@isc.meiji.ac.jp`

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF TAMPERE, 33014 TAMPEREEN
YLIOPISTO, FINLAND
E-mail address: `Eero.Hyry@uta.fi`