AN ENDPOINT ESTIMATE FOR THE CONE MULTIPLIER

YARYONG HEO, SUNGGEUM HONG, AND CHAN WOO YANG

(Communicated by Michael T. Lacey)

ABSTRACT. In this paper we consider an endpoint estimate for high-dimensional cone multipliers.

1. INTRODUCTION AND STATEMENTS OF THE RESULTS

Let $\widehat{\psi} \in C_0^\infty(\mathbb{R})$ be real-valued and supported in $\{1 < \tau < 2\}$. Then for each $\delta > 0$, we consider the convolution operators T^δ associated with the smooth cone multipliers given by

$$\widehat{T^\delta f}(\xi, \tau) = \widehat{f}(\xi, \tau) \left(1 - \frac{|\xi|^2}{|\tau|^2}\right)_+^\delta \widehat{\psi}(\tau), \quad (\xi, \tau) \in \mathbb{R}^d \times \mathbb{R}.$$

In the case $\delta > (d-1)/2$, the convolution kernel belongs to L^1 ; hence T^δ is an L^p -bounded operator for $1 \leq p \leq \infty$. In the case $0 < \delta < (d-1)/2$, it is conjectured that T^δ is an L^p -bounded operator for

$$\delta > \delta(p) := d|1/p - 1/2| - 1/2, \quad 1 < p < \infty.$$

This conjectured range is the same as for the d -dimensional Bochner-Riesz multiplier problem. Note that the cone multiplier problem implies the Bochner-Riesz multiplier problem. By now the Bochner-Riesz multiplier problem is understood in the range $p < (2d+4)/(d+4)$, $d \geq 2$ (see [8]). But compared to the Bochner-Riesz multiplier, little is known about the cone multiplier, and this conjecture still remains open for any $d \geq 2$. There are some partial results for this conjecture (see [3, 4, 5, 6, 7, 10]). In particular, in [10], the first author proved that T^δ is an L^p -bounded operator for $1 \leq p \leq 2(d-1)/(d+1)$, $\delta > \delta(p)$ and $d \geq 4$. This is the most recent result for high-dimensional cone multipliers. In this paper we consider an endpoint case $\delta = \delta(1) = (d-1)/2$.

Theorem 1.1. *If $d \geq 4$, then $T^{(d-1)/2}$ maps $L^1(\mathbb{R}^{d+1})$ to $L^{1,\infty}(\mathbb{R}^{d+1})$.*

Received by the editors October 6, 2008, and, in revised form, July 8, 2009.

2000 *Mathematics Subject Classification.* Primary 42B15.

Key words and phrases. Cone multipliers.

The first author was supported by the Korea Research Foundation Grant funded by the Korean Government (KRF-2008-357-C00002).

The second author was supported by the Korea Research Foundation Grant funded by the Korean Government (MEST) (No. 2009-0065011).

The third author was supported by the Korea Research Foundation Grant funded by the Korean Government (KRF-2008-331-C00016).

©2009 American Mathematical Society
 Reverts to public domain 28 years from publication

In the case of the Bochner-Riesz means, this result is well known for any dimension $d \geq 2$ (see [1, 2]). But for the cone multiplier there are some additional difficulties; i.e., in the case of the Bochner-Riesz multiplier, the main contribution of the convolution kernel comes from

$$(1.1) \quad |x|^{-\delta - \frac{d+1}{2}} e^{\pm 2\pi i |x|}, \quad |x| > 1, \quad x \in \mathbb{R}^d,$$

and in the case of the cone multiplier, the main contribution comes from

$$(1.2) \quad |x|^{-\delta - \frac{d+1}{2}} \psi(t \pm |x|), \quad |x| > 1, \quad (x, t) \in \mathbb{R}^d \times \mathbb{R}.$$

As we can see in Lemma 2.3 and [1, 2], the term $\psi(t \pm |x|)$ in (1.2) plays a similar role as $e^{\pm 2\pi i |x|}$ in (1.1). But the convolution kernel (1.2) of the cone multiplier is defined on $\mathbb{R}^d \times \mathbb{R}$. This makes the cone multiplier problem more difficult than the Bochner-Riesz problem. Even though (1.2) is defined on $\mathbb{R}^d \times \mathbb{R}$, we can see that it is essentially supported in the cone $|t \pm |x|| \leq 1$, and we can use this advantage together with M. Christ's stopping time arguments (Lemma 2.2).

Remark 1.2. Recently, the second author, F. Nazarov and A. Seeger [12] obtained $L^p(\mathbb{R}^{d+1}) \rightarrow L^{p,\infty}(\mathbb{R}^{d+1})$ inequalities for the cone multiplier $(1 - |\xi|^2/\tau^2)_+^{\delta(p)}$, $(\xi, \tau) \in \mathbb{R}^d \times \mathbb{R}$ with $1 < p < 2(d-1)/(d+1)$ and $d \geq 4$. See also [11] for the improvements upon the existing results in the so-called local smoothing problem for the wave equation in high dimensions.

2. REDUCTIONS AND THE PROOF OF THEOREM 1.1

Notation. If q is a dyadic cube in $\mathbb{R}^{d+1}$ with side-length 2^j , we write $\ell(q) = j$. For each $j \in \mathbb{Z}$, $\mathbf{D}_j$ denotes the collection of dyadic cubes $q \in \mathbb{R}^{d+1}$ with $\ell(q) = j$, and for each $q \in \mathbf{D}_j$, $2q$ denotes $q + [-2^j, 2^j]^{d+1}$. For two quantities A and B we shall write $A \lesssim B$ if $A \leq CB$ for some absolute positive constant C . Also we shall write $A \sim B$ if $A \lesssim B$ and $B \lesssim A$. The Lebesgue measure on $\mathbb{R}^{d+1}$ of a subset E will be denoted by $|E|$.

We need to show that

$$(2.1) \quad \left| \{(x, t) : |T^{(d-1)/2} f(x, t)| > \alpha\} \right| \leq C\alpha^{-1} \|f\|_1,$$

for each $\alpha > 0$. We may assume $f \geq 0$. Also by limiting arguments we may assume that $f \in L^1(\mathbb{R}^{d+1})$ have the form of a finite sum

$$(2.2) \quad f(x, t) = \sum_{q \in \mathbf{q}} \alpha_q \chi_q(x, t),$$

where $\alpha_q > 0$ and $\mathbf{q}$ is a finite, disjoint collection of dyadic cubes. Moreover if $\ell(q) \geq 0$, then by dividing q as smaller dyadic subcubes we may assume $\ell(q) = \ell \ll 0$ for all $q \in \mathbf{q}$. Let us explain the limiting arguments. Let $f \in L^1(\mathbb{R}^{d+1})$ with $f \geq 0$. Then there exists a sequence $\{f_m\}_{m=1}^\infty$ of functions such that each f_m has the form as in (2.2) and $\lim_{m \rightarrow \infty} \|f_m - f\|_1 = 0$. Also by choosing an appropriate subsequence we may assume

$$\|f_m\|_1 \leq 2\|f\|_1, \quad \|f_{m+1} - f_m\|_1 \leq 2^{-2m} \|f\|_1 \quad \text{for each } m.$$

Note that

$$\begin{aligned} T^{\frac{d-1}{2}} f(x, t) &= \lim_{m \rightarrow \infty} T^{\frac{d-1}{2}} f_m(x, t) \\ &= T^{\frac{d-1}{2}} f_1(x, t) + \sum_{m=1}^{\infty} T^{\frac{d-1}{2}} (f_{m+1} - f_m)(x, t) \end{aligned}$$

for almost every (x, t) . Hence if we prove (2.1) under the condition (2.2), then

$$\begin{aligned} &|\{ |T^{\frac{d-1}{2}} f(x, t)| > 2\alpha \}| \\ &\leq |\{ |T^{\frac{d-1}{2}} f_1(x, t)| > \alpha \}| + \sum_{m=1}^{\infty} |\{ |T^{\frac{d-1}{2}} (f_{m+1} - f_m)(x, t)| > 2^{-m}\alpha \}| \\ &\leq C\alpha^{-1} \|f_1\|_1 + \sum_{m=1}^{\infty} C(2^{-m}\alpha)^{-1} \|f_{m+1} - f_m\|_1 \\ &\leq C\alpha^{-1} \|f\|_1. \end{aligned}$$

Therefore, from now on we assume that

$$(2.3) \quad f = \sum_{q \in \mathbf{q}} \alpha_q |q|^{\frac{\chi_q}{|q|}} = \sum_{q \in \mathbf{q}} \lambda_q a_q,$$

where $\lambda_q = \alpha_q |q|$, $a_q = \chi_q / |q|$ and $\mathbf{q}$ is a finite, disjoint collection of dyadic cubes in $\mathbf{D}_\ell$ for some $\ell \ll 0$.

Next we compute the inverse Fourier transform $K^\delta(x, t)$ of $(1 - |\xi/\tau|^2)_+^\delta \widehat{\psi}(\tau)$. First note that

$$K^\delta(x, t) = \int_{\mathbb{R}^d} \int_{\mathbb{R}} e^{2\pi i \tau(x \cdot \xi + t)} (1 - |\xi|^2)_+^\delta \tau^d \widehat{\psi}(\tau) d\tau d\xi,$$

and by integration by parts via

$$\frac{\partial}{\partial \tau} e^{2\pi i \tau(x \cdot \xi + t)} = 2\pi i (x \cdot \xi + t) e^{2\pi i \tau(x \cdot \xi + t)},$$

we have

$$|K^\delta(x, t)| \leq C_N \sup_{|\xi| \leq 1} (1 + |x \cdot \xi + t|)^{-N} \quad \text{for each } N > 0.$$

From this, it is easy to see that $K^\delta(x, t) \chi_{(|x| \lesssim 1)} \in L^1(\mathbb{R}^{d+1})$. Therefore, from now on, we assume $|x| \gtrsim 1$. The inverse Fourier transform of the Bochner-Riesz multiplier $(1 - |\xi|^2)_+^\delta$, $\xi \in \mathbb{R}^d$ is given by

$$\pi^{-\delta} \Gamma(1 + \delta) |x|^{-\frac{d}{2} - \delta} J_{d/2 + \delta}(2\pi |x|),$$

where $J_{d/2 + \delta}$ is the Bessel function of order $d/2 + \delta$. So the inverse Fourier transform $K^\delta(x, t)$ of $(1 - |\xi/\tau|^2)_+^\delta \widehat{\psi}(\tau)$ is given by

$$K^\delta(x, t) = \int_{\mathbb{R}} e^{2\pi i \tau t} \left(\pi^{-\delta} \Gamma(1 + \delta) |\tau x|^{-\delta - d/2} J_{d/2 + \delta}(2\pi \tau |x|) \right) \tau^d \widehat{\psi}(\tau) d\tau.$$

It is well-known that, for all nonnegative integers N and n , as $r \rightarrow \infty$,

$$J_m(r) = e^{ir} \left[r^{-1/2} \sum_{j=0}^N a_j r^{-j/2} + A_N(r) \right] + e^{-ir} \left[r^{-1/2} \sum_{j=0}^N b_j r^{-j/2} + B_N(r) \right],$$

$$\frac{d^n}{dr^n} A_N(r) = O(r^{-n-\frac{N+1}{2}}), \quad \frac{d^n}{dr^n} B_N(r) = O(r^{-n-\frac{N+1}{2}}).$$

For reference, see pages 334 – 338 in Stein's book [13]. Therefore from the asymptotic expansion of the Bessel function $J_{d/2+\delta}$, for any positive integers N and M we have

$$K^\delta(x, t) = \sum_{j=0}^N \alpha_j^\pm |x|^{-(\delta+\frac{d+1+j}{2})} \mathcal{F}^{-1} \left[\tau^{-\delta+\frac{d-1-j}{2}} \widehat{\psi}(\tau) \right] (t \pm |x|) + F_N^\delta(x, t),$$

as $|x| \rightarrow \infty$; here $\mathcal{F}^{-1}$ denotes the inverse Fourier transform and

$$|F_N^\delta(x, t)| \leq C_{\delta, N, M} |x|^{-\delta-\frac{d+1+N}{2}} (1 + ||t| - |x||)^{-M}.$$

Note that

$$F_N^{(d-1)/2} \chi_{(|x| \geq 1)} \in L^1(\mathbb{R}^{d+1}) \quad \text{if } N \geq 1 \text{ and } M > 1.$$

Therefore it suffices to consider the terms

$$|x|^{-\delta-\frac{d+1+j}{2}} \mathcal{F}^{-1} \left[\tau^{-\delta+\frac{d-1-j}{2}} \widehat{\psi}(\tau) \right] (t \pm |x|), \quad |x| \geq 1,$$

with $j = 0, 1$, and $\delta = (d-1)/2$. From now on we will concentrate on the term

$$|x|^{-\delta-\frac{d+1}{2}} \mathcal{F}^{-1} \left[\tau^{-\delta+\frac{d-1}{2}} \widehat{\psi}(\tau) \right] (t - |x|), \quad |x| \geq 1, \delta = (d-1)/2,$$

and the other cases can be treated similarly. Now, we should treat the operation of convolution with

$$|x|^{-d} \psi(t - |x|), \quad |x| > 1.$$

For technical reasons, to obtain the convolution estimates in Lemma 2.3, fix a finite C^∞ partition of unity $\{\omega_i\}$ on the unit sphere $\mathbb{S}^{d-1}$, with each ω_i having very small support. Let ω be one of the C^∞ partition of unity $\{\omega_i\}$. Next choose $\eta \in C_0^\infty(\mathbb{R}^d)$, real-valued, radial and supported in $\{1/2 \leq |x| \leq 2\}$, so that $\sum_{j \in \mathbb{Z}} \eta(2^{-j}x) = 1$ on $\mathbb{R}^d \setminus \{0\}$. Also choose $\phi \in C_0^\infty(\mathbb{R})$, real-valued and supported in $\{|t| \leq 2\}$, so that

$$\sum_{n \in \mathbb{Z}} \phi(t - n) := \sum_{n \in \mathbb{Z}} \phi_n(t) = 1.$$

Then it suffices to treat the operation of convolution with

$$\sum_{n \in \mathbb{Z}} \sum_{j \geq 0} |x|^{-d} \psi(t - |x|) \omega(x/|x|) \phi(t - |x| - n) \eta(2^{-j}x) := \sum_{n \in \mathbb{Z}} \sum_{j \geq 0} K_j^n(x, t).$$

Let f be as in (2.3). Then it suffices to show that

$$\left| \left\{ (x, t) : \left| \left(\sum_{j \geq 0} K_j^n \right) * f(x, t) \right| > \alpha \right\} \right| \leq C_N (1 + |n|)^{-N} \alpha^{-1} \|f\|_1,$$

for each $n \in \mathbb{Z}$ and $\alpha > 0$. The following Lemma 2.1 is the standard Calderón-Zygmund decomposition, and we omit the proof (see Lemma 4.1 in [2]).

Lemma 2.1. *Suppose $\beta > 0$ is given. Then for any finite collection $\mathbf{q}$ of dyadic cubes q and associated positive scalars λ_q , there exists a collection of pairwise*

disjoint dyadic cubes $\{S : S \in \mathbf{S}\}$ such that

- (1) $\sum_{q: q \subset S} \lambda_q \leq 2^{d+1} \beta |S|,$
- (2) $\sum_{S: S \in \mathbf{S}} |S| \leq \beta^{-1} \sum_{q \in \mathbf{q}} \lambda_q,$
- (3) $\left\| \sum_{q: \text{not contained in any } S} \frac{\lambda_q}{|q|} \chi_q \right\|_{\infty} \leq \beta.$

Let $\mathbf{C}$ be the collection of dyadic cubes $q \in \mathbf{q}$ which are contained in some $S \in \mathbf{S}$. For each $q \in \mathbf{C}$ we define $S(q)$ as the unique $S \in \mathbf{S}$ containing q . The following is a refined Calderón-Zygmund decomposition whose proof relies on a stopping time argument.

Lemma 2.2 (cf. Lemma 5.1 in [2] or Lemma 5 in [9]). *Given $\beta > 0$ there exists a function $\Gamma : \mathbf{C} \rightarrow \mathbb{Z}$ and a measurable set E such that*

- (1) $|E| \leq C \left(\beta^{-1} \sum_{q \in \mathbf{C}} \lambda_q + \sum_{S \in \mathbf{S}} |S| \right),$
- (2) $\{q + \text{supp}(K_j^n)\} \subset E$ for all $j < \Gamma(q)$ and $q \in \mathbf{C},$
- (3) $\ell(S(q)) < \Gamma(q)$ for each $q \in \mathbf{C},$
- (4) for each $\tau, \ell \in \mathbb{Z}$ with $\ell \leq \tau$, and any $Q \in \mathbf{D}_{\ell}$ we have

$$\sum_{q \subset Q, q \in \mathbf{C}, \Gamma(q) \leq \tau} \lambda_q \leq \beta 2^{d(\tau+1)+\max(0, \ell)}.$$

For the proof we use the usual two-parameter stopping time arguments, and we construct an exceptional set E by combining stopping time arguments with the support condition of the kernel K_j^n ; i.e., if $\ell(Q) \leq \tau$ and $\tau \geq 0$, then

$$(2.4) \quad \left| \bigcup_{j=0}^{\tau} \left(Q + \text{supp}(K_j^n) \right) \right| \leq C 2^{d\tau + \max(0, \ell(Q))}.$$

If $S \in \mathbf{S}$ has side-length 2^j , then (1) of Lemma 2.1 says that

$$\sum_{q: q \subset S} \lambda_q \leq 2^{d+1} \beta |S|.$$

But if $Q \subset S$, then by (4) of Lemma 2.2 we have a more delicate estimate:

$$\sum_{q \subset Q, q \in \mathbf{C}, \Gamma(q) \leq j} \lambda_q \leq 2^d \beta 2^{jd + \max(0, \ell(Q))}.$$

This is why we referred to Lemma 2.2 as a refined Calderón-Zygmund decomposition. The proof will be given in Section 4.

Lemma 2.3. *Let $\tilde{K}_j^n$ denote the conjugate of K_j^n . Then for each $i < j$ and every $N > 0$ we have*

- (1) $|K_j^n * \tilde{K}_j^n(x, t)| \leq C_N (1 + |n|)^{-2N} 2^{-dj} (1 + |(x, t)|)^{\frac{1-d}{2}} \chi_{(|(x, t)| \leq 2^{j+4}}(x, t),$
- (2) $|K_j^n * \tilde{K}_i^n(x, t)| \leq C_N (1 + |n|)^{-2N} 2^{-dj} 2^{\frac{1-d}{2}i} \chi_{(|x| \leq 2^{j+4}}(x) \chi_{(|t - |x|| \leq 2^{i+4}}(x, t).$

The proof of Lemma 2.3 will be given in Section 3. For the moment we assume Lemmas 2.2 and 2.3 and prove Theorem 1.1. Let f be as in (2.3). We need to show that

$$(2.5) \quad \left| \left\{ (x, t) : \left| \left(\sum_{j \geq 0} K_j^n \right) * f(x, t) \right| > \alpha \right\} \right| \leq C_N (1 + |n|)^{-N} \alpha^{-1} \sum \lambda_q,$$

for each $\alpha > 0$. Apply Lemma 2.1 to the collection of dyadic cubes q and associated λ_q appearing in the definition of f with $\beta = (1 + |n|)^N \alpha$. Let $\mathbf{S}$ be as in Lemma 2.1 and define

$$b = \sum_{S \in \mathbf{S}} \sum_{q \subset S} \lambda_q a_q, \quad g = f - b.$$

Then $\|g\|_\infty \leq \beta$ and so by (1) of Lemma 2.3, for $d \geq 4$ we have

$$\begin{aligned} \left\| \left(\sum_j K_j^n \right) * g \right\|_2 &\leq \sum_j \|g\|_2 \|K_j^n * \tilde{K}_j^n\|_1^{1/2} \\ &\leq \|g\|_2 \sum_j \left(C_N (1 + |n|)^{-N} 2^{-j(d-3)/4} \right) \\ &\leq C_N (1 + |n|)^{-N} \beta^{1/2} \|f\|_1^{1/2} \\ &\leq C_N (1 + |n|)^{-N/2} \alpha^{1/2} \|f\|_1^{1/2}. \end{aligned}$$

Therefore by Tchebychev's inequality we have

$$(2.6) \quad \left| \left\{ (x, t) : \left| \left(\sum_j K_j^n \right) * g(x, t) \right| > \alpha \right\} \right| \leq \alpha^{-2} \left\| \left(\sum_j K_j^n \right) * g \right\|_2^2 \leq C_N (1 + |n|)^{-N} \alpha^{-1} \|f\|_1.$$

Let $\mathbf{S}$ be as above and $\mathbf{C}$ be the collection of q 's appearing in the definition of b . Then apply Lemma 2.2 with $\mathbf{S}$, $\mathbf{C}$. By (1) of Lemma 2.2 we have an exceptional set E such that

$$(2.7) \quad |E| \leq C \left(\beta^{-1} \sum_{q \in \mathbf{C}} \lambda_q + \sum_{S \in \mathbf{S}} |S| \right) \leq C \beta^{-1} \|f\|_1 \leq C (1 + |n|)^{-N} \alpha^{-1} \|f\|_1.$$

By (2.6) and (2.7), (2.5) will follow from

$$(2.8) \quad \left| \left\{ (x, t) \in \mathbb{R}^{d+1} \setminus E : \left| \left(\sum_j K_j^n \right) * b(x, t) \right| > \alpha \right\} \right| \leq C_N (1 + |n|)^{-N} \alpha^{-1} \sum_q \lambda_q.$$

By Tchebychev's inequality, (2.8) will follow from

$$(2.9) \quad \left\| \left(\sum_j K_j^n \right) * b \right\|_{L^2(\mathbb{R}^{d+1} \setminus E)}^2 \leq C_N (1 + |n|)^{-N} \alpha \sum_q \lambda_q.$$

By (2) of Lemma 2.2, for each $q \in \mathbf{C}$, $K_j^n * a_q$ is supported in E unless $j \geq \Gamma(q)$. Thus we have

$$\begin{aligned} &\left\| \sum_j b * K_j^n \right\|_{L^2(\mathbb{R}^{d+1} \setminus E)} \\ (2.10) \quad &\leq \left\| \sum_j \left(\sum_{\Gamma(q) \leq j} \lambda_q a_q \right) * K_j^n \right\|_2 \\ &\leq \left\| \sum_j \left(\sum_{\Gamma(q) \leq 0} \lambda_q a_q \right) * K_j^n \right\|_2 + \left\| \sum_j \left(\sum_{0 < \Gamma(q) \leq j} \lambda_q a_q \right) * K_j^n \right\|_2 \end{aligned}$$

and

$$\left\| \sum_j \left(\sum_{0 < \Gamma(q) \leq j} \lambda_q a_q \right) * K_j^n \right\|_2 = \left\| \sum_j \sum_{0 \leq s < j} \left(\sum_{\Gamma(q)=j-s} \lambda_q a_q \right) * K_j^n \right\|_2.$$

Note that

$$\sum_j \sum_{0 \leq s < j} = \sum_{s \geq 0} \sum_{j > s};$$

hence we have

$$(2.11) \quad \left\| \sum_j \sum_{0 \leq s < j} \left(\sum_{\Gamma(q)=j-s} \lambda_q a_q \right) * K_j^n \right\|_2 \leq \sum_{s \geq 0} \left\| \sum_{j > s} \left(\sum_{\Gamma(q)=j-s} \lambda_q a_q \right) * K_j^n \right\|_2.$$

Now by (2.10) and (2.11) we have

$$\left\| \sum_j b * K_j^n \right\|_{L^2(\mathbb{R}^{d+1} \setminus E)} \leq F_1 + F_2,$$

where

$$F_1 = \sum_j \left\| \left(\sum_{\Gamma(q) \leq 0} \lambda_q a_q \right) * K_j^n \right\|_2, \quad F_2 = \sum_{s \geq 0} \left\| \sum_{j > s} \left(\sum_{\Gamma(q)=j-s} \lambda_q a_q \right) * K_j^n \right\|_2.$$

Estimation of part F_1 . Let $B_0 = \sum_{\Gamma(q) \leq 0} \lambda_q a_q$. For each $q \in \mathbf{C}$ there exists a unique $S(q) \in \mathbf{S}$ containing q ; hence if $\Gamma(q) \leq 0$, then by the condition $\ell(S(q)) \leq \Gamma(q)$ we have $\ell(S(q)) \leq 0$. Therefore by (1) of Lemma 2.1, we can see that

$$(2.12) \quad \|B_0 \chi_Q\|_1 = \sum_{q \subset Q, \Gamma(q) \leq 0} \lambda_q \leq C\beta \quad \text{for each } Q \in \mathbf{D}_0.$$

By (1) of Lemma 2.3, we have

$$\|K_j^n * \tilde{K}_j^n * B_0\|_\infty \leq C_N (1 + |n|)^{-2N} 2^{-jd} \sup_{x \in \mathbb{R}^{d+1}} \left(\int_{|x-y| \lesssim 2^j} \frac{B_0(y)}{(1 + |x-y|)^{\frac{d-1}{2}}} dy \right).$$

By (2.12) it is easy to see that

$$\begin{aligned} \left(\int_{|x-y| \lesssim 2^j} \frac{B_0(y)}{(1 + |x-y|)^{\frac{d-1}{2}}} dy \right) &\leq \left(\int_{|x-y| \lesssim 2^j} \frac{C\beta}{(1 + |x-y|)^{\frac{d-1}{2}}} dy \right) \\ &\leq C\beta 2^{\frac{d+3}{2}j}. \end{aligned}$$

Therefore we have

$$\|K_j^n * \tilde{K}_j^n * B_0\|_\infty \leq C_N (1 + |n|)^{-2N} \beta 2^{\frac{3-d}{2}j},$$

and so for $d \geq 4$,

$$\begin{aligned} F_1 &= \sum_j \|B_0 * K_j^n\|_2 = \sum_j |\langle B_0, K_j^n * \tilde{K}_j^n * B_0 \rangle|^{1/2} \\ (2.13) \quad &\leq \|B_0\|_1^{1/2} \sum_j \|K_j^n * \tilde{K}_j^n * B_0\|_\infty^{1/2} \\ &\leq C_N (1 + |n|)^{-N} \beta^{1/2} \|f\|_1^{1/2}. \end{aligned}$$

Estimation of part F_2 . For each $l > 0$, let $B_l = \sum_{\Gamma(q)=l} \lambda_q a_q$. Then

$$\begin{aligned} \left\| \sum_{j>s} B_{j-s} * K_j^n \right\|_2^2 &\leq \sum_{j>s} \|B_{j-s} * K_j^n\|_2^2 \\ &\quad + 2 \sum_{j>s} |\langle B_{j-s} * K_j^n, B_{j-1-s} * K_{j-1}^n \rangle| \\ &\quad + 2 \sum_{j>s} \sum_{s<i<j-1} |\langle B_{i-s} * K_i^n * \tilde{K}_j^n, B_{j-s} \rangle| \\ &:= A_1(s) + A_2(s) + A_3(s). \end{aligned}$$

For the part $A_1(s)$, we write

$$\begin{aligned} \|B_{j-s} * K_j^n\|_2^2 &= \sum_{\Gamma(q), \Gamma(q')=j-s} \lambda_q \lambda_{q'} \langle a_{q'} * K_j^n, a_q * K_j^n \rangle \\ &\leq \sum_{m=0}^{j+4} \sum_{\substack{\Gamma(q), \Gamma(q')=j-s \\ 1+\text{dist}(q, q') \sim 2^m}} \lambda_q \lambda_{q'} |\langle a_{q'} * K_j^n * \tilde{K}_j^n, a_q \rangle| \\ &\leq \sum_{m=0}^{j-s+2} \sum_{\substack{\Gamma(q), \Gamma(q')=j-s \\ 1+\text{dist}(q, q') \sim 2^m}} + \sum_{m=j-s+3}^{j+4} \sum_{\substack{\Gamma(q), \Gamma(q')=j-s \\ 1+\text{dist}(q, q') \sim 2^m}} \\ &:= \text{I} + \text{II}. \end{aligned}$$

Estimation of part I. For each fixed m and q' consider the contribution of all λ_q over all q so that $1 + \text{dist}(q, q') \sim 2^m$. All such q are contained in the union of a fixed number of $\mathbf{D}_m$. Hence when $0 \leq m \leq j-s+2$, by (1) of Lemma 2.3 we have

$$\text{I} \leq C_N (1 + |n|)^{-2N} 2^{-jd} \sum_{q'} \lambda_{q'} \sum_{m=0}^{j-s+2} \sum_{q: 1+\text{dist}(q, q') \sim 2^m} 2^{\frac{1-d}{2}m} \lambda_q.$$

By (4) of Lemma 2.2 we have

$$\sum_{q: 1+\text{dist}(q, q') \sim 2^m, \Gamma(q)=j-s} \lambda_q \leq C \beta 2^{(j-s)d+m},$$

and so for $d \geq 4$,

$$\text{I} \leq C_N (1 + |n|)^{-2N} 2^{-ds} \beta \sum_{\Gamma(q')=j-s} \lambda_{q'}.$$

Estimation of part II. Next, consider all q with $\text{dist}(q, q') \sim 2^m$ and $j-s+3 \leq m \leq j+4$. Recall that each $q \in \mathbf{C}$ is contained in some $S(q) \in \mathbf{S}$. Since $\ell(S(q)) < \Gamma(q) = j-s$, we have $\text{dist}(S(q), q') \sim 2^m$ and so

$$\begin{aligned} \text{II} &\leq C_N (1 + |n|)^{-2N} 2^{-jd} \sum_{q'} \lambda_{q'} \sum_{m=j-s+3}^{j+4} \sum_{q: \text{dist}(q, q') \sim 2^m} 2^{\frac{1-d}{2}m} \lambda_q \\ &\leq C_N (1 + |n|)^{-2N} 2^{-jd} \sum_{q'} \lambda_{q'} \sum_{\substack{S \in \mathbf{S} \\ 2j-s+2 \leq \text{dist}(S, q') \leq 2j+5}} \sum_{q \in S} \text{dist}(S, q')^{\frac{1-d}{2}} \lambda_q. \end{aligned}$$

By (1) of Lemma 2.1, for each $S \in \mathbf{S}$ we have $\sum_{q \subset S} \lambda_q \leq C\beta|S|$, and so

$$\begin{aligned} \sum_{\substack{S \in \mathbf{S} \\ 2^{j-s}+2 \leq \text{dist}(S, q') \leq 2^{j+5}}} \sum_{q \subset S} \text{dist}(S, q')^{\frac{1-d}{2}} \lambda_q &\leq C \sum_{\substack{S \in \mathbf{S} \\ 2^{j-s}+2 \leq \text{dist}(S, q') \leq 2^{j+5}}} \text{dist}(S, q')^{\frac{1-d}{2}} (\beta|S|) \\ &\leq C\beta \int_{y \in \mathbb{R}^{d+1}, 2^{j-s}+2 \leq |y| \leq 2^{j+5}} |y|^{\frac{1-d}{2}} dy \\ &\leq C2^{\frac{3-d}{2}j} \beta. \end{aligned}$$

Now we have

$$\Pi \leq C_N(1+|n|)^{-2N} 2^{\frac{3-d}{2}j} \beta \sum_{\Gamma(q')=j-s} \lambda_{q'},$$

and so for $d \geq 4$,

$$\begin{aligned} (2.14) \quad A_1(s) &\leq \sum_{j>s} C_N(1+|n|)^{-2N} \left(2^{-sd} + 2^{\frac{3-d}{2}j}\right) \beta \sum_{\Gamma(q')=j-s} \lambda_{q'} \\ &\leq C_N(1+|n|)^{-2N} 2^{\frac{3-d}{2}s} \beta \sum_{q'} \lambda_{q'}. \end{aligned}$$

Similarly we have

$$(2.15) \quad A_2(s) \leq C_N(1+|n|)^{-2N} 2^{\frac{3-d}{2}s} \beta \sum_{q'} \lambda_{q'}.$$

For the part $A_3(s)$, if $\Gamma(q) = i-s$, then $\ell(q), S(q) \leq i-s$. So by (1) of Lemma 2.1 and (2) of Lemma (2.3) together with the condition

$$\text{supp}(K_i^n * \tilde{K}_j^n) \subset \{(x, t) : |x| \leq 2^{j+4}, |t - |x|| \leq 2^{i+4}\},$$

we have

$$\begin{aligned} \|B_{i-s} * K_i^n * \tilde{K}_j^n\|_\infty &\leq C_N(1+|n|)^{-2N} 2^{-jd} 2^{\frac{1-d}{2}i} \sum_{S \in \mathbf{S}} \sum_{\substack{q \subset S \\ \Gamma(q)=i-s}} \lambda_q \\ &\leq C_N(1+|n|)^{-2N} 2^{-jd} 2^{\frac{1-d}{2}i} \sum_{S \in \mathbf{S}} \beta|S|, \end{aligned}$$

where the sums are taken over all $S \in \mathbf{S}$ such that

$$S \subset \{(x, t) : |x| \leq 2^{j+4}, |t - |x|| \leq 2^{i+4}\}.$$

So we have $\sum_{S \in \mathbf{S}} |S| \leq C2^{jd+i}$ and

$$\|B_{i-s} * K_i^n * \tilde{K}_j^n\|_\infty \leq C_N(1+|n|)^{-2N} 2^{\frac{3-d}{2}i} \beta.$$

Therefore for $d \geq 4$,

$$\begin{aligned} (2.16) \quad A_3(s) &= \sum_{j>s} \sum_{s<i<j-1} |\langle B_{i-s} * K_i^n * \tilde{K}_j^n, B_{j-s} \rangle| \\ &\leq C_N(1+|n|)^{-2N} \sum_{j>s} \sum_{i>s} 2^{\frac{3-d}{2}i} \beta \|B_{j-s}\|_1 \\ &\leq C_N(1+|n|)^{-2N} 2^{\frac{3-d}{2}s} \beta \sum_q \lambda_q. \end{aligned}$$

Finally from (2.13), (2.14), (2.15) and (2.16), for $d \geq 4$ we have

$$F_2 \leq \sum_{s \geq 0} (A_1(s) + A_2(s) + A_3(s))^{1/2} \leq C_N (1 + |n|)^{-N} \beta^{1/2} \left(\sum_q \lambda_q \right)^{1/2},$$

and we are done.

3. PROOF OF LEMMA 2.3

The proof is similar to Lemma 3.1 in [2]. For (1), let $\eta_1(x) := \eta(x)|x|^{-d}$. Then we have

$$K_j^n * \tilde{K}_j^n(x, t) = 2^{-2dj} \int_{\mathbb{R}} \int_0^\infty F_j(x, t, s, r) \overline{\psi}(s-r) \eta_1(2^{-j}r) \phi_n(s-r) r^{d-1} dr ds,$$

where

$$F_j(x, t, s, r) = \int_{\mathbb{S}^{d-1}} \psi(t + s - |x + r\theta|) \omega[(x + r\theta)/|x + r\theta|] \omega(\theta) \\ \times \phi_n(t + s - |x + r\theta|) \eta_1[2^{-j}(x + r\theta)] d\theta.$$

Note that

$$\psi(t + s - |x + r\theta|) = \int_{\mathbb{R}} e^{2\pi i \tau(t+s-|x+r\theta|)} \widehat{\psi}(\tau) d\tau.$$

Hence we have

$$(3.1) \quad F_j(x, t, s, r) = \int_{\mathbb{R}} e^{2\pi i \tau(t+s)} \widehat{\psi}(\tau) G_j(x, t, s, r, \tau) d\tau,$$

where

$$(3.2) \quad G_j(x, t, s, r, \tau) = \int_{\mathbb{S}^{d-1}} e^{-2\pi i \tau|x+r\theta|} \omega[(x + r\theta)/|x + r\theta|] \omega(\theta) \\ \times \phi_n(t + s - |x + r\theta|) \eta_1[2^{-j}(x + r\theta)] d\theta.$$

It suffices to show that

$$|G_j(x, t, s, r, \tau)| \leq C (1 + |(x, t)|)^{(1-d)/2},$$

uniformly for $\tau \sim 1$, s and $r \sim 2^j$. Suppose that the function ω from the partition of unity has sufficiently small support about $(0, \dots, 0, 1)$. Then we use the local coordinate chart

$$\theta = (\theta_1, \dots, \theta_d) = (\theta_1, \dots, \theta_{d-1}, \sqrt{1 - \theta_1^2 - \dots - \theta_{d-1}^2}).$$

Then by direct calculation, for $1 \leq l \leq d-1$,

$$(3.3) \quad \frac{\partial}{\partial \theta_l} (|x + r\theta|) = r \left(\frac{\theta_d x_l - \theta_l x_d}{|x + r\theta| \theta_d} \right),$$

$$(3.4) \quad \frac{\partial^2 |x + r\theta|}{\partial \theta_l^2} = r \left(\frac{|x + r\theta|^2 (\theta_d^2 - \theta_l^2) (-x_d) - r \theta_d |\theta_d x_l - \theta_l x_d|^2}{|x + r\theta|^3 \theta_d^3} \right),$$

and for $1 \leq l \neq m \leq d-1$,

$$(3.5) \quad \frac{\partial^2 |x + r\theta|}{\partial \theta_l \partial \theta_m} = -r \left(\frac{\theta_l \theta_m x_d |x + r\theta|^2 + r \theta_d (\theta_d x_m - \theta_m x_d) (\theta_d x_l - \theta_l x_d)}{|x + r\theta|^3 \theta_d^3} \right).$$

Note that

$$(3.6) \quad \frac{\partial^2 |x + r\theta|}{\partial \theta_l^2} (0, \dots, 0, 1) = r \left(\frac{|x + r\theta|^2 (-x_d) - x_l^2}{|x + r\theta|^3} \right)$$

and

$$(3.7) \quad \frac{\partial^2 |x + r\theta|}{\partial \theta_l \partial \theta_m}(0, \dots, 0, 1) = \frac{-r^2 x_l x_m}{|x + r\theta|^3}, \quad 1 \leq l \neq m \leq d-1.$$

In the case $|x_l| \gtrsim |x|$ for some $1 \leq l \leq d-1$, by (3.3),

$$\left| \frac{\partial |x + r\theta|}{\partial \theta_l} \right| \gtrsim |x_l|$$

uniformly for $|x + r\theta| \sim 2^j$, $r \sim 2^j$. By integrating by parts via

$$\frac{\partial}{\partial \theta_l} \left(e^{-2\pi i \tau |x + r\theta|} \right) = -2\pi i \tau \frac{\partial |x + r\theta|}{\partial \theta_l} \left(e^{-2\pi i \tau |x + r\theta|} \right),$$

we have

$$|G_j(x, t, s, r, \tau)| \leq C_N (1 + |x_l|)^{-N} \leq C_N (1 + |x|)^{-N}.$$

In the case $|x_d| \gtrsim |x|$, from (3.4), (3.5), (3.6), (3.7), together with the conditions $|x + r\theta| \sim 2^j$, $r \sim 2^j$, we can see that each absolute value of the eigenvalue of the Hessian matrix

$$\frac{\partial^2 |x + r\theta|}{\partial^2 \theta}$$

is bounded below by $C|x_d|$ uniformly for $2^{j-1} < r < 2^{j+1}$ when θ and $(x + r\theta)/|x + r\theta|$ are both in the support of ω . By the methods of stationary phase, we have

$$|G_j(x, t, s, r, \tau)| \leq C (1 + |x_d|)^{(1-d)/2} \leq (1 + |x|)^{(1-d)/2}.$$

Hence we have

$$(3.8) \quad |K_j^n * \tilde{K}_j^n(x, t)| \leq C_N (1 + |n|)^{-2N} 2^{-jd} (1 + |x|)^{(1-d)/2}.$$

Next recall that

$$\begin{aligned} K_j^n * \tilde{K}_j^n(x, t) &= 2^{-2dj} \int_{\mathbb{R}} \int_{\mathbb{R}^d} \psi(t + s - |x + y|) \bar{\psi}(s - |y|) \omega[(x + y)/|x + y|] \\ &\quad \times \omega(y/|y|) \phi_n(t + s - |x + y|) \phi_n(s - |y|) \eta_1[2^{-j}(x + y)] \eta_1(2^{-j}y) dy ds. \end{aligned}$$

Hence $K_j^n * \tilde{K}_j^n(x, t)$ is supported in

$$|t + s - |x + y| - n| \leq 2, \quad |s - |y| - n| \leq 2,$$

and by the triangle inequality, this implies that

$$(3.9) \quad |t| \leq |t + s - |x + y| - n| + |-s + |y| + n| + |-|y| + |x - y|| \leq 2 + 2 + |x|.$$

Therefore if $|x| \leq 2$, then $|(x, t)| \leq 6$, and (3.8) implies that

$$|K_j^n * \tilde{K}_j^n(x, t)| \leq C_N (1 + |n|)^{-2N} 2^{-jd} (1 + |(x, t)|)^{-N}.$$

If $|x| \geq 2$, then by (3.9) we have $|t| \leq 3|x|$, and (3.8) implies that

$$|K_j^n * \tilde{K}_j^n(x, t)| \leq C_N (1 + |n|)^{-2N} 2^{-jd} (1 + |(x, t)|)^{(1-d)/2}.$$

For (2), let $F_j(x, t, s, r)$ and $G_j(x, t, s, r, \tau)$ be the same as in (3.1) and (3.2). Then we have

$$K_j^n * \tilde{K}_i^n(x, t) = 2^{-dj-di} \int_{\mathbb{R}} \int_0^\infty F_j(x, t, s, r) \bar{\psi}(s - r) \eta_1(2^{-i}r) \phi_n(s - r) r^{d-1} dr ds.$$

Since $|x + r\theta| \sim 2^j$, $r \sim 2^i$, we have $|x| \sim 2^j$. Hence in the case $|x_l| \gtrsim |x|$ for some $1 \leq l \leq d-1$, from (3.3) we can see that

$$\left| \frac{\partial |x + r\theta|}{\partial \theta_l} \right| \gtrsim 2^i,$$

uniformly for $|x + r\theta| \sim 2^j$ and $r \sim 2^i$, and by integrating by parts via

$$\frac{\partial}{\partial \theta_l} \left(e^{-2\pi i \tau |x + r\theta|} \right) = -2\pi i \tau \frac{\partial |x + r\theta|}{\partial \theta_l} \left(e^{-2\pi i \tau |x + r\theta|} \right),$$

we have

$$|G_j(x, t, s, r, \tau)| \leq C_N 2^{-Ni}.$$

In the case $|x_d| \gtrsim |x|$, each absolute value of the eigenvalue of the Hessian matrix

$$\frac{\partial^2 |x + r\theta|}{\partial^2 \theta}$$

is bounded below by $C2^i$ uniformly for $r \sim 2^i$ when θ and $(x + r\theta)/|x + r\theta|$ are both in support of ω . By the methods of stationary phase, we have

$$|G_j(x, t, s, r, \tau)| \leq C2^{\frac{1-d}{2}i}.$$

Hence in the case $|x_d| \gtrsim |x|$ we have

$$(3.10) \quad |K_j^n * \tilde{K}_i^n(x, t)| \leq C_N (1 + |n|)^{-2N} 2^{-jd} 2^{\frac{1-d}{2}i}.$$

Note that $K_j^n * \tilde{K}_i^n(x, t)$ is supported in

$$|t + s - |x + y| - n| \leq 2, \quad |s - |y| - n| \leq 2, \quad |x + y| \leq 2^{j+1}, \quad |y| \leq 2^{i+1}.$$

Hence $K_j^n * \tilde{K}_i^n(x, t)$ is supported in

$$(3.11) \quad \begin{aligned} |t - |x|| &\leq |t + s - |x + y| - n| + ||x + y| - |x|| + |-s + n| \\ &\leq 2 + |y| + (2 + |y|) \leq 2^{i+4}, \end{aligned}$$

and from (3.10) and (3.11) we have (2) of Lemma 2.3.

4. PROOF OF LEMMA 2.2

The following are the usual two-parameter stopping-time arguments. These will be discussed in more detail below. We will combine these arguments with the support condition of the kernel K_j^n to construct an exceptional set E . Let $m = \min\{\ell(q) : q \in \mathbf{C}\}$. Select an integer τ_0 such that

$$\tau_0 > \max\{\ell(q) : q \in \mathbf{C}\}, \quad \sum_{q \in \mathbf{C}} \lambda_q < \beta 2^{d\tau_0 + \max(0, m)}.$$

For each fixed $\tau \in \mathbb{Z}$ with $\tau \leq \tau_0$, we will define a sequence of functions $\Lambda_{\tau, \ell} : \mathbf{D}_\ell \rightarrow \mathbb{R}$ by a descending induction on $\ell \in \mathbb{Z}$ with $\ell \leq \tau$, and proceed with the same construction by a descending induction on τ . At each step, we will define subsets $\mathbf{C}_1, \mathbf{C}_2$ of $\mathbf{C}$ which will increase as we proceed. Let $\mathbf{C}_1, \mathbf{C}_2 \subset \mathbf{C}$ and $\tau \in \mathbb{Z}$ be fixed for the moment, and define **Inner loop** as

Inner loop. Define $\Lambda_{\tau,\ell} : \mathbf{D}_\ell \rightarrow \mathbb{R}$ with $\ell \leq \tau$. For each $Q \in \mathbf{D}_\ell$, define

$$\Lambda_{\tau,\ell}(Q) = \sum_{q \subset Q: q \notin \mathbf{C}_1 \cup \mathbf{C}_2} \lambda_q.$$

Begin with $\ell = \tau$. If

$$(4.1) \quad \Lambda_{\tau,\ell}(Q) > \beta 2^{d\tau + \max(0,\ell)},$$

then we say that “ Q is selected at step (τ, ℓ) ”. Put into $\mathbf{C}_1$ every q such that $q \subset Q$ and define $\Gamma(q) = 1 + \tau$. Repeat until $\ell < \min\{\ell(q) : q \in \mathbf{C}\}$. Actually this part of the process will be terminated once ℓ is smaller than m . Put into $\mathbf{C}_2$ every $q \in \mathbf{C} \setminus \mathbf{C}_1$ such that $\ell(q) \geq \tau$ and for such q define $\Gamma(q) = 1 + \ell(S(q))$. Actually every $q \in \mathbf{C} \setminus \mathbf{C}_1 \cup \mathbf{C}_2$ satisfies $\ell(q) \leq \tau - 1$.

Perform **Inner loop** with $\mathbf{C}_1 = \emptyset = \mathbf{C}_2$ and $\tau = \tau_0$. Next replace τ by $\tau - 1$ and repeat **Inner loop**. Repeat until $\tau = m - 1$. After this process we obtain $\mathbf{C} = \mathbf{C}_1 \cup \mathbf{C}_2$. Clearly all selected Q are disjoint and Γ is well defined. Note that there is the usual stopping-time condition

$$(4.2) \quad \Lambda_{\tau,\ell}(Q) \leq \beta 2^{d(\tau+1) + \max(0,\ell)},$$

which holds for all $Q \in \mathbf{D}_\ell$ when $\ell \leq \tau \leq \tau_0$. This is because if $\tau = \tau_0$, then the condition is clear from the choice of τ_0 and m . When $\ell \leq \tau \leq \tau_0$, suppose this fails. Then $\Lambda_{\tau+1,\ell}(Q) \geq \Lambda_{\tau,\ell}(Q) > \beta 2^{d(\tau+1) + \max(0,\ell)}$. This means that Q is selected at step $(\tau + 1, \ell)$; hence $\Lambda_{\tau,\ell}(Q) = 0$, and we have the contradiction.

Next we show (4), which says, for each $Q \in \mathbf{D}_\ell$ with $\ell \leq \tau$,

$$\sum_{q \subset Q: \Gamma(q) \leq \tau} \lambda_q \leq \beta 2^{d(\tau+1) + \max(0,\ell)}.$$

When $\tau \geq \tau_0$, this property is clear from the initial choice of τ_0 . When $\tau < \tau_0$, if we show that

$$(4.3) \quad \Lambda_{\tau,\ell}(Q) = \sum_{q \subset Q: q \notin \mathbf{C}_1 \cup \mathbf{C}_2} \lambda_q \geq \sum_{q \subset Q: \Gamma(q) \leq \tau} \lambda_q$$

for each $Q \in \mathbf{D}_\ell$ with $\ell \leq \tau < \tau_0$, then by combining (4.2) and (4.3), we have (4). Now (4.3) follows from the definition

$$\Lambda_{\tau,\ell}(Q) = \sum_{q \subset Q: q \notin \mathbf{C}_1 \cup \mathbf{C}_2} \lambda_q$$

and the fact that

$$\Gamma(q) \leq \tau \Rightarrow q \notin \mathbf{C}_1 \cup \mathbf{C}_2 \text{ at the beginning of step } (\tau, \ell).$$

This is because, if $q \in \mathbf{C}_1$, then $\Gamma(q) \geq 1 + \tau > \tau$, and if $q \in \mathbf{C}_2$, then $\Gamma(q) = 1 + \ell(S(q)) \geq 1 + (1 + \tau) > \tau$. Hence $\Gamma(q) \leq \tau$ implies $q \notin \mathbf{C}_1 \cup \mathbf{C}_2$, and so we have (4.3).

Next, we construct an exceptional set E by using the above stopping-time arguments. If Q is selected at step (τ, ℓ) , then we define $\tau(Q) = \tau$. If $\tau(Q) \geq 0$, then we define the tendril $T(Q)$ associated to Q by

$$T(Q) = \bigcup_{j < \tau(Q) + 1} (Q + \text{supp}(K_j^n)).$$

Also we define

$$E = E_1 \cup E_2, \quad E_1 = \bigcup_{S \in \mathbf{S}} 2S, \quad E_2 = \bigcup_{Q: \text{selected}, \tau(Q) \geq 0} T(Q).$$

Thus we have

$$|E_1| \leq C \sum_{S \in \mathbf{S}} |S|.$$

By the measure condition (2.4) of $T(Q)$ and the condition (4.1) of selected Q ,

$$\begin{aligned} |E_2| &\leq \sum_{Q: \text{selected}, \tau(Q) \geq 0} |T(Q)| \leq C \sum_{Q: \text{selected}} 2^{d\tau(Q) + \max(0, \ell(Q))} \\ &\leq C \frac{1}{\beta} \sum_{Q: \text{selected}} \Lambda_{\tau, \ell}(Q) \leq C \frac{1}{\beta} \sum \lambda_q. \end{aligned}$$

So we have (1). For (2), if $q \in \mathbf{C}_1$, then q belongs to some selected Q and $\Gamma(q) = \tau(Q) + 1$. Hence if $\Gamma(q) > j$, then

$$\bigcup_{j < \Gamma(q)} (q + \text{supp}(K_j^n)) \subset T(Q) \subset E_2,$$

and if $q \in \mathbf{C}_2$, then q belongs to some $S(q) \in \mathbf{S}$ and $\Gamma(q) = 1 + \ell(S(q))$. Hence if $\Gamma(q) > j$, then

$$\bigcup_{j < \Gamma(q)} \{q + \text{supp}(K_j^n)\} \subset 2S(q) \subset E_1.$$

For (3), let $\Gamma' := \Gamma$ and redefine

$$\Gamma(q) = \max\{\Gamma'(q), 1 + \ell(S(q))\}.$$

Then (1) and (3) are satisfied. We must check (2) and (4). For (2), if $\Gamma(q) = \Gamma'(q)$, then there is no problem. If $\Gamma(q) = 1 + \ell(S(q)) > j$, then the argument is the same as above. (4) follows from

$$\sum_{q \subset Q: \Gamma(q) \leq \tau} \lambda_q \leq \sum_{q \subset Q: \Gamma'(q) \leq \tau} \lambda_q,$$

and we have Lemma 2.2.

REFERENCES

1. M. Christ and C. D. Sogge, *On the L^1 behavior of eigenfunction expansions and singular integral operators*, in *Miniconferences on harmonic analysis and operator algebras (Canberra, 1987)*, 29–50, Austral. Nat. Univ., Canberra, 1988. MR0953981 (89k:42024)
2. M. Christ, *Weak type $(1, 1)$ bounds for rough operators*, Ann. of Math. (2) **128** (1988), no. 1, 19–42. MR0951506 (89m:42013)
3. G. Mockenhaupt, *A note on the cone multiplier*, Proc. Amer. Math. Soc. **117** (1993), no. 1, 145–152. MR1098404 (93c:42015)
4. G. Garrigós and A. Seeger, *On plate decompositions of cone multipliers*, Proceedings of the Conference on Harmonic Analysis and Its Applications, Hokkaido University, Sapporo (2005).
5. J. Bourgain, *Estimates for cone multipliers*, in *Geometric aspects of functional analysis (Israel, 1992–1994)*, 41–60, Birkhäuser, Basel, 1995. MR1353448 (96m:42022)
6. T. Wolff, *Local smoothing type estimates on L^p for large p* , Geom. Funct. Anal. **10** (2000), no. 5, 1237–1288. MR1800068 (2001k:42030)
7. I. Laba and T. Wolff, *A local smoothing estimate in higher dimensions*, J. Anal. Math. **88** (2002), 149–171. MR1956533 (2005b:35015)
8. S. Lee, *Improved bounds for Bochner-Riesz and maximal Bochner-Riesz operators*, Duke Math. J. **122** (2004), no. 1, 205–232. MR2046812 (2005e:42042)

9. Y. R. Heo, *An endpoint estimate for some maximal operators associated to submanifolds of low codimension*, Pacific J. Math. **201** (2001), no. 2, 323–338. MR1875897 (2002j:42030)
10. Y. Heo, *Improved bounds for high dimensional cone multipliers*, Indiana Univ. Math. J. **58** (2009), no. 3, 1187–1202. MR2541363
11. Y. Heo, F. Nazarov and A. Seeger, *Radial Fourier multipliers in high dimensions*, preprint.
12. Y. Heo, F. Nazarov and A. Seeger, *On radial and conical Fourier multipliers*, preprint.
13. E. M. Stein, *Harmonic analysis: Real-variable methods, orthogonality, and oscillatory integrals*, Princeton Univ. Press, Princeton, NJ, 1993. MR1232192 (95c:42002)

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WISCONSIN-MADISON, MADISON, WISCONSIN 53706

E-mail address: heo@math.wisc.edu

Current address: Pohang Mathematics Institute, Pohang University of Science & Technology, Pohang 790-784, Korea

E-mail address: heo@postech.ac.kr

DEPARTMENT OF MATHEMATICS, CHOSUN UNIVERSITY, GWANGJU 501-759, REPUBLIC OF KOREA

E-mail address: skhong@mail.chosun.ac.kr

DEPARTMENT OF MATHEMATICS, KOREA UNIVERSITY, SEOUL 136-701, REPUBLIC OF KOREA

E-mail address: cw.yang@korea.ac.kr