A NUCLEAR FRÉCHET SPACE CONSISTING OF C^∞ -FUNCTIONS AND FAILING THE BOUNDED APPROXIMATION PROPERTY

DIETMAR VOGT

(Communicated by Nigel J. Kalton)

ABSTRACT. An easy and transparent example is given of a nuclear Fréchet space failing the bounded approximation property and consisting of C^∞ -functions on a subset of $\mathbb{R}^3$.

The problem of Grothendieck whether every nuclear Fréchet space has the bounded approximation property was open for quite a while. The first counterexample is due to Dubinsky [1, 2]. Another much simpler example was given by the author in [7]. Other examples are due to Floret [4] and Moscatelli [6]. In Dubinsky-Vogt [3] it was shown that every infinite dimensional nuclear Fréchet space not isomorphic to $\mathbb{K}^\mathbb{N}$, where $\mathbb{K} = \mathbb{R}$ or $\mathbb{C}$ is the scalar field, has a quotient failing the bounded approximation property.

For a nuclear Fréchet space E the bounded approximation property means the existence of a sequence $(\varphi_n)_{n \in \mathbb{N}}$ of continuous finite rank operators in E so that $\varphi_n x \rightarrow x$ for all $x \in E$.

In [7] the following criterion was established for the bounded approximation property. In [3] it was shown that it is equivalent to E being countably normed. The observation that a non-countably normed Fréchet space admitting a continuous norm will fail the bounded approximation property is due to Pełczyński.

Criterion 1. If E has the bounded approximation property and a continuous norm, then the following holds.

- (*) There exists p_0 such that for every $p \geq p_0$ we have a $q \geq p$ with the following property:

Every sequence in E which is Cauchy with respect to $\|\cdot\|_q$ and convergent to 0 with respect to $\|\cdot\|_{p_0}$ even converges to 0 with respect to $\|\cdot\|_p$.

Here $\|\cdot\|_0 \leq \|\cdot\|_1 \leq \dots$ denotes a fundamental system of continuous seminorms on E .

We use Criterion 1 to develop a scheme, how to construct nuclear Fréchet spaces without the bounded approximation property.

Let F and G be nuclear Fréchet spaces. Let again $\|\cdot\|_0 \leq \|\cdot\|_1 \leq \dots$ denote fundamental systems of continuous seminorms on F and G respectively. We assume that G admits a continuous norm, which we may assume to be $\|\cdot\|_0$. By F_k , $k \in \mathbb{N}_0$ we denote the local Banach space with respect to $\|\cdot\|_k$, i.e. the completion of

Received by the editors January 11, 2009, and, in revised form, August 13, 2009.

2010 *Mathematics Subject Classification.* Primary 46A04; Secondary 46A11.

©2009 American Mathematical Society
 Reverts to public domain 28 years from publication

$(F/\ker \|\cdot\|_k, \|\cdot\|_k)$ and by $j_m^k : F_m \rightarrow F_k$ for $m \geq k$, $j^k : F \rightarrow F_k$ the natural linking maps (see [5, p. 294]).

Let $A : F \supset D(A) \rightarrow G$ be an injective closed linear map; i.e. the graph $\Gamma(A)$ is closed in $F \times G$. We assume that for any p_0 there is $p > p_0$ such that for any $q \geq p$ there is a sequence $(x_n)_n$ in F with the following properties:

- (1) $x_n \rightarrow 0$ with respect to $\|\cdot\|_{p_0}$;
- (2) $Ax_n \rightarrow 0$ with respect to $\|\cdot\|_q$;
- (3) there is $x \in F_q$ with $j_q^p x \neq 0$, so that $j^q x_n \rightarrow x$ in F_q .

Lemma 2. *Then $E := D(A)$ with graph topology given by the norms $\|x\|_k + \|Ax\|_k$, $k \in \mathbb{N}_0$, is a nuclear Fréchet space without the bounded approximation property.*

Proof. E is isomorphic to a closed subspace of $F \times G$, hence a nuclear Fréchet space. Then, because of (2) and (3) the sequence (x_n, Ax_n) , $n \in \mathbb{N}$, is a $\|\cdot\|_q$ -Cauchy sequence. Because of (1) and (2) it is a $\|\cdot\|_{p_0}$ -null sequence. And, because of (2) and (3), it does not converge to 0 with respect to $\|\cdot\|_p$. Since A is injective, E is a normed space. $\square$

We now proceed to the construction of a concrete example. We identify $\mathbb{R}^3 = \mathbb{C} \times \mathbb{R}$ with the variables (z, t) . We apply the lemma to

- (1) $\mathbb{M} = \{(z, t) : |z| \leq 1, |t| < 1\}$,
- (2) $F = \{f \in C^\infty(\mathbb{M}) : f^{(\alpha)}(0, t) = 0, \text{ for all } t \text{ and } \alpha\}$,
- (3) $G = C^\infty(\overline{\mathbb{M}})$,
- (4) $A = \frac{\partial}{\partial \bar{z}}$ with $D(A) = \{f \in F : \frac{\partial f}{\partial \bar{z}} \in G\}$.

Here we identified $C^\infty(\overline{\mathbb{M}})$ with a linear subspace of $C^\infty(\mathbb{M})$. For the nuclearity of F and G we refer to e.g. [5, 28.6 and 28.9] and remark that, by Whitney's extension theorem, $C^\infty(\mathbb{M})$ is a quotient of $C^\infty(\Omega)$, where $\Omega = \{(z, t) \in \mathbb{R}^3 : |t| < 1\}$ is an open subset of $\mathbb{R}^3 = \mathbb{C} \times \mathbb{R}$. A is injective since $\ker A$ consists of functions, which are holomorphic in z for $|z| < 1$ and flat in $(0, t)$ for all t .

Theorem 3. *The space $E = \{f \in F : \frac{\partial f}{\partial \bar{z}} \in G\}$ with its natural Fréchet topology is a nuclear Fréchet space without the bounded approximation property.*

Proof. We may choose the fundamental system of norms

$$\|f\|_k = \sup\{|f^{(\alpha)}(z, t)| : |\alpha| \leq k, |z| \leq 1, |t| \leq t_k\}$$

on F , where $t_k = \frac{k}{k+1}$, and

$$|f|_k = \sup\{|f^{(\alpha)}(z, t)| : |\alpha| \leq k, (z, t) \in \mathbb{M}\}$$

on G . Both spaces are nuclear and clearly A is closed and injective.

Let p_0 be given. We put $p = p_0 + 1$ and assume $q \geq p$.

We choose $\varphi \in \mathcal{D}(\mathbb{D})$, $\mathbb{D}$ the unit disc, such that $\varphi \equiv 1$ in a neighborhood of 0, and $\psi \in \mathcal{D}[t_{p_0}, 1[$ with $\psi(t_p) \neq 0$. We define

$$f_n(z, t) = \psi(t)(1 - \varphi(nz))z^{q+2}.$$

1. $\|f_n\|_{p_0} = 0$ for all n .
2. We have with a suitable constant C :

$$|f_{n, \bar{z}}|_q = |\psi(t)\varphi(nz)\bar{z}z^{q+2}|_q \leq C n^{-1}.$$

That means $|Af_n|_q \rightarrow 0$.

3. With a suitable constant C , we have

$$\|f_n(z, t) - \psi(t) z^{q+2}\|_q = \|\psi(t)\varphi(nz)z^{q+2}\|_q \leq C n^{-2}.$$

That means $f_n \rightarrow \psi z^{q+2} \in F_q$ with respect to $\|\cdot\|_q$ where

$$\|\psi(t) z^{q+2}\|_p \neq 0.$$

The result follows from Lemma 2. $\square$

Let us finally remark that in [8] there was given an easy example of a nuclear Fréchet space consisting of C^∞ -functions which has no basis, namely the space

$$E = \{f \in C^\infty(\mathbb{R}^2) : f|_M \in \mathcal{S}(M)\}$$

with its natural Fréchet topology, where

$$M = \{(x, y) \in \mathbb{R}^2 : x \geq 0, |\sin y| \leq e^{1-\frac{1}{x}}\}.$$

Here $\mathcal{S}(M)$ denotes the space of all C^∞ -functions on M which are rapidly decreasing for $|x| \rightarrow \infty$ with all their derivatives and we set $\exp(-\frac{1}{0}) = 0$. It might be useful to remark that the existence of a basis implies the bounded approximation property. So the present example, being likewise easy, provides much sharper properties.

REFERENCES

1. Dubinsky, E., *The structure of nuclear Fréchet spaces*, Lecture Notes in Mathematics, 720, Berlin-Heidelberg-New York: Springer, 1979. MR537039 (81f:46004)
2. Dubinsky, E., *Nuclear Fréchet spaces without the bounded approximation property*, Studia Math. **71** (1981/82), 85-105. MR651326 (83e:46007)
3. Dubinsky, E., Vogt, D., *Fréchet spaces with quotients failing the bounded approximation property*, Studia Math. **81** (1985), 71-77. MR818172 (87e:46005)
4. Floret, K., *Continuous norms on locally convex strict inductive limit spaces*, Math. Z. **188** (1984), 75-88. MR767364 (86d:46002)
5. Meise, R., Vogt, D., *Introduction to functional analysis*, The Clarendon Press, Oxford Univ. Press, New York, 1997. MR1483073 (98g:46001)
6. Moscatelli, V. B., *Nuclear Fréchet spaces without the bounded approximation property*, Results Math. **19** (1991), 143-146. MR1091962 (92i:46006)
7. Vogt, D., *An example of a nuclear Fréchet space without the bounded approximation property*, Math. Z. **182** (1983), 265-267. MR689304 (84m:46009)
8. Vogt, D., *A nuclear Fréchet space of C^∞ -functions which has no basis*, Note Mat. **25** (2005/06), 187-190. MR2259965 (2007g:46004)

FACHBEREICH C-MATHEMATIK UND NATURWISSENSCHAFTEN, BERGISCHE UNIVERSITÄT WUPPERTAL, D-42097 WUPPERTAL, GERMANY

E-mail address: `dvogt@math.uni-wuppertal.de`