NON-CROSSING LINKED PARTITIONS, THE PARTIAL ORDER $\ll$ ON $NC(n)$, AND THE S -TRANSFORM

ALEXANDRU NICA

(Communicated by Marius Junge)

ABSTRACT. The paper establishes a connection between two recent combinatorial developments in free probability: the non-crossing linked partitions introduced by Dykema in 2007 to study the S -transform, and the partial order $\ll$ on $NC(n)$ introduced by Belinschi and Nica in 2008 in order to study relations between free and Boolean probability. More precisely, one has a canonical bijection between $NCL(n)$ (the set of all non-crossing linked partitions of $\{1, \dots, n\}$) and the set $\{(\alpha, \beta) \mid \alpha, \beta \in NC(n), \alpha \ll \beta\}$. As a consequence of this bijection, one gets an alternative description of Dykema's formula expressing the moments of a non-commutative random variable a in terms of the coefficients of the reciprocal S -transform $1/S_a$. Moreover, due to the Boolean features of $\ll$, this formula can be simplified to a form which resembles the moment-cumulant formula from c -free probability.

1. INTRODUCTION

This paper puts into evidence a connection between two seemingly unrelated combinatorial developments in free probability, which appeared recently in the papers [1] and [5].

The basic structure for combinatorial considerations in free probability is the poset $(NC(n), \leq)$, where $NC(n)$ is the set of all non-crossing partitions of $\{1, \dots, n\}$, and " $\leq$ " is the reverse refinement order for such partitions. $NC(n)$ enters free probability theory via the free cumulants introduced in [10]; a detailed exposition of how this happens can be found in Part II of the monograph [7]. Both papers [1], [5] build on the basic combinatorics of $NC(n)$, but appear to go in different directions, following different goals.

On the one hand, in 2007, Dykema's paper [5] introduced the concept of "non-crossing linked partition" and used it to study the S -transform (an important transform introduced by Voiculescu [12] in order to treat the operation of multiplying free random variables). In particular, it was shown in [5] how the moment of order n of a non-commutative random variable a can be expressed in terms of the coefficients of the series $1/S_a(z)$ via a summation over $NCL(n)$, where $S_a(z)$ is the S -transform of a and where $NCL(n)$ denotes the set of all non-crossing linked

Received by the editors January 29, 2009.

2010 *Mathematics Subject Classification.* Primary 46L54; Secondary 05A18.

Research supported by a Discovery Grant from NSERC, Canada.

©2009 American Mathematical Society
 Reverts to public domain 28 years from publication

partitions of $\{1, \dots, n\}$. Following up [5], further work related to $NCL(n)$ was done in [4], [8]. A review of $NCL(n)$ and of its connection to the S -transform is given in Section 2B and Section 4 below.

On the other hand, in 2008, the paper [1] by Belinschi and Nica introduced a partial order “ $\ll$ ” on $NC(n)$, coarser than the reverse refinement order. The partial order $\ll$ was used in [1] and subsequently in [2] in order to study connections between free probability and Boolean probability (another brand of non-commutative probability, where instead of the $NC(n)$ ’s one works with Boolean posets). A useful fact which illustrates how $\ll$ mixes together features from the non-crossing and the Boolean worlds is the following: for a fixed $\beta \in NC(n)$, the set $\{\alpha \in NC(n) \mid \alpha \ll \beta\}$ can be identified as an interval with respect to the reverse refinement order, and is consequently counted by a product of Catalan numbers. But for a fixed $\alpha \in NC(n)$, the set $\{\beta \in NC(n) \mid \beta \gg \alpha\}$ is isomorphic to a Boolean poset and is thus counted by a power of 2. (See the review in Propositions 2.4 and 2.5 below.)

The present paper puts into evidence a canonical bijection from $NCL(n)$ onto the set $\{(\alpha, \beta) \mid \alpha, \beta \in NC(n), \alpha \ll \beta\}$. If $\pi \in NCL(n)$ is mapped by this bijection to (α, β) , then:

- β is the partition generated by the blocks of π . (Note that the blocks of a linked partition are allowed, under certain special circumstances, to not be disjoint. Hence π does not have to be itself a partition, and it may happen that $\beta \neq \pi$.)
- α is obtained by applying a certain “cycling” procedure to a partition $\check{\pi} \in NC(n)$, which is defined in [5] and is called the “unlinking of π ”.

The precise details of how this works are given in Theorem 3.4 of this paper.

As a consequence of the above bijection, one gets an alternative description of Dykema’s formula expressing the moments of a non-commutative random variable a in terms of the coefficients of the reciprocal S -transform $1/S_a$. Moreover, due to the Boolean features of $\ll$, this formula can be simplified to a form which expresses the moment of order n of a by using a summation over $NC(n)$ (instead of $NCL(n)$), and where the term indexed by $\alpha \in NC(n)$ in the summation can still be canonically written as a product over the set of blocks of α . The precise description of how this goes is given in Theorem 4.5 below. It is worth noting that equation (4.27) from Theorem 4.5 has a close resemblance to the moment-cumulant formula developed in [3] for the framework of conditionally free random variables (another framework where the free and Boolean probability worlds interact).

Besides the introduction, the paper has three other sections. Section 2 contains a review of the background and notation, Section 3 presents the canonical bijection advertised above, and Section 4 examines the application of this bijection to the reciprocal S -transform.

2. BACKGROUND AND NOTATION

2A. Non-crossing partitions and the partial order $\ll$. We will use the standard conventions of notation for non-crossing partitions (as in [9], or in Lecture 9 of [7]). The partial order given by reverse refinement on $NC(n)$ will be simply denoted as “ $\leq$ ”; in other words, for $\alpha, \beta \in NC(n)$, we write “ $\alpha \leq \beta$ ” to mean that every block of β is a union of blocks of α . The minimal and maximal elements of $(NC(n), \leq)$ are denoted by 0_n (the partition of $\{1, \dots, n\}$ into n blocks of 1 element each) and respectively 1_n (the partition of $\{1, \dots, n\}$ into 1 block of n elements).

The coarser partial order $\ll$ on $NC(n)$ was defined in [1] as follows.

Definition 2.1. For $\alpha, \beta \in NC(n)$ we write “ $\alpha \ll \beta$ ” to mean that $\alpha \leq \beta$ and that, in addition, the following condition is fulfilled:

$$(2.1) \quad \begin{cases} \text{For every block } W \text{ of } \beta \text{ there exists a block} \\ V \text{ of } \alpha \text{ such that } \min(W), \max(W) \in V. \end{cases}$$

It is immediate that if $\alpha \leq \beta$ in $NC(n)$ and if V, W are as in (2.1), then one must have $V \subseteq W$ and $\min(V) = \min(W)$, $\max(V) = \max(W)$. It will be convenient to give a special name to the blocks V of α that can be matched to a block W of β in this way.

Definition 2.2. Let α, β be partitions in $NC(n)$ such that $\alpha \ll \beta$. A block V of α will be said to be β -special when there exists a block W of β such that $\min(V) = \min(W)$ and $\max(V) = \max(W)$.

Remark 2.3. Let $\alpha, \beta \in NC(n)$ be such that $\alpha \ll \beta$.

1. The correspondence $V \mapsto W$ from Definition 2.2 clearly gives a bijection from the set $\{V \text{ block of } \alpha \mid V \text{ is } \beta\text{-special}\}$ onto the set of all blocks of β .

2. Let us recall that a block V of α is said to be *inner* (respectively *outer*) when there exists (respectively when there does not exist) another block V' of α such that $\min(V') < \min(V)$ and $\max(V') > \max(V)$. It is easily seen that every outer block of α is β -special; moreover, the correspondence $V \mapsto W$ from Definition 2.2 induces a bijection between the outer blocks of α and those of β ; cf. Remarks 2.9 and 2.12 in [1].

The next two propositions state in more detail the facts mentioned in the introduction about sets of the form $\{\alpha \in NC(n) \mid \alpha \ll \beta\}$ and $\{\beta \in NC(n) \mid \beta \gg \alpha\}$.

Proposition 2.4. Let $\beta = \{W_1, \dots, W_q\}$ be a partition in $NC(n)$. Consider the partition $\beta_0 \in NC(n)$ obtained by refining β as follows: every block W_j with $|W_j| \leq 2$ is left intact, while every block W_j with $|W_j| \geq 3$ is broken into the doubleton $\{\min(W_j), \max(W_j)\}$ and $|W_j| - 2$ singletons. (Thus every block of β_0 has either 1 or 2 elements.) Then

$$(2.2) \quad \{\alpha \in NC(n) \mid \alpha \ll \beta\} = \{\alpha \in NC(n) \mid \beta_0 \leq \alpha \leq \beta\}.$$

As a consequence, one has that

$$(2.3) \quad |\{\alpha \in NC(n) \mid \alpha \ll \beta\}| = \prod_{j=1}^q \text{Cat}_{|W_j|-1},$$

where, for every $k \geq 0$, we denote $\text{Cat}_k := (2k)!/(k!(k+1)!)$ (the k th Catalan number).

Proof. The equality in (2.2) follows immediately from how the partial order $\ll$ is defined. The right-hand side of (2.2) is the interval $[\beta_0, \beta]$ with respect to the reverse refinement order, and every interval of $(NC(n), \leq)$ is known to be canonically isomorphic (as a poset) to a direct product of lattices $NC(m)$, $2 \leq m \leq n$; see the detailed discussion on pp. 149-150 of [7], which also gives a concrete algorithm for how to obtain the canonical factorization of the interval. By following this

algorithm it is immediately found that

$$(2.4) \quad [\beta_0, \beta] \simeq \prod_{\substack{1 \leq j \leq q \text{ such} \\ \text{that } |W_j| \geq 3}} NC(|W_j| - 1),$$

and (2.3) follows by taking cardinalities in (2.4). $\square$

The next proposition uses the abbreviations $V \in \alpha$ for “ V is a block of α ” and $\mathcal{V} \subseteq \alpha$ for “ $\mathcal{V}$ is a set of blocks of α ”, where α is a partition in $NC(n)$. For the proof of this proposition, the reader is referred to Proposition 2.13 and Remark 2.14 of [1].

Proposition 2.5. *Let α be in $NC(n)$ and consider the set of partitions*

$$(2.5) \quad \{\beta \in NC(n) \mid \beta \gg \alpha\}.$$

Then $\beta \mapsto \{V \in \alpha \mid V \text{ is } \beta\text{-special}\}$ is a one-to-one map from the set (2.5) to the set of subsets of α . The image of this map is equal to $\{\mathcal{V} \subseteq \alpha \mid \mathcal{V} \text{ contains all outer blocks of } \alpha\}$.

2B. Linked partitions and $NCL(n)$. Following [5], we will use the term “*linked partition* of $\{1, \dots, n\}$ ” for a set $\pi = \{A_1, \dots, A_p\}$ of non-empty subsets of $\{1, \dots, n\}$ such that $A_1 \cup \dots \cup A_p = \{1, \dots, n\}$, and where for every $i \neq j$ ($1 \leq i, j \leq p$) one has that either $A_i \cap A_j = \emptyset$ or that the following holds:

$$(2.6) \quad \begin{cases} |A_i| \geq 2, |A_j| \geq 2, |A_i \cap A_j| = 1, \min(A_i) \neq \min(A_j), \text{ and} \\ \text{the unique element of } A_i \cap A_j \text{ is one of } \min(A_i), \min(A_j). \end{cases}$$

Obviously, every partition of $\{1, \dots, n\}$ (in the usual sense of the term) is a linked partition of $\{1, \dots, n\}$, but the converse is not true. Throughout this paper we use the letters $\alpha, \beta, \dots$ to denote partitions, and $\pi, \rho, \dots$ to denote linked partitions (which may or may not be partitions). A few more terms and basic facts from Section 5 of [5] are reviewed next.

Review 2.6. 1. If $\pi = \{A_1, \dots, A_p\}$ is a linked partition of $\{1, \dots, n\}$, then $A_1, \dots, A_p$ are called the *blocks* of π . It is easy to see that every $m \in \{1, \dots, n\}$ belongs either to exactly one or to exactly two blocks A_i . In the first case one says that m is *singly-covered* by π , and in the second case one says that m is *doubly-covered* by π .

2. Let π be a linked partition of $\{1, \dots, n\}$. The partition of $\{1, \dots, n\}$ which is generated by π will be denoted as $\hat{\pi}$. In other words, $\hat{\pi}$ is the smallest (with respect to the reverse refinement order) among all partitions β of $\{1, \dots, n\}$ which have the following property: “for every block A of π there exists a block V of β such that $A \subseteq V$ ”.

3. Let $\pi = \{A_1, \dots, A_p\}$ be a linked partition of $\{1, \dots, n\}$. For every $1 \leq i \leq p$ define

$$(2.7) \quad V_i = \begin{cases} A_i, & \text{if } \min(A_i) \text{ is singly-covered by } \pi, \\ A_i \setminus \{\min(A_i)\}, & \text{if } \min(A_i) \text{ is doubly-covered by } \pi. \end{cases}$$

Then $\{V_1, \dots, V_p\}$ is a partition of $\{1, \dots, n\}$ called the *unlinking* of π and denoted as $\tilde{\pi}$.

4. A linked partition π of $\{1, \dots, n\}$ is said to be *non-crossing* if it is not possible to find two distinct blocks A, B of π and elements $a, a' \in A$, $b, b' \in B$ such that $a < a' < b < b'$.

5. Let n be a positive integer. The set of all non-crossing linked partitions of $\{1, \dots, n\}$ is denoted by $NCL(n)$. It is not hard to see that if $\pi \in NCL(n)$, then the partitions $\hat{\pi}$ and $\check{\pi}$ defined in 2 and 3 above belong to $NC(n)$.

Remark 2.7 (Restrictions of linked partitions). It is immediate that the above discussion about linked partitions could be carried out without any modifications in the larger framework where instead of $\{1, \dots, n\}$ one uses an abstract finite totally ordered set F , and one considers the set $NCL(F)$ of non-crossing linked partitions of F (instead of just sticking to the $NCL(n)$). This doesn't really bring anything new, since it is obvious that $NCL(F)$ is canonically identified to $NCL(|F|)$ via the map $\{A_1, \dots, A_p\} \mapsto \{f(A_1), \dots, f(A_p)\}$, where f is the unique order-preserving bijection from F onto $\{1, \dots, |F|\}$. But in the subsequent discussion it will be nevertheless convenient to allow linked partitions for a slightly more general kind of set F (specifically, for F being a non-empty finite subset of $\mathbb{N}$), in order to simplify the notation for restrictions of linked partitions.

So let $E \subseteq F$ be non-empty subsets of $\mathbb{N}$, and let $\pi \in NCL(F)$ be such that E is saturated with respect to π (which means that whenever A is a block of π and $A \cap E \neq \emptyset$, it follows that $A \subseteq E$). Then π is of the form $\{A_1, \dots, A_p, A'_1, \dots, A'_q\}$ with $A_1, \dots, A_p \subseteq E$ and $A'_1, \dots, A'_q \subseteq F \setminus E$, and one defines the *restriction* of π to E to be

$$(2.8) \quad \pi|_E := \{A_1, \dots, A_p\}.$$

It is immediate that $\pi|_E \in NCL(E)$. Moreover, it is easily seen that the operation of restriction is well-behaved with respect to the maps $\pi \mapsto \hat{\pi}$ and $\pi \mapsto \check{\pi}$ from 2 and 3 of Review 2.6, in the sense that one has

$$(2.9) \quad \left(\pi|_E\right)^{\wedge} = (\hat{\pi})|_E \quad \text{and} \quad \left(\pi|_E\right)^{\vee} = (\check{\pi})|_E.$$

Let us next record the observation (cf. [5], Corollary 5.13 and its proof) that non-crossing linked partitions can be broken into “irreducible” pieces, as follows.

Proposition 2.8. *Let n be a positive integer and let $\beta = \{W_1, \dots, W_q\}$ be a partition in $NC(n)$. The map*

$$\pi \mapsto \left(\pi|_{W_1}, \dots, \pi|_{W_q}\right)$$

is a bijection from $\{\pi \in NCL(n) \mid \hat{\pi} = \beta\}$ onto $\prod_{j=1}^q \{\pi_j \in NCL(W_j) \mid \hat{\pi}_j = 1_{W_j}\}$ (where $1_{W_j} \in NC(W_j)$ is the partition of W_j into only one block).

So, after suitably renumbering every W_j from the preceding proposition as $\{1, \dots, |W_j|\}$, one is reduced in the end to looking at sets of non-crossing linked partitions of the form $\{\pi \in NCL(m) \mid \hat{\pi} = 1_m\}$. In Proposition 5.11 of [5] it is pointed out that the latter sets can be identified with sets of usual non-crossing partitions, as follows.

Proposition 2.9. *For every $n \geq 2$, the map $\pi \mapsto \check{\pi}$ is a bijection from $\{\pi \in NCL(n) \mid \hat{\pi} = 1_n\}$ onto $\{\alpha \in NC(n) \mid 1 \text{ and } 2 \text{ belong to the same block of } \alpha\}$.*

3. THE CANONICAL BIJECTION RELATING $NCL(n)$ TO $\ll$

This section is devoted to proving the bijection announced in the introduction. The main result is Theorem 3.4. We start by recalling some connections between set-partitions and permutations and by defining precisely what is the “cycled un-linking” of a linked partition $\pi \in NCL(n)$.

Notation 3.1. 1. If τ is a permutation of $\{1, \dots, n\}$ and if $\alpha = \{V_1, \dots, V_p\}$ is a partition of $\{1, \dots, n\}$, then we denote

$$(3.10) \quad \tau \cdot \alpha := \{\tau(V_1), \dots, \tau(V_p)\} \quad (\text{a new partition of } \{1, \dots, n\}).$$

It is clear that formula (3.10) defines an action of the symmetric group $\mathcal{S}_n$ on the set of all partitions of $\{1, \dots, n\}$.

2. Every partition $\alpha \in NC(n)$ has associated to it a permutation of $\{1, \dots, n\}$, which is denoted by P_α and is defined by the following prescription: for every block $V = \{i_1, \dots, i_m\}$ of π , with $i_1 < \dots < i_m$, one creates a cycle of P_α by putting

$$P_\alpha(i_1) = i_2, \dots, P_\alpha(i_{m-1}) = i_m, P_\alpha(i_m) = i_1.$$

Definition 3.2. Let π be in $NCL(n)$. Consider the partitions $\hat{\pi}, \check{\pi} \in NC(n)$ (as in Review 2.6) and form the new partition

$$(3.11) \quad \overset{\circ}{\pi} := P_{\hat{\pi}}^{-1} \cdot \check{\pi}$$

(where the permutation $P_{\hat{\pi}}$ is defined as in Notation 3.1.2). The partition $\overset{\circ}{\pi}$ will be called the *cycled unlinking* of π .

Example 3.3 (A concrete example). Say for instance that $n = 11$ and that

$$\pi = \{ \{1, 2, 4\}, \{2, 3\}, \{4, 5, 6\}, \{6, 7\}, \{8, 9, 11\}, \{9, 10\} \} \in NCL(11).$$

Then one has

$$\hat{\pi} = \{ \{1, 2, 3, 4, 5, 6, 7\}, \{8, 9, 10, 11\} \}, \quad \check{\pi} = \{ \{1, 2, 4\}, \{3\}, \{5, 6\}, \{7\}, \{8, 9, 11\}, \{10\} \}.$$

The permutation associated to $\hat{\pi}$ is

$$P_{\hat{\pi}} = (1, 2, 3, 4, 5, 6, 7) (8, 9, 10, 11) \quad (\text{written in cycle notation});$$

hence the cycled unlinking of π is

$$\overset{\circ}{\pi} = P_{\hat{\pi}}^{-1} \cdot \check{\pi} = \{ \{1, 3, 7\}, \{2\}, \{4, 5\}, \{6\}, \{8, 10, 11\}, \{9\} \}.$$

Note that $\overset{\circ}{\pi} \in NC(11)$ and $\overset{\circ}{\pi} \ll \hat{\pi}$ (we will see that this must generally be the case).

Theorem 3.4. Let n be a positive integer. The map $\pi \mapsto (\overset{\circ}{\pi}, \hat{\pi})$ is a bijection from $NCL(n)$ onto the set $\{(\alpha, \beta) \mid \alpha, \beta \in NC(n), \alpha \ll \beta\}$.

For the proof of Theorem 3.4, it is convenient to first establish some lemmas.

Lemma 3.5. Let β be a partition in $NC(n)$. The map $\alpha \mapsto P_\beta^{-1} \cdot \alpha$ sends the set $\{\alpha \in NC(n) \mid \alpha \leq \beta\}$ into itself.

Proof. Let us fix an $\alpha \in NC(n)$ such that $\alpha \leq \beta$, and let us consider the partition $\alpha' := P_\beta^{-1} \cdot \alpha$. Observe that $\alpha' \leq \beta$ in reverse refinement order, i.e. that every block V' of α' is contained in a block of β . Indeed, V' can be written as $V' = P_\beta^{-1}(V)$, where V is a block of α , and the block W of β which contains V must contain V' as well (since $V' \subseteq P_\beta^{-1}(W) = W$).

It remains to show that the partition α' is non-crossing. Let us fix two distinct blocks V'_1 and V'_2 of α' , and let us prove that V'_1 and V'_2 do not cross. Consider the blocks W_1, W_2 of β such that $W_1 \supseteq V'_1$ and $W_2 \supseteq V'_2$, and distinguish the following two cases.

Case 1. $W_1 \neq W_2$. In this case, W_1 and W_2 do not cross, so the smaller sets V'_1 and V'_2 can't cross either.

Case 2. $W_1 = W_2 =: W$. In this case, consider the sets $V_1 := P_\beta(V'_1)$ and $V_2 := P_\beta(V'_2)$. These are two distinct blocks of α , which are still contained in W . We know that V_1 and V_2 don't cross (since α is in $NC(n)$), and we will conclude the proof by arguing that if V'_1 and V'_2 would cross, then so would V_1 and V_2 . Indeed, suppose that $i_1, j_1 \in V'_1$ and $i_2, j_2 \in V'_2$ are such that $i_1 < i_2 < j_1 < j_2$. If j_2 is not the maximal element of W , then it follows that $P_\beta(i_1) < P_\beta(i_2) < P_\beta(j_1) < P_\beta(j_2)$, which is a crossing between V_1 and V_2 . In the opposite situation when $j_2 = \max(W)$, it follows that $P_\beta(j_2) < P_\beta(i_1) < P_\beta(i_2) < P_\beta(j_1)$, and we get a crossing between V_2 and V_1 (since $P_\beta(j_2), P_\beta(i_2) \in V_2$ and $P_\beta(i_1), P_\beta(j_1) \in V_1$). $\square$

Lemma 3.6. *Let π be in $NCL(n)$. Then the partition $\overset{\circ}{\pi}$ is in $NC(n)$, and $\overset{\circ}{\pi} \ll \hat{\pi}$.*

Proof. Since $\check{\pi}, \hat{\pi} \in NC(n)$ and $\check{\pi} \leq \hat{\pi}$, the preceding lemma gives us that $\overset{\circ}{\pi} = P_{\hat{\pi}}^{-1} \cdot \check{\pi}$ is in $NC(n)$ and that $\overset{\circ}{\pi} \leq \hat{\pi}$.

It remains to show that, for every block W of $\hat{\pi}$, the numbers $\min(W)$ and $\max(W)$ belong to the same block of $\overset{\circ}{\pi}$. For the rest of the proof we fix such a W , with $|W| =: m \geq 2$ (the case $|W| = 1$ is obvious). Let us write explicitly $W = \{i_1, i_2, \dots, i_m\}$, with $i_1 < i_2 < \dots < i_m$; our goal is then to prove that i_1 and i_m belong to the same block of $\overset{\circ}{\pi}$.

Consider the linked partition $\pi|_W \in NCL(W)$, which has $(\pi|_W)^\wedge = \hat{\pi}|_W = 1_W$. Proposition 2.9 gives us (after going through the suitable identification of $i_1, i_2, \dots, i_m$ with $1, 2, \dots, m$) that i_1 and i_2 belong to the same block of $(\pi|_W)^\vee$. But $(\pi|_W)^\vee = \check{\pi}|_W$ (cf. equation (2.9)); hence it follows that i_1 and i_2 belong to the same block V of $\check{\pi}$. This implies in turn that $P_{\hat{\pi}}^{-1}(i_1)$ and $P_{\hat{\pi}}^{-1}(i_2)$ belong to the same block $P_{\hat{\pi}}^{-1}(V)$ of $\overset{\circ}{\pi}$, but $P_{\hat{\pi}}^{-1}(i_1) = i_m$ and $P_{\hat{\pi}}^{-1}(i_2) = i_1$, so we are done. $\square$

Lemma 3.7. *Let $\pi, \rho \in NCL(n)$ be such that $\hat{\pi} = \hat{\rho}$ and $\check{\pi} = \check{\rho}$. Then $\pi = \rho$.*

Proof. Let us denote $\hat{\pi} = \hat{\rho} =: \beta \in NC(n)$, and let us write explicitly $\beta = \{W_1, \dots, W_q\}$. In order to prove that $\pi = \rho$ it suffices to verify that they have the same image by the bijection from Proposition 2.8, i.e. that $\pi|_{W_j} = \rho|_{W_j} \in NCL(W_j)$ for every $1 \leq j \leq q$.

So let us fix j ($1 \leq j \leq q$). Observe that $(\pi|_{W_j})^\wedge = \hat{\pi}|_{W_j} = \beta|_{W_j} = 1_{W_j}$, and similarly $(\rho|_{W_j})^\wedge = 1_{W_j}$. Now, by invoking Proposition 2.9 (and by using the suitable renumbering of W_j into $\{1, \dots, |W_j|\}$) we see that the map $\theta \mapsto \check{\theta}$ is one-to-one on $\{\theta \in NCL(W_j) \mid \hat{\theta} = 1_{W_j}\}$. Thus the required fact that $\pi|_{W_j} = \rho|_{W_j}$ will follow if we can prove that $(\pi|_{W_j})^\vee = (\rho|_{W_j})^\vee$. But the latter equality amounts (in view of (2.9)) to $\check{\pi}|_{W_j} = \check{\rho}|_{W_j}$, and thus it follows from the hypothesis that $\check{\pi} = \check{\rho}$. $\square$

Proof of Theorem 3.4. From Lemma 3.6 it follows that the map $\pi \mapsto (\overset{\circ}{\pi}, \hat{\pi})$ is well-defined (with target set as described in the theorem). In order to prove the injectivity of this map, consider two linked partitions $\pi, \rho \in NCL(n)$ such that $\overset{\circ}{\pi} = \overset{\circ}{\rho} =: \alpha$ and $\hat{\pi} = \hat{\rho} =: \beta$. Then $\check{\pi} = \check{\rho} = P_\beta \cdot \alpha$; hence π and ρ satisfy the hypotheses of Lemma 3.7, and it follows that $\pi = \rho$.

In order to complete the proof of the theorem, it now suffices to verify that the sets $NCL(n)$ and $\{(\alpha, \beta) \mid \alpha, \beta \in NC(n), \alpha \ll \beta\}$ have the same cardinality. For

$NCL(n)$ we can invoke Corollary 5.13 of [5], which gives us the formula

$$(3.12) \quad |NCL(n)| = \sum_{\beta \in NC(n)} \left(\prod_{W \in \beta} \text{Cat}_{|W|-1} \right).$$

But on the other hand we have

$$|\{(\alpha, \beta) \mid \alpha, \beta \in NC(n), \alpha \ll \beta\}| = \sum_{\beta \in NC(n)} |\{\alpha \mid \alpha \in NC(n), \alpha \ll \beta\}|,$$

which is indeed equal to the right-hand side of (3.12), due to Proposition 2.4. $\square$

Remark 3.8. Theorem 3.4 allows one to transfer enumeration properties in between the two sets involved in the bijection of the theorem. In particular, as it is known that $NCL(n)$ is counted by the $(n-1)$ th Schröder number r_{n-1} (Theorem 8.3 in [5]; see also the discussion in Section 2 of [4]), it follows that the same is true for $\{(\alpha, \beta) \mid \alpha, \beta \in NC(n), \alpha \ll \beta\}$. By writing the latter set as $\bigcup_{\alpha} \{\beta \in NC(n) \mid \beta \gg \alpha\}$ and by invoking Proposition 2.5, one thus comes to the following amusing (possibly already known) enumerative interpretation of the Schröder number r_{n-1} : it counts coloured non-crossing partitions $\alpha \in NC(n)$, where every block of α is coloured in either red or blue, and all the outer blocks are red. For a nice collection of other enumerative interpretations of the Schröder numbers, see Example 6.2.8 and Exercise 6.39 of the monograph [11].

4. APPLICATION TO THE RECIPROCAL S -TRANSFORM

Let $(\mathcal{A}, \varphi)$ be an algebraic non-commutative probability space (which simply means that $\mathcal{A}$ is a unital algebra over $\mathbb{C}$ and $\varphi : \mathcal{A} \rightarrow \mathbb{C}$ is a linear functional such that $\varphi(1_{\mathcal{A}}) = 1$), and let $a \in \mathcal{A}$ be such that $\varphi(a) = 1$. The S -transform of a is the series

$$(4.13) \quad S_a(z) := \frac{1+z}{z} M_a^{\langle -1 \rangle}(z) \in \mathbb{C}[[z]],$$

where $M_a(z) := \sum_{n=1}^{\infty} \varphi(a^n) z^n$ (the *moment series* of a), and where $M_a^{\langle -1 \rangle}$ is the inverse of M_a under composition. The S -transform plays a fundamental role in the study of multiplication of free random variables (see Section 3.6 of [13] or Lecture 18 of [7] for the details of how this goes).

It is immediate that the series $S_a(z)$ from (4.13) has constant term equal to 1. Hence one can consider its reciprocal (i.e. inverse under multiplication) $1/S_a(z)$, which is another series with constant term equal to 1:

$$(4.14) \quad 1/S_a(z) = \sum_{n=0}^{\infty} t_n z^n, \quad \text{with } t_0 = 1.$$

In the paper [5] the series $1/S_a$ goes under the name of “ T -transform of a ”. It is observed there that the formula giving back the family of moments $\{\varphi(a^n) \mid n \geq 1\}$ of a in terms of the family $\{t_n \mid n \geq 0\}$ of coefficients of $1/S_a$ only uses positive integer coefficients:

$$(4.15) \quad \begin{cases} \varphi(a) = 1, \quad \varphi(a^2) = t_1 + 1, \quad \varphi(a^3) = t_2 + t_1^2 + 3t_1 + 1, \\ \varphi(a^4) = t_3 + 3t_2t_1 + t_1^3 + 4t_2 + 6t_1^2 + 6t_1 + 1, \dots \end{cases}$$

Moreover, [5] identifies precisely the combinatorial structure which indexes the sums in (4.15): this is nothing but $NCL(n)$, and the formula generalizing the special cases

from (4.15) is

$$(4.16) \quad \varphi(a^n) = \sum_{\pi \in NCL(n)} \left(\prod_{A \in \pi} t_{|A|-1} \right), \quad n \geq 1,$$

where “ $A \in \pi$ ” is an abbreviation for “ A is a block of π ”; see Proposition 8.1 in [5].

In view of Theorem 3.4 of the present paper, the formula giving $\varphi(a^n)$ in terms of the coefficients of $1/S_a$ can be equivalently understood as a summation over the set $\{(\alpha, \beta) \mid \alpha, \beta \in NC(n), \alpha \ll \beta\}$. This alternative version of the formula will be stated in Proposition 4.3 below. In preparation of that statement, we prove two lemmas.

Lemma 4.1. *Let π be a linked partition in $NCL(n)$ such that $\hat{\pi} = 1_n$ and let A be a block of π such that $A \not\# 1$. Then $\min(A)$ is doubly-covered by π .*

Proof. Denote $\min(A) =: m$. Since 1 and m belong to the same (unique) block of $\hat{\pi}$, there have to exist $p \geq 1$, some $m_0, m_1, \dots, m_p \in \{1, \dots, n\}$ and some blocks $A_1, \dots, A_p$ of π such that $m_0 = 1$, $m_p = m$, and such that

$$(4.17) \quad m_0, m_1 \in A_1; m_1, m_2 \in A_2; \dots, m_{p-1}, m_p \in A_p.$$

Let us suppose moreover that in (4.17) p is picked to be as small as possible. It is then immediate that $m_{j-1} \neq m_j$ for every $1 \leq j \leq p$, and that $A_{j-1} \neq A_j$ for every $2 \leq j \leq p$.

Observe that $m_1 \in A_1 \cap A_2$ and m_1 is not the minimum of A_1 (since $m_0 \in A_1$, and $m_0 = 1 < m_1$); from the definition of a linked partition it follows that $m_1 = \min(A_2)$. We next observe that $m_2 \in A_2 \cap A_3$ and $m_2 \neq \min(A_2)$ (since $\min(A_2) = m_1 \neq m_2$), so the same argument as above applies to give us that $m_2 = \min(A_3)$. Continuing like this by induction we find that $m_j = \min(A_{j+1})$ for every $1 \leq j \leq p-1$, and in particular that $m_{p-1} = \min(A_p)$.¹ Thus the block A_p of π contains $m_p = m$, and A_p is different from A (because $\min(A_p) = m_{p-1} \neq m_p = \min(A)$). This shows that m is doubly-covered by π , as required. $\square$

Lemma 4.2. *Let π be a linked partition in $NCL(n)$, let A be a block of π , and let W be the unique block of $\hat{\pi}$ such that $W \supseteq A$. Then*

$$(4.18) \quad \left(\begin{array}{c} \min(A) \text{ is} \\ \text{singly-covered by } \pi \end{array} \right) \Leftrightarrow \left(\min(A) = \min(W) \right).$$

Proof. By replacing π with $\pi|_W$ and by redenoting the elements of W as $\{1, 2, \dots, |W|\}$ in increasing order, we may assume without loss of generality that $\hat{\pi} = 1_n$ and hence that $W = \{1, \dots, n\}$. The statement on the right-hand side of equivalence (4.18) becomes “ $\min(A) = 1$ ”. The implication “ $\Leftarrow$ ” in this equivalence is then immediate (as the definition of a linked partition implies that 1 always is singly-covered), while the implication “ $\Rightarrow$ ” follows from Lemma 4.1. $\square$

Proposition 4.3. *Let $(\mathcal{A}, \varphi)$ be an algebraic non-commutative probability space, let $a \in \mathcal{A}$ be such that $\varphi(a) = 1$, and consider the reciprocal S -transform $1/S_a(z) =$*

¹ This argument was run by assuming that $p \geq 2$. If $p = 1$, then the conclusion that $m_{p-1} = \min(A_p)$ still holds, as we must have that $m_0 = 1 = \min(A_1)$.

$\sum_{n=0}^{\infty} t_n z^n$. Then for every $n \geq 1$ one has

$$(4.19) \quad \varphi(a^n) = \sum_{\substack{\alpha, \beta \in NC(n) \\ \text{such that } \alpha \ll \beta}} \left(\prod_{\substack{U \in \alpha, \\ \beta\text{-special}}} t_{|U|-1} \right) \cdot \left(\prod_{\substack{V \in \alpha, \\ \text{not } \beta\text{-special}}} t_{|V|} \right)$$

(where the concept of a β -special block of α is as in Definition 2.2).

Proof. We will verify that the sums on the right-hand sides of (4.16) and (4.19) are identified term by term when one uses the bijection $\pi \leftrightarrow (\alpha, \beta)$ from Theorem 3.4. We thus fix for the whole proof $\pi \in NCL(n)$ and $\alpha, \beta \in NC(n)$ such that $\pi \leftrightarrow (\alpha, \beta)$ in Theorem 3.4, and our goal is to show that

$$(4.20) \quad \prod_{A \in \pi} t_{|A|-1} = \left(\prod_{\substack{U \in \alpha, \\ \beta\text{-special}}} t_{|U|-1} \right) \cdot \left(\prod_{\substack{V \in \alpha, \\ \text{not } \beta\text{-special}}} t_{|V|} \right).$$

The fact that $\pi \leftrightarrow (\alpha, \beta)$ in Theorem 3.4 means of course that $\alpha = \overset{\circ}{\pi}$ and $\beta = \hat{\pi}$. Let us write explicitly $\beta = \{W_1, \dots, W_q\}$. In view of Lemma 4.2 we see that π can then be written in the form $\pi = \{A_1, \dots, A_q, B_1, \dots, B_r\}$, where $\min(A_1) = \min(W_1), \dots, \min(A_q) = \min(W_q)$ and where $\min(A_1), \dots, \min(A_q)$ are singly-covered by π , while $\min(B_1), \dots, \min(B_r)$ are doubly-covered by π . From how the unlinking $\tilde{\pi}$ is defined (cf. Review 2.6.3) we next infer that

$$\begin{cases} \tilde{\pi} = \{U_1, \dots, U_q, V_1, \dots, V_r\}, & \text{where} \\ U_j = A_j, \forall 1 \leq j \leq q \text{ and } V_k = B_k \setminus \{\min(B_k)\}, \forall 1 \leq k \leq r. \end{cases}$$

From Definition 3.2 we further infer that the partition $\alpha = \overset{\circ}{\pi}$ can be written in the form

$$\begin{cases} \overset{\circ}{\pi} = \{U'_1, \dots, U'_q, V'_1, \dots, V'_r\}, & \text{where} \\ U'_j = P_{\beta}^{-1}(U_j), \forall 1 \leq j \leq q \text{ and } V'_k = P_{\beta}^{-1}(V_k), \forall 1 \leq k \leq r. \end{cases}$$

We next observe that

$$(4.21) \quad \min(W_j), \max(W_j) \in U'_j, \quad \forall 1 \leq j \leq q.$$

Indeed, this statement is clear in the case $|W_j| = 1$ (when one has $W_j = U_j = U'_j$). In the case when $|W_j| =: m \geq 2$ we write $W_j = \{i_1, i_2, \dots, i_m\}$ with $i_1 < i_2 < \dots < i_m$ and we observe that the argument used in the proof of Lemma 3.6 applies, giving us that $i_1, i_2 \in U_j$, and hence that $i_1, i_m \in U'_j$.

From (4.21) it follows that the β -special blocks of α are precisely $U'_1, \dots, U'_q$. The right-hand side of (4.20) thus takes the form

$$(4.22) \quad \left(\prod_{j=1}^q t_{|U'_j|-1} \right) \cdot \left(\prod_{k=1}^r t_{|V'_k|} \right).$$

But it is clear that $|U'_j| = |U_j| = |A_j|$, $1 \leq j \leq q$, and that $|V'_k| = |V_k| = |B_k| - 1$, $1 \leq k \leq r$. Hence the product from (4.22) equals $\left(\prod_{j=1}^q t_{|A_j|-1} \right) \cdot \left(\prod_{k=1}^r t_{|B_k|-1} \right)$, and (4.20) follows. $\square$

It is in fact fairly easy, in hindsight, to give a direct proof of Proposition 4.3 by using the R -transform R_a (another important transform of free probability) as an intermediate for passing from the series $1/S_a$ to the moments of a .

Second proof of Proposition 4.3. Let us consider the R -transform of a . This is the series $R_a(z) = \sum_{n=1}^{\infty} \kappa_n z^n$ whose coefficients $(\kappa_n)_{n=1}^{\infty}$, called the *free cumulants* of a , are uniquely determined by the fact that they satisfy the relations

$$(4.23) \quad \varphi(a^n) = \sum_{\beta \in NC(n)} \left(\prod_{W \in \beta} \kappa_{|W|} \right), \quad \forall n \geq 1.$$

(For instance $\kappa_1 = \varphi(a)(= 1)$, $\kappa_2 = \varphi(a^2) - (\varphi(a))^2$ and $\kappa_3 = \varphi(a^3) - 3\varphi(a)\varphi(a^2) + 2(\varphi(a))^3$. For a more detailed discussion of equation (4.23), see e.g. Lecture 12 of [7].) It was recently observed in [6] that the formula expressing the free cumulants of a in terms of the coefficients of $1/S_a(z)$ is merely a shifted version of equation (4.23):

$$(4.24) \quad \kappa_n = \sum_{\gamma \in NC(n-1)} \left(\prod_{V \in \gamma} t_{|V|} \right), \quad \forall n \geq 2.$$

Formula (4.24) arises when comparing the various functional equations satisfied by the series M_a , R_a and S_a ; see Lemma 6.2 of [6].

Now, for every $n \geq 2$ one has a natural identification between $NC(n-1)$ and $\{\alpha \in NC(n) \mid \alpha \ll 1_n\}$. Indeed, the latter set consists of the partitions α in $NC(n)$ such that 1 and n belong to the same block of α ; and any such partition is uniquely obtained from a $\gamma \in NC(n-1)$ by adjoining the number n to the block of γ which contains 1. By using this identification, the summation on the right-hand side of (4.24) can be turned into a summation over $\{\alpha \in NC(n) \mid \alpha \ll 1_n\}$. It is suggestive to write the ensuing equation in the form

$$(4.25) \quad \kappa_n = \sum_{\substack{\alpha \in NC(n), \\ \alpha \ll 1_n}} \left(\prod_{\substack{U \in \alpha, \\ 1_n\text{-special}}} t_{|U|-1} \right) \cdot \left(\prod_{\substack{V \in \alpha, \\ \text{not } 1_n\text{-special}}} t_{|V|} \right),$$

where the product over U has in fact only one factor (a partition $\alpha \ll 1_n$ has a unique 1_n -special block, the one which contains 1 and n). It is a straightforward exercise, left to the reader, to verify that (4.25) can be upgraded to the statement that for every $n \geq 1$ and every $\beta \in NC(n)$ one has

$$(4.26) \quad \prod_{W \in \beta} \kappa_{|W|} = \sum_{\substack{\alpha \in NC(n), \\ \alpha \ll \beta}} \left(\prod_{\substack{U \in \alpha, \\ \beta\text{-special}}} t_{|U|-1} \right) \cdot \left(\prod_{\substack{V \in \alpha, \\ \text{not } \beta\text{-special}}} t_{|V|} \right).$$

Finally, we sum over $\beta \in NC(n)$ on both sides of (4.26) and we invoke (4.23) on the left-hand side, and (4.19) follows. $\square$

Remark 4.4. The double sum over α and β from equation (4.19) can be treated as an iterated sum in two ways. One of them (sum first over α , then over β) has in fact just been invoked at the end of the preceding proof. But actually it is the other order of summation (with β first) which brings the double sum to a simpler form, because it allows one to take advantage of the Boolean features of the partial order $\ll$; specifically, one arrives at equation (4.27) stated in the next theorem. It is remarkable that (4.27) closely resembles the formula used to define the concept of “ c -free cumulants” in the theory of conditionally free convolution; compare e.g. to the third displayed equation on p. 366 of [3].

Theorem 4.5. Let $(\mathcal{A}, \varphi)$ be an algebraic non-commutative probability space, let $a \in \mathcal{A}$ be such that $\varphi(a) = 1$, and consider the reciprocal S -transform $1/S_a(z) = \sum_{n=0}^{\infty} t_n z^n$. Then for every $n \geq 1$ one has

$$(4.27) \quad \varphi(a^n) = \sum_{\alpha \in NC(n)} \left(\prod_{\substack{U \in \alpha, \\ \text{outer}}} t_{|U|-1} \right) \cdot \left(\prod_{\substack{V \in \alpha, \\ \text{inner}}} t_{|V|-1} + t_{|V|} \right).$$

Proof. Let α be a partition in $NC(n)$, and let $\mathcal{I}, \mathcal{O}$ denote the set of inner blocks and respectively the set of outer blocks of α . Proposition 2.5 implies that

$$(4.28) \quad \begin{aligned} & \sum_{\substack{\beta \in NC(n), \\ \beta \gg \alpha}} \left(\prod_{\substack{U \in \alpha, \\ \beta\text{-special}}} t_{|U|-1} \right) \cdot \left(\prod_{\substack{V \in \alpha, \\ \text{not } \beta\text{-special}}} t_{|V|} \right) \\ &= \sum_{\substack{\mathcal{V} \subseteq \alpha \text{ such} \\ \text{that } \mathcal{V} \supseteq \mathcal{O}}} \left(\prod_{U \in \mathcal{V}} t_{|U|-1} \right) \cdot \left(\prod_{V \in \alpha \setminus \mathcal{V}} t_{|V|} \right). \end{aligned}$$

By performing the substitution $\mathcal{V} \setminus \mathcal{O} =: \mathcal{U}$ in the sum on the right-hand side of (4.28), this can be continued with

$$= \left(\prod_{U \in \mathcal{O}} t_{|U|-1} \right) \cdot \left[\sum_{\mathcal{U} \subseteq \mathcal{I}} \left(\prod_{U \in \mathcal{U}} t_{|U|-1} \right) \cdot \left(\prod_{V \in \mathcal{I} \setminus \mathcal{U}} t_{|V|} \right) \right].$$

But it is clear that the latter sum over $\mathcal{U} \subseteq \mathcal{I}$ is precisely the expansion of the product $\prod_{V \in \mathcal{I}} (t_{|V|-1} + t_{|V|})$. So, altogether, what we have obtained is that

$$(4.29) \quad \begin{aligned} & \sum_{\substack{\beta \in NC(n), \\ \beta \gg \alpha}} \left(\prod_{\substack{U \in \alpha, \\ \beta\text{-special}}} t_{|U|-1} \right) \cdot \left(\prod_{\substack{V \in \alpha, \\ \text{not } \beta\text{-special}}} t_{|V|} \right) \\ &= \left(\prod_{U \in \mathcal{O}} t_{|U|-1} \right) \cdot \left(\prod_{V \in \mathcal{I}} t_{|V|-1} + t_{|V|} \right). \end{aligned}$$

Equation (4.29) holds for every $\alpha \in NC(n)$. Let us sum over α on both its sides. Then on the left-hand side we obtain exactly the double sum over α and β which is known from Proposition 4.3 to be equal to $\varphi(a^n)$, and the required formula (4.27) follows. $\square$

REFERENCES

- [1] S.T. Belinschi, A. Nica. η -series and a Boolean Bercovici-Pata bijection for bounded k -tuples, *Advances in Mathematics* 217 (2008), 1-41. MR2357321 (2009c:46088)
- [2] S.T. Belinschi, A. Nica. Free Brownian motion and evolution towards $\boxplus$ -infinite divisibility for k -tuples, *International Journal of Mathematics* 20 (2009), 309-338. MR2500073
- [3] M. Bozejko, M. Leinert, R. Speicher. Convolution and limit theorems for conditionally free random variables, *Pacific Journal of Mathematics* 175 (1996), 357-388. MR1432836 (98j:46069)
- [4] W.Y.C. Chen, S.Y.J. Wu, C.H. Yan. Linked partitions and linked cycles, *European Journal of Combinatorics* 29 (2008), 1408-1426. MR2423730 (2009e:05026)
- [5] K. Dykema. Multilinear function series and transforms in free probability theory, *Advances in Mathematics* 208 (2007), 351-407. MR2304321 (2008k:46193)
- [6] M. Mastnak, A. Nica. Hopf algebras and the logarithm of the S -transform, to appear in *Transactions of the American Mathematical Society*, arXiv:0807.4169.

- [7] A. Nica, R. Speicher. *Lectures on the combinatorics of free probability*, London Mathematical Society Lecture Note Series, 335, Cambridge University Press, 2006. MR2266879 (2008k:46198)
- [8] M. Popa. Non-crossing linked partitions and multiplication of free random variables, preprint, 2008, arXiv:0812.2064.
- [9] R. Simion. Noncrossing partitions, *Discrete Mathematics* 217 (2000), 367-409. MR1766277 (2001g:05011)
- [10] R. Speicher. Multiplicative functions on the lattice of non-crossing partitions and free convolution, *Mathematische Annalen* 298 (1994), 611-628. MR1268597 (95h:05012)
- [11] R.P. Stanley. *Enumerative combinatorics*, volume 2, Cambridge Studies in Advanced Mathematics, 62, Cambridge University Press, 1999. MR1676282 (2000k:05026)
- [12] D. Voiculescu. Multiplication of certain noncommuting random variables, *Journal of Operator Theory* 18 (1987), 223-235. MR915507 (89b:46076)
- [13] D.V. Voiculescu, K.J. Dykema, A. Nica. *Free random variables*, CRM Monograph Series, 1, American Mathematical Society, 1992. MR1217253 (94c:46133)

DEPARTMENT OF PURE MATHEMATICS, UNIVERSITY OF WATERLOO, WATERLOO, ONTARIO
N2L 3G1, CANADA

E-mail address: `anica@math.uwaterloo.ca`