THE PINCHING CONSTANT OF MINIMAL HYPERSURFACES IN THE UNIT SPHERES

QIN ZHANG

(Communicated by Richard A. Wentworth)

ABSTRACT. In this paper, we prove that if M^n ($n \leq 8$) is a closed minimal hypersurface in a unit sphere $S^{n+1}(1)$, then there exists a positive constant $\alpha(n)$ depending only on n such that if $n \leq S \leq n + \alpha(n)$, then M is isometric to a Clifford torus, where S is the squared norm of the second fundamental form of M .

1. INTRODUCTION

Let M^n be an n -dimensional closed minimal hypersurface in a unit sphere $S^{n+1}(1)$ of dimension $n + 1$. Denote by S the squared norm of the second fundamental form of M^n . Lawson [2], Simons [1] and Chern, do Carmo, Kobayashi [3] obtained independently the famous rigidity theorem, which says that if $S \leq n$, then $S \equiv 0$ or $S \equiv n$; i.e., M^n is the great sphere $S^n(1)$ or the Clifford torus. Further discussions in this direction have been carried out by many other authors [6, 8, 9, 10, 11, 12, 13]. In [4], Peng and Terng proved that if the scalar curvature of M is constant, then there exists a positive constant $\alpha(n)$ depending only on n such that if $n \leq S \leq n + \alpha(n)$, then $S \equiv n$. Later, Cheng and Yang [14] improved the pinching constant $\alpha(n)$ to $n/3$. More generally, Peng and Terng [5] proved that if M^n ($n \leq 5$) is a closed minimal hypersurface in S^{n+1} , then there exists a positive constant $\alpha(n)$ depending only on n such that if $n \leq S \leq n + \alpha(n)$, then $S \equiv n$. So they proposed the following problem.

Let M^n ($n \geq 6$) be a closed minimal hypersurface in S^{n+1} . Does there exist a positive constant $\alpha(n)$ depending only on n such that if $n \leq S \leq n + \alpha(n)$, then $S \equiv n$?

In [15], Cheng gives a positive answer under the additional condition that M has only two distinct principal curvatures. Later, Cheng and Ishikawa [6] improved the result of Peng and Terng [5] when $n \leq 5$.

In this paper, we solve the problem proposed by Peng and Terng [5] for $n \leq 8$.

Theorem 1.1. *Let M^n ($n \leq 8$) be a closed minimal hypersurface in $S^{n+1}(1)$. If $n \leq S \leq n + \alpha(n)$, then $S \equiv n$ and M^n is isometric to a Clifford torus $S^m(\sqrt{\frac{m}{n}}) \times S^{n-m}(\sqrt{\frac{n-m}{n}})$, where $\alpha(n) = \frac{2(n+4)(3-n\delta)}{9n+30}$, $\delta(3) = 0$, $\delta(4) = 0.16$, $\delta(5) = 0.23$, $\delta(6) = 0.28$, $\delta(7) = 0.32$ and $\delta(8) = 0.34$.*

Received by the editors June 7, 2009, and, in revised form, August 18, 2009.

2000 *Mathematics Subject Classification.* Primary 53C40.

Key words and phrases. Minimal hypersurface, Clifford torus, second fundamental form.

For $n \leq 5$, Cheng and Ishikawa [6] proved the following: Let M be an n -dimensional ($n \leq 5$) closed minimal hypersurface of a unit sphere $S^{n+1}(1)$. If $n \leq S \leq n + \alpha(n)$, then $S = n$, where $\alpha(3) = \frac{42}{85}$, $\alpha(4) = \frac{8}{31}$ and $\alpha(5) = \frac{3(21 - 5\sqrt{17})}{28 + 3\sqrt{17}}$. It is obvious that our pinching constant is larger than theirs. Up to now, the open problem for $n \geq 9$ is still open and it is a very hard problem.

2. FUNDAMENTAL FORMULAS

Let M^n be an n -dimensional hypersurface in an $(n+1)$ -dimensional unit sphere $S^{n+1}(1)$. We choose a local orthonormal frame field $e_1, \dots, e_{n+1}$ in $S^{n+1}(1)$, restricted to M^n , so that $e_1, \dots, e_n$ are tangent to M^n . Let $\omega_1, \dots, \omega_{n+1}$ denote the dual coframe field in $S^{n+1}(1)$. Then in M^n , $\omega_{n+1} = 0$. It follows from Cartan's Lemma that

$$(2.1) \quad \omega_{n+1i} = \sum_j h_{ij} \omega_j.$$

The second fundamental form α and the mean curvature H of M^n are defined by

$$(2.2) \quad \alpha = \sum_{i,j} h_{ij} \omega_i \omega_j e_{n+1}, \quad nH = \sum_i h_{ii},$$

respectively. If M^n is a minimal hypersurface, then $\sum_i h_{ii} = 0$. The connection form ω_{ij} is characterized by the structure equations

$$(2.3) \quad d\omega_i + \sum_j \omega_{ij} \wedge \omega_j = 0, \quad \omega_{ij} + \omega_{ji} = 0,$$

$$(2.4) \quad d\omega_{ij} + \sum_k \omega_{ik} \wedge \omega_{kj} = \Omega_{ij},$$

$$(2.5) \quad \Omega_{ij} = \frac{1}{2} \sum_{k,l} R_{ijkl} \omega_k \wedge \omega_l,$$

where Ω_{ij} (resp. R_{ijkl}) denotes the curvature form (resp. the components of the curvature tensor) of M^n . The Gauss equation is given by

$$(2.6) \quad R_{ijkl} = (\delta_{ik}\delta_{jl} - \delta_{il}\delta_{jk}) + (h_{ik}h_{jl} - h_{il}h_{jk}).$$

Denote by h_{ijk} , h_{ijkl} , h_{ijklm} the components of the first, second and third covariant derivatives of the second fundamental form, respectively. Then

$$(2.7) \quad h_{ijk} = h_{ikj} = h_{jik},$$

$$(2.8) \quad h_{ijkl} - h_{ijlk} = \sum_m h_{im} R_{mjkl} + \sum_m h_{mj} R_{mikl},$$

$$(2.9) \quad h_{ijklm} - h_{ijkml} = \sum_r h_{rjk} R_{ril m} + \sum_r h_{irk} R_{rjlm} + \sum_r h_{ijr} R_{rklm}.$$

For any fixed point p in M^n , we take a local orthonormal frame field $e_1, \dots, e_n$ such that

$$(2.10) \quad h_{ij} = \begin{cases} \lambda_i, & i = j, \\ 0, & i \neq j. \end{cases}$$

Let $S := \sum_{i,j} h_{ij}^2 = \sum_i \lambda_i^2$. The following formulas can be obtained by a direct computation (cf. [7]):

$$(2.11) \quad \frac{1}{2} \Delta S = \sum_{i,j,k} h_{ijk}^2 - S(S-n),$$

$$(2.12) \quad \frac{1}{2} \sum_{i,j,k} h_{ijk}^2 = \sum_{i,j,k,l} h_{ijkl}^2 + (2n+3-S) \sum_{i,j,k} h_{ijk}^2 + 3(2B-A) - \frac{3}{2} |\nabla S|^2,$$

where $A = \sum_{i,j,k} \lambda_i^2 h_{ijk}^2$, $B = \sum_{i,j,k} \lambda_i \lambda_j h_{ijk}^2$.

3. PROOF OF THEOREM 1.1

At first we give a proposition and some lemmas which will play a crucial role in the proof of our theorem. For convenience, we let

$$b_i = h_{ii1}, b = \sum_{i \neq 1} b_i^2 + \frac{1}{3} b_1^2, f = \sum_{i \neq 1} (\lambda_1^2 - 4\lambda_1 \lambda_i) b_i^2 - \lambda_1^2 b_1^2.$$

Proposition 3.1. *Let M^n be a closed minimal hypersurface in $S^{n+1}(1)$. Suppose that*

$$3(A-2B) \leq [2 + \delta(n)] \sum_{i,j,k} h_{ijk}^2,$$

where $\delta(n)$ is a number depending only on n such that $0 \leq \delta(n) < \min\{\frac{1}{2}, \frac{3}{n}\}$. Then there exists a positive constant $\alpha(n)$ depending only on n such that if $n \leq S \leq n + \alpha(n)$, then $S \equiv n$; i.e., M^n is isometric to a Clifford torus $S^m(\sqrt{\frac{m}{n}}) \times S^{n-m}(\sqrt{\frac{n-m}{n}})$. Here, $\alpha(n) = \frac{2(n+4)(3-n\delta)}{9n+30}$.

Proof. Since M^n is a minimal hypersurface in $S^{n+1}(1)$, from (2.11) and (2.12) we have

$$(3.1) \quad \int_M \sum_{i,j,k} h_{ijk}^2 dM = \int_M S(S-n) dM,$$

$$(3.2) \quad -\frac{1}{2} \int_M |\nabla S|^2 dM = \int_M \left[S \sum_{i,j,k} h_{ijk}^2 - S^2(S-n) \right] dM,$$

$$(3.3) \quad \int_M \sum_{i,j,k,l} h_{ijkl}^2 dM = \int_M \left[(S-2n-3) \sum_{i,j,k} h_{ijk}^2 + 3(A-2B) + \frac{3}{2} |\nabla S|^2 \right] dM.$$

Letting $f_3 = \sum_i \lambda_i^3$ and $f_4 = \sum_i \lambda_i^4$, we have (cf. [6])

$$(3.4) \quad \sum_{i,j,k,l} h_{ijkl}^2 \geq \frac{3}{2} (Sf_4 - f_3^2 - 2S^2 + nS) + \frac{3S(S-n)^2}{2(n+4)},$$

$$(3.5) \quad \int_M (A-2B) dM = \int_M \left[Sf_4 - f_3^2 - S^2 - \frac{1}{4} |\nabla S|^2 \right] dM.$$

From (3.3), (3.4) and (3.5), we have

$$(3.6) \quad \int_M \left[(S-2n-3) \sum_{i,j,k} h_{ijk}^2 + \frac{3}{2} (A-2B) + \frac{3}{2} S(S-n) + \frac{9}{8} |\nabla S|^2 - \frac{3S(S-n)^2}{2(n+4)} \right] dM \geq 0.$$

Noticing that $S^2 = S(S - n) + nS$, from (3.1), (3.2) and (3.6), we have

$$(3.7) \quad \int_M \left[\frac{3}{2}(A - 2B) + \frac{9n + 30}{4(n + 4)}S(S - n)^2 - \left(\frac{5}{4}S - \frac{n}{4} + \frac{3}{2} \right) \sum_{i,j,k} h_{ijk}^2 \right] dM \geq 0.$$

Suppose $3(A - 2B) \leq [2 + \delta(n)]S \sum_{i,j,k} h_{ijk}^2$ and $n \leq S \leq n + \alpha(n)$. From the above inequality, we have

$$(3.8) \quad \int_M \left\{ \frac{9n + 30}{4(n + 4)}\alpha(n) - \frac{1 - 2\delta(n)}{4}(S - n) - \frac{3 - n\delta(n)}{2} \right\} \sum_{i,j,k} h_{ijk}^2 dM \geq 0.$$

Since $\alpha(n) = \frac{2(n + 4)(3 - n\delta)}{9n + 30}$ and $\delta(n) < \min \left\{ \frac{1}{2}, \frac{3}{n} \right\}$, we have

$$(3.9) \quad - \int_M (S - n) \sum_{i,j,k} h_{ijk}^2 dM \geq 0.$$

Hence, $S \equiv n$. This completes the proof of Proposition 3.1. $\square$

Lemma 3.2. *Let M^n be a closed minimal hypersurface in $S^{n+1}(1)$. If $\lambda_1^2 - 4\lambda_1\lambda_2 \geq tS$ ($t \geq 2$), then $(\lambda_1^2 - 4\lambda_1\lambda_2) - (\lambda_1^2 - 4\lambda_1\lambda_i) \geq rS$ ($i \neq 1, 2$). Here, $r = \frac{16t - 8 - 12\sqrt{-2t^2 + 2t + 8}}{17}$.*

Proof. Let $\lambda_1^2 = x^2S$, $\lambda_2^2 = y^2S$ ($x, y > 0$). Since $\lambda_1^2 - 4\lambda_1\lambda_2 \geq tS$ ($t \geq 2$), we have $x^2 + 4xy \geq t$, that is, $y \geq \frac{t - x^2}{4x}$. Hence, we have

$$\begin{aligned} \lambda_1^2 - 4\lambda_1\lambda_i &\leq \left(x^2 + 4x\sqrt{1 - x^2 - y^2} \right) S \\ &\leq \left\{ x^2 + \sqrt{16x^2 - 16x^4 - (t - x^2)^2} \right\} S \\ &= \left\{ x^2 + \sqrt{-17x^4 + (16 + 2t)x^2 - t^2} \right\} S. \end{aligned}$$

Let $g(z) = z + \sqrt{-17z^2 + (16 + 2t)z - t^2}$ ($0 < z < 1$). Then

$$g'(z) = 1 - \frac{17z - (8 + t)}{\sqrt{-17z^2 + (16 + 2t)z - t^2}}.$$

Letting $g'(z_0) = 0$, we have

$$z_0 = \frac{3(8 + t) + 2\sqrt{-2t^2 + 2t + 8}}{51}.$$

Hence we have

$$g(z) \leq g(z_0) = \frac{t + 8 + 12\sqrt{-2t^2 + 2t + 8}}{17},$$

which implies that

$$\lambda_1^2 - 4\lambda_1\lambda_i \leq \frac{t + 8 + 12\sqrt{-2t^2 + 2t + 8}}{17}.$$

Since $\lambda_1^2 - 4\lambda_1\lambda_2 \geq tS$, we have

$$(\lambda_1^2 - 4\lambda_1\lambda_2) - (\lambda_1^2 - 4\lambda_1\lambda_i) \geq \frac{16t - 8 - 12\sqrt{-2t^2 + 2t + 8}}{17}.$$

This completes the proof of Lemma 3.2. $\square$

Lemma 3.3. *Let $f_n(t) = 17[t - 2 - \delta(n)][3(n - 2)t + (n + 2)\delta(n) + 10 - 4n]$ and $g_n(t) = [8 + 16\delta(n)](4t - 2 - 3\sqrt{-2t^2 + 2t + 8})$. Then*

$$h_n(t) = f_n(t) - g_n(t) \leq 0 \quad (t \geq 2, 4 \leq n \leq 8).$$

Here, $\delta(4) = 0.16$, $\delta(5) = 0.23$, $\delta(6) = 0.28$, $\delta(7) = 0.32$ and $\delta(8) = 0.34$.

Proof. By a direct computation, we obtain

$$h_n''(t) = 51 \left\{ 2(n - 2) - [8 + 16\delta(n)](-2t^2 + 2t + 8)^{-\frac{3}{2}} \right\}$$

and

$$h_n'''(t) = -153[8 + 16\delta(n)](2t - 1)(-2t^2 + 2t + 8)^{-\frac{5}{2}} < 0 \quad (t \geq 2).$$

On the other hand, $h_n''(2) > 0$ and $h_n(2) < 0$. Hence, if there exist real numbers $t_i > 2$ ($i = 1, 2$) such that $h_n'(t_1) > 0$, $h_n'(t_2) < 0$ and $h_n(t) \leq 0$ ($t_1 \leq t \leq t_2$), then $h_n(t) \leq 0$ ($\forall t \geq 2$).

In the case $n = 4$, since $\delta(4) = 0.16$, we have

$$\begin{aligned} f_4(t) &= 17(6t^2 - 18t + 10.8864), \\ g_4(t) &= 10.56(4t - 2 - 3\sqrt{-2t^2 + 2t + 8}), \\ h_4'(t) &= 204t - 348.24 - 31.68(2t - 1)(-2t^2 + 2t + 8)^{-\frac{1}{2}}. \end{aligned}$$

By a direct computation, we obtain

$$(3.10) \quad h_4'(2.48) > 0, \quad h_4'(2.5) < 0.$$

On the other hand, when $2.48 \leq t \leq 2.5$, we have

$$f_4(t) \leq f_4(2.5) \leq 57.6, \quad g_4(t) \geq g_4(2.48) \geq 57.8.$$

This implies that

$$(3.11) \quad h_4(t) \leq 0 \quad (2.48 \leq t \leq 2.5).$$

From (3.10) and (3.11), we know that Lemma 3.3 is true in the case $n = 4$.

In the case $n = 5$, since $\delta(5) = 0.23$, we have

$$\begin{aligned} f_5(t) &= 17(9t^2 - 28.46t + 18.7097), \\ g_5(t) &= 11.68(4t - 2 - 3\sqrt{-2t^2 + 2t + 8}), \\ h_5'(t) &= 306t - 530.54 - 35.04(2t - 1)(-2t^2 + 2t + 8)^{-\frac{1}{2}}. \end{aligned}$$

By a direct computation, we obtain

$$(3.12) \quad h_5'(2.51) > 0, \quad h_5'(2.52) < 0.$$

On the other hand, when $2.51 \leq t \leq 2.52$, we have

$$f_5(t) \leq f_5(2.52) \leq 70.5, \quad g_5(t) \geq g_5(2.51) \geq 71.$$

This implies that

$$(3.13) \quad h_5(t) \leq 0 \quad (2.51 \leq t \leq 2.52).$$

From (3.12) and (3.13), we know that Lemma 3.3 is true in the case $n = 5$.

In the case $n = 6$, since $\delta(6) = 0.28$, we have

$$\begin{aligned} f_6(t) &= 17(12t^2 - 39.12t + 26.8128), \\ g_6(t) &= 12.48(4t - 2 - 3\sqrt{-2t^2 + 2t + 8}), \\ h_6'(t) &= 408t - 714.96 - 37.44(2t - 1)(-2t^2 + 2t + 8)^{-\frac{1}{2}}. \end{aligned}$$

By a direct computation, we obtain

$$(3.14) \quad h'_6(2.53) > 0, \quad h'_6(2.535) < 0.$$

On the other hand, when $2.53 \leq t \leq 2.535$, we have

$$f_6(t) \leq f_6(2.535) \leq 81, \quad g_6(t) \geq g_6(2.51) \geq 82.$$

This implies that

$$(3.15) \quad h_6(t) \leq 0 \quad (2.53 \leq t \leq 2.535).$$

From (3.14) and (3.15), we know that Lemma 3.3 is true in the case $n = 6$.

In the case $n = 7$, since $\delta(7) = 0.32$, we have

$$\begin{aligned} f_7(t) &= 17(15t^2 - 49.92t + 35.0784), \\ g_7(t) &= 13.12(4t - 2 - 3\sqrt{-2t^2 + 2t + 8}), \\ h'_7(t) &= 510t - 901.12 - 39.36(2t - 1)(-2t^2 + 2t + 8)^{-\frac{1}{2}}. \end{aligned}$$

By a direct computation, we obtain

$$(3.16) \quad h'_7(2.54) > 0, \quad h'_7(2.544) < 0.$$

On the other hand, when $2.54 \leq t \leq 2.544$, we have

$$f_7(t) \leq f_7(2.544) \leq 88, \quad g_7(t) \geq g_7(2.54) \geq 90.$$

This implies that

$$(3.17) \quad h_7(t) \leq 0 \quad (2.54 \leq t \leq 2.544).$$

From (3.16) and (3.17), we know that Lemma 3.3 is true in the case $n = 7$.

In the case $n = 8$, since $\delta(8) = 0.34$, we have

$$\begin{aligned} f_8(t) &= 17(18t^2 - 60.72t + 43.524), \\ g_8(t) &= 13.44(4t - 2 - 3\sqrt{-2t^2 + 2t + 8}), \\ h'_8(t) &= 612t - 1086 - 40.32(2t - 1)(-2t^2 + 2t + 8)^{-\frac{1}{2}}. \end{aligned}$$

By a direct computation, we obtain

$$(3.18) \quad h'_8(2.5465) > 0, \quad h'_8(2.5468) < 0.$$

On the other hand, when $2.5465 \leq t \leq 2.5468$, we have

$$f_8(t) \leq f_8(2.5468) \leq 95.78, \quad g_8(t) \geq g_8(2.5465) \geq 95.8.$$

This implies that

$$(3.19) \quad h_8(t) \leq 0 \quad (2.5465 \leq t \leq 2.5468).$$

From (3.18) and (3.19), we know that Lemma 3.3 is true in the case $n = 8$. This completes the proof of Lemma 3.3. $\square$

Lemma 3.4. *Let M^n be a closed minimal hypersurface in $S^{n+1}(1)$. Then*

$$f \leq [2 + \delta(n)]Sb, \quad 3 \leq n \leq 8.$$

Here, $\delta(3) = 0$, $\delta(4) = 0.16$, $\delta(5) = 0.23$, $\delta(6) = 0.28$, $\delta(7) = 0.32$ and $\delta(8) = 0.34$.

Proof. In the case $n = 3$, if $b_1 = 0$, then $b_2^2 = b_3^2 = \frac{1}{2}b$. Hence

$$\begin{aligned} f &= \frac{1}{2}(\lambda_1^2 - 4\lambda_1\lambda_2 + \lambda_1^2 - 4\lambda_1\lambda_3)b \\ &= \left\{ \lambda_1^2 - 2 \cdot \frac{\lambda_1}{\sqrt{2}} \cdot \sqrt{2}\lambda_2 - 2 \cdot \frac{\lambda_1}{\sqrt{2}} \cdot \sqrt{2}\lambda_3 \right\} b \\ &\leq 2(\lambda_1^2 + \lambda_2^2 + \lambda_3^2)b \leq 2Sb. \end{aligned}$$

When $b_1 \neq 0$, we let $b_2 = (x - \frac{1}{2})b_1$ and $\lambda_2 = (y - \frac{1}{2})\lambda_1$. Then

$$\begin{aligned} f &= \frac{36x^2 + 48xy + 3}{48x^2y^2 + 36x^2 + 20y^2 + 15} \cdot 2Sb \\ &= \frac{36x^2 + 48xy + 3}{36x^2 + 48xy + 3 + 48(xy - 1/2)^2 + 20y^2} \cdot 2Sb \\ &\leq 2Sb. \end{aligned}$$

From the above discussion, we know that Lemma 3.4 is true in the case $n = 3$.

In the case $4 \leq n \leq 8$, if $\lambda_1^2 - 4\lambda_1\lambda_i \leq [2 + \delta(n)]S$ ($\forall i \neq 1$), then Lemma 3.4 is true. Otherwise, without loss of generality, we suppose that $\lambda_1^2 - 4\lambda_1\lambda_2 = tS$ ($t \geq 2$) and $b_1 = xb_2$. Then

$$\sum_{i \neq 1, 2} b_i^2 \geq \frac{(1+x)^2}{n-2} b_2^2, \quad \lambda_1^2 \geq (t-2)S.$$

From the above inequalities and Lemma 3.2, we have

$$\begin{aligned} f - [2 + \delta(n)]Sb &\leq [t - 2 - \delta(n)]Sb_2^2 + [t - r - 2 - \delta(n)]S \sum_{i \neq 1, 2} b_i^2 \\ &\quad - \left\{ t - 2 + \frac{2 + \delta(n)}{3} \right\} Sb_1^2 \\ &\leq [t - 2 - \delta(n)]Sb_2^2 + \frac{t - r - 2 - \delta(n)}{n-2} (1+x)^2 Sb_2^2 \\ &\quad - \left\{ t - 2 + \frac{2 + \delta(n)}{3} \right\} x^2 Sb_2^2. \end{aligned}$$

Here, $r = \frac{16t - 8 - 12\sqrt{-2t^2 + 2t + 8}}{17}$.

Let $F(n, t, x) = t - 2 - \delta(n) + \frac{t - r - 2 - \delta(n)}{n-2} (1+x)^2 - \left\{ t - 2 + \frac{2 + \delta(n)}{3} \right\} x^2$.

The above inequality becomes

$$(3.20) \quad f - [2 + \delta(n)]Sb \leq F(n, t, x)Sb_2^2.$$

Since

$$\frac{1}{2} \frac{\partial F(n, t, x)}{\partial x} = \frac{t - r - 2 - \delta(n)}{n-2} (1+x) - \left\{ t - 2 + \frac{2 + \delta(n)}{3} \right\} x,$$

we have

(3.21)

$$\begin{aligned} F(n, t, x) &\leq F(n, t, x_0) \\ &= \frac{t - 2 - \delta(n)}{G} \left\{ (n-2)t + \frac{(n+2)\delta(n) + 10 - 4n}{3} \right\} - \frac{2 + 4\delta(n)}{3G} r. \end{aligned}$$

Here, $-x_0 = \frac{3[r+2+\delta(n)-t]}{3r+3(n-3)t+14+(n+1)\delta(n)-4n}$, $\frac{\partial F(n,t,x_0)}{\partial x} = 0$, and $G = r+2+\delta(n)-t+(n-2)\left(t-2+\frac{2+\delta(n)}{3}\right)$.

Notice that $h_n(t) = 51F(n,t,x_0)G$, where $h_n(t)$ is defined as in Lemma 3.3. From (3.20), (3.21) and Lemma 3.3, we have

$$f \leq [2+\delta(n)]Sb.$$

This completes the proof of Lemma 3.4. $\square$

Now we are in a position to give the proof of Theorem 1.1.

Proof of Theorem 1.1. From Lemma 3.4, we have obtained

$$f = \sum_{i \neq 1} (\lambda_1^2 - 4\lambda_1\lambda_i)h_{ii1}^2 - \lambda_1^2 h_{111}^2 \leq (2+\delta)S\left(\sum_{i \neq 1} h_{ii1}^2 + \frac{1}{3}h_{111}^2\right).$$

In general,

$$f_j = \sum_{i \neq j} (\lambda_j^2 - 4\lambda_i\lambda_j)h_{iij}^2 - \lambda_j^2 h_{jjj}^2 \leq (2+\delta)S\left(\sum_{i \neq j} h_{iij}^2 + \frac{1}{3}h_{jjj}^2\right), \forall j.$$

Hence we get

$$\begin{aligned} 3(A-2B) &= \sum_{i \neq j \neq k \neq i} [2(\lambda_i^2 + \lambda_j^2 + \lambda_k^2) - (\lambda_i + \lambda_j + \lambda_k)^2] h_{ijk}^2 \\ &\quad - 3 \sum_i \lambda_i^2 h_{iii}^2 + 3 \sum_{i \neq j} (\lambda_j^2 - 4\lambda_i\lambda_j) h_{iij}^2 \\ &\leq 2S \sum_{i \neq j \neq k \neq i} h_{ijk}^2 + 3 \sum_{i \neq j} [(\lambda_j^2 - 4\lambda_i\lambda_j) h_{iij}^2 - \lambda_j^2 h_{jjj}^2] \\ &\leq (2+\delta)S \left\{ \sum_{i \neq j \neq k \neq i} h_{ijk}^2 + 3 \sum_{i \neq j} h_{iij}^2 + \sum_j h_{jjj}^2 \right\} \\ &= (2+\delta)S \sum_{i,j,k} h_{ijk}^2. \end{aligned}$$

Notice that $\delta(3) = 0$, $\delta(4) = 0.16$, $\delta(5) = 0.23$, $\delta(6) = 0.28$, $\delta(7) = 0.32$, $\delta(8) = 0.34$ and $\alpha(n) = \frac{2(n+4)(3-n\delta)}{9n+30}$. We conclude from Proposition 3.1 that $S \equiv n$. This completes the proof of Theorem 1.1. $\square$

REFERENCES

1. J. Simons, *Minimal varieties in Riemannian manifolds*, Ann. of Math. (2), **88** (1968), 62–105. MR0233295 (38:1617)
2. H. B. Lawson Jr., *Local rigidity theorems for minimal hypersurfaces*, Ann. of Math. (2), **89** (1969), 187–197. MR0238229 (38:6505)
3. S. S. Chern, M. do Carmo, S. Kobayashi, *Minimal submanifolds of a sphere with second fundamental form of constant length*, Functional analysis and related fields, Springer, New York, 1970, pp. 59–75. MR0273546 (42:8424)
4. C. K. Peng, C. L. Terng, *Minimal hypersurfaces of spheres with constant scalar curvature*, Ann. of Math. Stud., 103, Princeton University Press, 1983, 179–198. MR795235 (87k:53143)
5. C. K. Peng, C. L. Terng, *The scalar curvature of minimal hypersurfaces in spheres*, Math. Ann., **266** (1983), 105–113. MR722930 (85c:53099)
6. Q. M. Cheng, S. Ishikawa, *A characterization of the Clifford torus*, Proc. Amer. Math. Soc., **127** (1999), 819–828. MR1636934 (99g:53064)

7. Q. M. Cheng, *The classification of complete hypersurfaces with nonzero constant mean curvature of space form of dimension 4*, Mem. Fac. Sci. Kyushu Univ. Ser. A, **47** (1993), 79–102. MR1222356 (94h:53067)
8. S. Y. Cheng, S. T. Yau, *Hypersurfaces with constant scalar curvature*, Math. Ann., **225** (1977), no. 3, 195–204. MR0431043 (55:4045)
9. A. M. Li, J. M. Li, *An intrinsic rigidity theorem for minimal submanifolds in a sphere*, Arch. Math. (Basel), **58** (1992), no. 6, 582–594. MR1161925 (93b:53050)
10. H. W. Xu, *A rigidity theorem for submanifolds with parallel mean curvature in a sphere*, Arch. Math. (Basel), **61** (1993), no. 5, 489–496. MR1241055 (94m:53084)
11. H. W. Xu, *On closed minimal submanifolds in pinched Riemannian manifolds*, Trans. Amer. Math. Soc., **347** (1995), no. 5, 1743–1751. MR1243175 (95h:53088)
12. H. W. Xu, W. Fang, F. Xiang, *A generalization of Gauchman's rigidity theorem*, Pacific J. Math., **228** (2006), no. 1, 185–199. MR2263029 (2007j:53048)
13. S. T. Yau, *Submanifolds with constant mean curvature. I, II*, Amer. J. Math., **96** (1974), 346–366; *ibid.*, **97** (1975), 76–100. MR0370443 (51:6670)
14. Q. M. Cheng, H. C. Yang, *Chern's conjecture on minimal hypersurfaces*, Math. Z., **227** (1998), no. 3, 377–390. MR1612653 (99c:53070)
15. Q. M. Cheng, *The rigidity of Clifford torus $S^1(\sqrt{\frac{1}{n}}) \times S^{n-1}(\sqrt{\frac{n-1}{n}})$* , Comment. Math. Helvetici, **71** (1996), 60–69. MR1371678 (97a:53094)
16. Y. T. Zhang, S. L. Xu, *Rigidity of the Clifford torus $S^m(\sqrt{m/n}) \times S^{n-m}(\sqrt{(n-m)/n})$* , Acta Mathematica Scientia Ser. A Chin. Ed., **28** (2008), 128–132. MR2392041 (2008m:53145)

INSTITUTE OF MATHEMATICS AND SOFTWARE SCIENCE, SICHUAN NORMAL UNIVERSITY,
CHENGDU 610066, PEOPLE'S REPUBLIC OF CHINA
E-mail address: zhangdiligence@126.com