UNIFORM ASYMPTOTIC EXPANSIONS OF THE TRICOMI-CARLITZ POLYNOMIALS

K. F. LEE AND R. WONG

(Communicated by Walter Van Assche)

Dedicated to the Lord

ABSTRACT. The Tricomi-Carlitz polynomials satisfy the second-order linear difference equation

$$(n+1)f_{n+1}^{(\alpha)}(x) - (n+\alpha)x f_n^{(\alpha)}(x) + f_{n-1}^{(\alpha)}(x) = 0, \quad n \geq 1,$$

with initial values $f_0^{(\alpha)}(x) = 1$ and $f_1^{(\alpha)}(x) = \alpha x$, where x is a real variable and α is a positive parameter. An asymptotic expansion is derived for these polynomials by using the turning-point theory for three-term recurrence relations developed by Wang and Wong [Numer. Math. **91**(2002) and **94**(2003)]. The result holds uniformly in regions containing the critical values $x = \pm 2/\sqrt{\nu}$, where $\nu = n + 2\alpha - 1/2$.

1. INTRODUCTION

The Tricomi polynomials are defined by

$$(1.1) \quad t_n^{(\alpha)}(x) = \sum_{k=0}^n (-1)^k \binom{x-\alpha}{k} \frac{x^{n-k}}{(n-k)!}, \quad n = 0, 1, 2, \dots,$$

which satisfy the recurrence relation

$$(1.2) \quad (n+1)t_{n+1}^{(\alpha)}(x) - (n+\alpha)t_n^{(\alpha)}(x) + x t_{n-1}^{(\alpha)}(x) = 0, \quad n \geq 1,$$

with initial values $t_0^{(\alpha)}(x) = 1$, $t_1^{(\alpha)}(x) = \alpha$. Tricomi [7] observed that $\{t_n^{(\alpha)}(x)\}$ is not a system of orthogonal polynomials, since the recurrence relation (1.2) fails to have the required form [6, p.135]. However, Carlitz [1] discovered that if one sets

$$(1.3) \quad f_n^{(\alpha)}(x) = x^n t_n^{(\alpha)}(x^{-2}),$$

then $f_n^{(\alpha)}(x)$ satisfies

$$(1.4) \quad (n+1)f_{n+1}^{(\alpha)}(x) - (n+\alpha)x f_n^{(\alpha)}(x) + f_{n-1}^{(\alpha)}(x) = 0, \quad n \geq 1,$$

Received by the editors June 23, 2009, and, in revised form, November 6, 2009.

2010 *Mathematics Subject Classification*. Primary 41A60, 39A10; Secondary 33C45.

Key words and phrases. Tricomi-Carlitz polynomials, uniform asymptotic expansion, difference equation.

©2010 American Mathematical Society
 Reverts to public domain 28 years from publication

with initial values $f_0^{(\alpha)}(x) = 1$, $f_1^{(\alpha)}(x) = \alpha x$. Furthermore, he gave the orthogonal relation

$$(1.5) \quad \int_{-\infty}^{\infty} f_m^{(\alpha)}(x) f_n^{(\alpha)}(x) d\psi^{(\alpha)}(x) = \frac{2e^\alpha}{(n+\alpha)n!} \delta_{mn},$$

where $\psi^{(\alpha)}(x)$ is the step function whose jumps are

$$(1.6) \quad d\psi^{(\alpha)}(x) = \frac{(k+\alpha)^{k-1}e^{-k}}{k!} \quad \text{at } x = x_k = \pm(k+\alpha)^{-1/2}, \quad k = 0, 1, 2, \dots$$

The generating function

$$(1.7) \quad \exp\left\{\frac{w}{x} + \frac{1-\alpha x^2}{x^2} \log(1-wx)\right\} = \sum_{n=0}^{\infty} f_n^{(\alpha)}(x) w^n$$

can be derived from the recurrence relation (1.4) when $x \neq 0$. The series in (1.7) is convergent for $|wx| < 1$.

Asymptotic behavior of $f_n^{(\alpha)}(x)$ has been investigated by Goh and Wimp [2, 3]. In their first paper, they used an elementary approach to show that

$$(1.8) \quad \frac{f_n^{(\alpha)}(y/\sqrt{\alpha})}{\alpha^{-n/2} y^n n^{-\alpha(1-y^2)/y^2-1}} \rightarrow \frac{e^{\alpha/y^2}}{\Gamma(-\alpha(1-y^2)/y^2)}, \quad \text{as } n \rightarrow \infty,$$

uniformly for all y in a compact set $K \subseteq \mathbb{C} \setminus [-1, 1]$. (There is a minor error in the statement of their result; the factor $\alpha^{-n/2}$ in the denominator on the left-hand side of (1.8) is missing in their equation (23) in [2].) The validity of (1.8) can be verified by a direct application of Darboux's method [10, p. 116]. Goh and Wimp also observed that all zeros of $f_n^{(\alpha)}(y/\sqrt{\alpha})$ lie in the interval $[-1, 1]$. In their second paper, they used a saddle-point method to study the asymptotics of $f_n^{(\alpha)}(z/\sqrt{n})$ for z in neighborhoods of $z = \pm i/2$. Note that the scales in their two papers are different; in [2] the scale is $y/\sqrt{\alpha}$, whereas in [3] the scale is $z/\sqrt{n}$. The behavior of $f_n^{(\alpha)}(x)$ as $\alpha \rightarrow \infty$ has been studied by López and Temme [4]. Their result is expressed in terms of Hermite polynomials.

The purpose of this paper is to present an asymptotic expansion for $f_n^{(\alpha)}(t/\sqrt{\nu})$, which holds uniformly for t in $[0, \infty)$, where $\nu = n + 2\alpha - 1/2$. In view of the reflection formula $f_n^{(\alpha)}(-x) = (-1)^n f_n^{(\alpha)}(x)$, a corresponding result can be given for t in $(-\infty, 0]$. Note that our result is not covered in the two papers of Goh and Wimp; in fact, it complements those in [2] and [3]. We also point out that the Tricomi-Carlitz polynomials do not satisfy a second-order differential equation; hence the powerful tools developed for differential equations (see, e.g., [5]) are not applicable. Our approach is to use a turning-point theory recently introduced by Wang and Wong [8, 9] for three-term recurrence relations.

2. DIFFERENCE EQUATION

Returning to (1.4), we write

$$(2.1) \quad f_{n+1}^{(\alpha)}(x) - \frac{n+\alpha}{n+1} x f_n^{(\alpha)}(x) + \frac{1}{n+1} f_{n-1}^{(\alpha)}(x) = 0$$

and introduce the sequence $\{K_n\}$ defined by

$$(2.2) \quad (n+1)K_{n+1} = K_{n-1}$$

with $K_0 = 1$ and $K_1 = \sqrt{2/\pi}$. Induction shows that

$$(2.3) \quad K_n = \frac{1}{2^{n/2} \Gamma(\frac{1}{2}n + 1)}.$$

With the notation

$$(2.4) \quad F_n^{(\alpha)}(x) := \frac{f_n^{(\alpha)}(x)}{K_n},$$

equation (2.1) can be put in the canonical form considered in [9],

$$(2.5) \quad F_{n+1}^{(\alpha)}(x) - (A_n x + B_n) F_n^{(\alpha)}(x) + F_{n-1}^{(\alpha)}(x) = 0$$

with

$$(2.6) \quad A_n = \frac{n + \alpha}{n + 1} \frac{K_n}{K_{n+1}} = \frac{n + \alpha}{\sqrt{2}} \frac{\Gamma(\frac{1}{2}(n + 1))}{\Gamma(\frac{1}{2}n + 1)}$$

and $B_n = 0$. To find an asymptotic expansion for A_n , we recall the well-known result [10, p. 47]

$$\frac{\Gamma(\frac{1}{2}n + \frac{1}{2})}{\Gamma(\frac{1}{2}n + 1)} \sim \sqrt{\frac{2}{n}} \left[1 - \frac{1}{4n} + \frac{1}{32n^2} + \cdots \right].$$

Thus,

$$(2.7) \quad A_n \sim n^{1/2} \left[1 + \left(\alpha - \frac{1}{4} \right) \frac{1}{n} + \left(\frac{1}{32} - \frac{\alpha}{4} \right) \frac{1}{n^2} + \cdots \right].$$

In terms of the notation

$$(2.8) \quad A_n \sim n^{-\theta} \sum_{s=0}^{\infty} \frac{\alpha_s}{n^s} \quad \text{and} \quad B_n \sim \sum_{s=0}^{\infty} \frac{\beta_s}{n^s},$$

used in [9], we have

$$(2.9) \quad \theta = -\frac{1}{2}, \quad \alpha_0 = 1, \quad \alpha_1 = \alpha - \frac{1}{4}, \quad \alpha_2 = \left(\frac{1}{32} - \frac{\alpha}{4} \right), \dots$$

and $\beta_0 = \beta_1 = \beta_2 = \cdots = 0$. If these expansions are recast in the form

$$(2.10) \quad A_n \sim \nu^{-\theta} \sum_{s=0}^{\infty} \frac{\alpha'_s}{\nu^s} \quad \text{and} \quad B_n \sim \sum_{s=0}^{\infty} \frac{\beta'_s}{\nu^s},$$

where $\nu = n + \tau_0$ and τ_0 is some fixed real number to be determined, it is easily found that

$$(2.11) \quad \alpha'_0 = 1, \quad \alpha'_1 = \left(\alpha - \frac{1}{4} \right) - \frac{\tau_0}{2}, \dots$$

and $\beta'_0 = \beta'_1 = \beta'_2 = \cdots = 0$. To apply the result in [9], we first choose τ_0 so that $\alpha'_1 = 0$. From (2.11), it is obvious that the choice is

$$(2.12) \quad \tau_0 = 2\alpha - \frac{1}{2}.$$

According to equation (2.4) in [9], the characteristic equation is

$$(2.13) \quad \lambda^2 - t\lambda + 1 = 0,$$

where t is the rescaled variable $x = \nu^{-\frac{1}{2}}t$. The two roots of this equation are

$$(2.14) \quad \lambda_{\pm} = \frac{1}{2}(t \pm \sqrt{t^2 - 4}).$$

The points $t_{\pm} = \pm 2$ where the two roots coincide are called the *turning points* of equation (2.5). In view of the symmetry relation $F_n^{(\alpha)}(-x) = (-1)^n F_n^{(\alpha)}(x)$, we may restrict ourselves just to the case $0 < t < \infty$.

We now define the function $\zeta(t)$ introduced in [9, (4.10)]. With $t_+ = 2$, $\theta = -\frac{1}{2}$, $\alpha'_0 = 1$ and $\beta'_0 = 0$, this function is given by

$$\frac{2}{3}[\zeta(t)]^{3/2} := \log \frac{t + \sqrt{t^2 - 4}}{2} - \frac{1}{t^2} \int_2^t \frac{s^2}{\sqrt{s^2 - 4}} ds, \quad t \geq 2,$$

and

$$\frac{2}{3}[-\zeta(t)]^{3/2} := t^{-2} \int_t^2 \frac{s^2}{\sqrt{4 - s^2}} ds - \cos^{-1} \frac{t}{2}, \quad 0 < t < 2.$$

By direct calculation, one obtains

$$(2.15) \quad \frac{2}{3}[\zeta(t)]^{3/2} = \log \frac{t + \sqrt{t^2 - 4}}{2} - \frac{1}{t^2} \left[\frac{1}{2} t \sqrt{t^2 - 4} + 2 \log |t + \sqrt{t^2 - 4}| - 2 \ln 2 \right],$$

for $t \geq 2$, and

$$(2.16) \quad \frac{2}{3}[-\zeta(t)]^{3/2} = \frac{1}{t^2} \left[-2 \sin^{-1} \frac{t}{2} + \pi + \frac{t}{2} \sqrt{4 - t^2} \right] - \cos^{-1} \frac{t}{2},$$

for $0 < t < 2$. We also define the functions $H_0(\zeta)$ and $\Phi(\zeta)$ introduced in [9, (4.19) and (4.28)]. In the present situation, these functions are given by

$$(2.17) \quad H_0(\zeta) = -\sqrt{\frac{t^2 - 4}{4\zeta}} \quad \text{and} \quad \Phi(\zeta) = 0,$$

where ζ is the function defined in (2.15) and (2.16). Note that in our special case, $\alpha'_1 = \beta'_1 = 0$; hence, according to the definition of $\Phi(\zeta)$ given in (4.28) of [9], the second equation in (2.17) holds for $0 < t < \infty$, instead of $t \geq \delta$, $0 < \delta < 2$.

With this preliminary work done, we can now apply the main result in [9] to conclude that there are constants $C_1(x)$ and $C_2(x)$ such that the polynomials $F_n^{(\alpha)}(x)$ in (2.4) can be expressed as

$$(2.18) \quad F_n^{(\alpha)}(x) = C_1(x)P_n(x) + C_2(x)Q_n(x),$$

where, with $x = \nu^{-\frac{1}{2}}t$, $P_n(x)$ and $Q_n(x)$ have the asymptotic expansions

$$(2.19) \quad P_n(\nu^{-\frac{1}{2}}t) \sim \left(\frac{4\zeta}{t^2 - 4} \right)^{\frac{1}{4}} \left[\text{Ai} \left(\nu^{\frac{2}{3}}\zeta \right) \sum_{s=0}^{\infty} \frac{\tilde{A}_s(\zeta)}{\nu^{s-\frac{1}{6}}} + \text{Ai}' \left(\nu^{\frac{2}{3}}\zeta \right) \sum_{s=0}^{\infty} \frac{\tilde{B}_s(\zeta)}{\nu^{s+\frac{1}{6}}} \right]$$

and

$$(2.20) \quad Q_n(\nu^{-\frac{1}{2}}t) \sim \left(\frac{4\zeta}{t^2 - 4} \right)^{\frac{1}{4}} \left[\text{Bi} \left(\nu^{\frac{2}{3}}\zeta \right) \sum_{s=0}^{\infty} \frac{\tilde{A}_s(\zeta)}{\nu^{s-\frac{1}{6}}} + \text{Bi}' \left(\nu^{\frac{2}{3}}\zeta \right) \sum_{s=0}^{\infty} \frac{\tilde{B}_s(\zeta)}{\nu^{s+\frac{1}{6}}} \right].$$

In (2.19) and (2.20), $\text{Ai}(\cdot)$ and $\text{Bi}(\cdot)$ are the Airy functions, the leading coefficients are given by

$$(2.21) \quad \tilde{A}_0(\zeta) = 1 \quad \text{and} \quad \tilde{B}_0(\zeta) = 0,$$

and the expansions hold uniformly for $0 \leq t < \infty$.

3. DETERMINATION OF $C_1(x)$ AND $C_2(x)$

First we examine the behavior of $f_n^{(\alpha)}(x)$ as $n \rightarrow \infty$. By using the classical steepest decent method, one can show that to leading order, we have

$$(3.1) \quad f_n^{(\alpha)}\left(\frac{t}{\sqrt{\nu}}\right) \sim \frac{\nu^{-\frac{n}{2}} e^{\nu\left(\frac{\lambda_+}{t} + \left(\frac{2}{t^2} - 1\right) \log \lambda_+\right)}}{\sqrt{2\pi\nu}(t^2 - 4)^{\frac{1}{4}}} \cos\left(\pi\alpha - \frac{\nu\pi}{t^2}\right) \\ + \frac{\nu^{-\frac{n}{2}} e^{\nu\left(\frac{\lambda_-}{t} + \left(\frac{2}{t^2} - 1\right) \log \lambda_-\right)}}{\sqrt{2\pi\nu}(t^2 - 4)^{\frac{1}{4}}} 2 \sin\left(\pi\alpha - \frac{\nu\pi}{t^2}\right),$$

when $t > 2$;

$$(3.2) \quad f_n^{(\alpha)}\left(\frac{t}{\sqrt{\nu}}\right) \sim \frac{\sqrt{2}\nu^{-\frac{n}{2}} e^{\frac{\nu}{2}}}{\sqrt{\pi\nu}(4 - t^2)^{\frac{1}{4}}} \sin\left[\nu\left(\frac{1}{2t}\sqrt{4 - t^2} + \frac{\pi}{t^2} - \frac{2}{t^2}\sin^{-1}\frac{t}{2} - \cos^{-1}\frac{t}{2}\right) \right. \\ \left. + \pi\alpha - \frac{\nu\pi}{t^2} + \frac{\pi}{4}\right],$$

when $0 < t < 2$ and

$$(3.3) \quad f_n^{(\alpha)}\left(\frac{2}{\sqrt{\nu}}\right) \sim \nu^{-\frac{n}{2}} e^{\frac{\nu}{2}} 3^{\frac{1}{3}} \nu^{-\frac{1}{3}} \left(\frac{1}{3\Gamma(\frac{2}{3})} \cos\left(\pi\alpha - \frac{\nu\pi}{4}\right) + \frac{\sqrt{3}}{3\Gamma(\frac{2}{3})} \sin\left(\pi\alpha - \frac{\nu\pi}{4}\right) \right),$$

when $t = 2$. Next we recall the well-known asymptotic formulas

$$(3.4) \quad \text{Ai}(\eta) \sim \frac{\eta^{-\frac{1}{4}}}{2\sqrt{\pi}} \exp\left(-\frac{2}{3}\eta^{\frac{3}{2}}\right)$$

and

$$(3.5) \quad \text{Bi}(\eta) \sim \frac{\eta^{-\frac{1}{4}}}{\sqrt{\pi}} \exp\left(\frac{2}{3}\eta^{\frac{3}{2}}\right)$$

as $\eta \rightarrow +\infty$; see [5, p. 392]. Furthermore, since $\nu = n + 2\alpha - 1/2$, it is readily seen from (3.4) and (3.5) that

$$\frac{P_n(\nu^{-\frac{1}{2}}t)}{2^{\frac{n}{2}}\Gamma(\frac{n}{2} + 1)} \sim \frac{\nu^{-\frac{n}{2}}\nu^{-\frac{1}{2}}}{\sqrt{2}\pi(t^2 - 4)^{\frac{1}{4}}} e^{\nu\left(\frac{\lambda_+}{t} + \left(\frac{2}{t^2} - 1\right) \log \lambda_+\right)},$$

$$\frac{Q_n(\nu^{-\frac{1}{2}}t)}{2^{\frac{n}{2}}\Gamma(\frac{n}{2} + 1)} \sim \frac{\sqrt{2}\nu^{-\frac{n}{2}}\nu^{-\frac{1}{2}}}{\pi(t^2 - 4)^{\frac{1}{4}}} e^{\nu\left(\frac{\lambda_-}{t} + \left(\frac{2}{t^2} - 1\right) \log \lambda_-\right)}.$$

Comparing the two sides of (2.18), one concludes that

$$(3.6) \quad C_1(x) = \sqrt{\pi} \cos\left(\pi\alpha - \frac{\pi}{x^2}\right), \quad C_2(x) = \sqrt{\pi} \sin\left(\pi\alpha - \frac{\pi}{x^2}\right).$$

Note that in obtaining (3.6), any one of the three formulas (3.1), (3.2) or (3.3) could have been used. In summary, we have from (2.4), (2.18), (2.19) and (2.20)

$$(3.7) \quad f_n^{(\alpha)}\left(\frac{t}{\sqrt{\nu}}\right) = \frac{\sqrt{\pi} \cos\left(\pi\alpha - \frac{\nu\pi}{t^2}\right)}{2^{\frac{1}{2}n}\Gamma(\frac{1}{2}n + 1)} \left(\frac{4\zeta}{t^2 - 4}\right)^{\frac{1}{4}} \nu^{\frac{1}{6}} \left[\text{Ai}\left(\nu^{\frac{2}{3}}\zeta\right) + O\left(\frac{1}{\nu}\right) \right] \\ + \frac{\sqrt{\pi} \sin\left(\pi\alpha - \frac{\nu\pi}{t^2}\right)}{2^{\frac{1}{2}n}\Gamma(\frac{1}{2}n + 1)} \left(\frac{4\zeta}{t^2 - 4}\right)^{\frac{1}{4}} \nu^{\frac{1}{6}} \left[\text{Bi}\left(\nu^{\frac{2}{3}}\zeta\right) + O\left(\frac{1}{\nu}\right) \right],$$

where $x = \nu^{-\frac{1}{2}}t$ and $\nu = n + 2\alpha - 1/2$. This result holds uniformly for $0 \leq t < \infty$. As a check on the validity of (3.7), we note that

$$(3.8) \quad f_{2n}^{(\alpha)}(0) = \frac{(-1)^n}{2^n n!}, \quad f_{2n+1}^{(\alpha)}(0) = 0, \quad n = 0, 1, 2, \dots$$

From (3.7), we obtain

$$(3.9) \quad f_n^{(\alpha)}(0) \sim \frac{e^{n/2}}{\sqrt{\pi n^{(n+1)/2}}} \sin\left(-\frac{n\pi}{2} + \frac{\pi}{2}\right).$$

In view of Stirling's formula, (3.9) agrees with (3.8).

4. NUMERICAL RESULTS

The expansion (3.7) is particularly useful near $t = 2$, where the two characteristic roots in (2.14) coincide. As an illustration, we take $\alpha = 1.9$ and $n = 100$. Table 1 provides exact and approximate values of $2^{n/2}\Gamma(n/2 + 1)f_n^{(\alpha)}(t/\sqrt{\nu})$. The last column of the table shows the percentage error of the approximations.

TABLE 1. Numerical Results

t	Exact values	Approximate values	Error
1.0	$-7.46841313 \times 10^{-1}$	$-7.37169075 \times 10^{-1}$	0.0130
1.2	$6.30560302 \times 10^{-1}$	$6.34644034 \times 10^{-1}$	0.0065
1.4	$8.46491439 \times 10^{-1}$	$8.31094179 \times 10^{-1}$	0.0182
1.6	-1.30620957	-1.29091632	0.0117
1.8	1.51877275	1.49426911	0.0161
2.0	1.90931070	1.89864389	0.0056
2.2	3.26791053×10^2	3.16621347×10^2	0.0311
2.4	-9.21372780×10^4	-8.83780892×10^4	0.0408
2.6	3.94171949×10^9	3.74229636×10^9	0.0506
2.8	$1.54972056 \times 10^{13}$	$1.45539824 \times 10^{13}$	0.0609
3.0	$7.21884326 \times 10^{16}$	$6.70106781 \times 10^{16}$	0.0717

REFERENCES

1. L. Carlitz, On some polynomials of Tricomi, *Boll. Un. Mat. Ital.*, **13**, 58-64, 1958. MR0103303 (21:2078)
2. W.M.Y. Goh and J. Wimp, On the asymptotics of the Tricomi-Carlitz polynomials and their zero distribution. I, *SIAM J. Math. Anal.*, **25**, 420-428, 1994. MR1266567 (95b:42025)
3. W.M.Y. Goh and J. Wimp, The zero distribution of the Tricomi-Carlitz polynomials. Approximation theory and applications, *Comput. Math. Appl.*, **33**, 119-127, 1997. MR1442066 (99e:33006)
4. J.L. López and N.M. Temme, Approximation of orthogonal polynomials in terms of Hermite polynomials. Dedicated to Richard A. Askey on the occasion of his 65th birthday, Part II, *Methods Appl. Anal.*, **6**, 131-146, 1999. MR1803886 (2001m:33013)
5. F.W.J. Olver, *Asymptotics and Special Functions*, Academic Press, New York, 1974. (Reprinted by A. K. Peters, Ltd., Wellesley, 1997.) MR0435697 (55:8655)
6. G. Szegő, *Orthogonal Polynomials*, 4th edition, *Amer. Math. Soc. Colloq. Publ.*, **23**, Amer. Math. Soc., Providence, RI, 1975. MR0372517 (51:8724)
7. F.G. Tricomi, A class of non-orthogonal polynomials related to those of Laguerre, *J. Analyse Math.*, **13**, 209-231, 1951. MR0051351 (14:466e)
8. Z. Wang and R. Wong, Uniform asymptotic expansion of $J_\nu(\nu a)$ via a difference equation, *Numer. Math.*, **91**, 147-193, 2002. MR1896091 (2003g:33008)

9. Z. Wang and R. Wong, Asymptotic expansions for second-order linear difference equations with a turning point, *Numer. Math.*, **94**, 147–194, 2003. MR1971216 (2004c:39012)
10. R. Wong, *Asymptotic Approximations of Integrals*, Academic Press, Boston, 1989. MR1016818 (90j:41061)

DEPARTMENT OF MATHEMATICS, CITY UNIVERSITY OF HONG KONG, 83 TAT CHEE AVENUE,
KOWLOON, HONG KONG

E-mail address: charleslwf8571@gmail.com

DEPARTMENT OF MATHEMATICS, CITY UNIVERSITY OF HONG KONG, 83 TAT CHEE AVENUE,
KOWLOON, HONG KONG

E-mail address: rscwong@cityu.edu.hk