A NON-FINITELY GENERATED ALGEBRA OF FROBENIUS MAPS

MORDECHAI KATZMAN

(Communicated by Bernd Ulrich)

ABSTRACT. This paper describes an Artinian module over a ring of prime characteristic whose algebra of Frobenius maps is not finitely generated. This settles a question raised by Lyubeznik and Smith.

1. INTRODUCTION

The purpose of this paper is to answer a question raised by Gennady Lyubeznik and Karen Smith in [LS]. This question involves the finite generation of a certain non-commutative algebra which we define below (cf. section 3 in [LS]).

Let S be any commutative algebra of prime characteristic p . For any S -module M and all $e \geq 0$ we let $\mathcal{F}^e(M)$ denote the set of all additive functions $\phi : M \rightarrow M$ with the property that $\phi(sm) = s^{p^e} \phi(m)$ for all $s \in S$ and $m \in M$. Note that for all $e_1, e_2 \geq 0$ and $\phi_1 \in \mathcal{F}^{e_1}(M)$, $\phi_2 \in \mathcal{F}^{e_2}(M)$, the composition $\phi_2 \circ \phi_1$ is in $\mathcal{F}^{e_1+e_2}(M)$. Note also that each $\mathcal{F}^e(M)$ is a module over $\mathcal{F}^0(M) = \text{Hom}_S(M, M)$ via $\phi_0 \phi = \phi_0 \circ \phi$. We now define $\mathcal{F}(M) = \bigoplus_{e \geq 0} \mathcal{F}^e(M)$ and endow it with the structure of a $\text{Hom}_S(M, M)$ -algebra with multiplication given by composition.

In section 2 we construct an example of an Artinian module over a complete local ring S for which $\mathcal{F}(M)$ is not a finitely generated $\text{Hom}_S(M, M)$ -algebra, thus giving a negative answer to the question raised in section 3 of [LS].

2. THE EXAMPLE

Let $\mathbb{K}$ be a field of characteristic $p > 0$, $R = \mathbb{K}[[x, y, z]]$, and let $I \subseteq R$ be an ideal. Let E be the injective hull of the residue field of R and let f denote the standard Frobenius map of E (cf. section 4 in [K]). Write $S = R/I$ and let E_S be the injective hull of the residue field of S .

Notice that as S is complete, $\mathcal{F}^0(E_S) = \text{Hom}_S(E_S, E_S) \cong S$; the S -module $\mathcal{F}^e(E_S)$ of p^e th Frobenius maps on E_S is given by $(I^{[p^e]} : I)f^e$ (cf. section 4 in [K]).

For all $e \geq 1$ write $K_e = (I^{[p^e]} : I)$. We define

$$L_e = \sum_{\substack{1 \leq \beta_1, \dots, \beta_s < e \\ \beta_1 + \dots + \beta_s = e}} K_{\beta_1} K_{\beta_2}^{[p^{\beta_1}]} K_{\beta_3}^{[p^{\beta_1 + \beta_2}]} \dots K_{\beta_s}^{[p^{\beta_1 + \dots + \beta_{s-1}}]}.$$

Proposition 2.1. *Fix any $e \geq 1$, and let $\mathcal{F}_{<e}$ be the S -subalgebra of $\mathcal{F}(E_S)$ generated by $\mathcal{F}^0(E_S), \dots, \mathcal{F}^{e-1}(E_S)$. We have that $\mathcal{F}_{<e} \cap \mathcal{F}^e(E_S) = L_e f^e$.*

Received by the editors June 5, 2009 and, in revised form, December 16, 2009.
 2010 *Mathematics Subject Classification.* Primary 13A35, 13E10.

©2010 American Mathematical Society
 Reverts to public domain 28 years from publication

Proof. Any element in $\mathcal{F}_{<e} \cap \mathcal{F}^e(E_S)$ can be written as a sum of elements of the form $\phi_1 \cdots \phi_s$, where for all $1 \leq j \leq s$ we have $\phi_j \in \mathcal{F}^{\beta_j}(E_S)$ ($1 \leq \beta_j < e$) and $\beta_1 + \cdots + \beta_s = e$. Each such ϕ_j equals $a_j f^{\beta_j}$ where $a_j \in K_{\beta_j}$, so

$$\begin{aligned} \phi_1 \cdots \phi_s &= a_1 f^{\beta_1} a_2 f^{\beta_2} a_3 f^{\beta_3} \cdots a_s f^{\beta_s} \\ &= a_1 a_2^{\beta_1} a_3^{p^{\beta_1+\beta_2}} \cdots a_s^{p^{\beta_1+\cdots+\beta_{s-1}}} f^{\beta_1+\cdots+\beta_s} \in L_e f^e, \end{aligned}$$

hence $\mathcal{F}_{<e} \cap \mathcal{F}^e(E_S) \subseteq L_e f^e$.

On the other hand, for all $1 \leq \beta_1, \dots, \beta_s < e$ such that $\beta_1 + \cdots + \beta_s = e$,

$$K_{\beta_1} K_{\beta_2}^{[p^{\beta_1}]} K_{\beta_3}^{[p^{\beta_1+\beta_2}]} \cdots K_{\beta_s}^{[p^{\beta_1+\cdots+\beta_{s-1}}]} \subseteq (I^{[p^{\beta_1+\cdots+\beta_s}]} : I) = (I^{[p^e]} : I),$$

so $L_e f^e \subseteq (I^{[p^e]} : I) f^e = \mathcal{F}^e(E_S)$. A similar argument to the one in the previous paragraph shows that we also have

$$K_{\beta_1} K_{\beta_2}^{[p^{\beta_1}]} K_{\beta_3}^{[p^{\beta_1+\beta_2}]} \cdots K_{\beta_s}^{[p^{\beta_1+\cdots+\beta_{s-1}}]} f^e \subseteq \mathcal{F}_{<e},$$

and we deduce that $L_e f^e \subseteq \mathcal{F}_{<e} \cap \mathcal{F}^e(E_S)$. $\square$

Now fix I to be the ideal generated by xy and yz . We show that $\mathcal{F}(M)$ is not a finitely generated S -algebra.

Proposition 2.2. *For all $e \geq 1$, K_e is generated by*

$$\left\{ x^{p^e} y^{p^e-1}, x^{p^e-1} y^{p^e-1} z^{p^e-1}, y^{p^e-1} z^{p^e} \right\}.$$

Proof. For any $q > 1$,

$$\begin{aligned} (x^q y^q, y^q z^q) : (xy, yz) &= ((x^q y^q, y^q z^q) : xy) \cap ((x^q y^q, y^q z^q) : yz) \\ &= (x^{q-1} y^{q-1}, y^{q-1} z^q) \cap (x^q y^{q-1}, y^{q-1} z^{q-1}) \\ &= (x^q y^{q-1}, x^{q-1} y^{q-1} z^{q-1}, x^q y^{q-1} z^q, y^{q-1} z^q) \\ &= (x^q y^{q-1}, x^{q-1} y^{q-1} z^{q-1}, y^{q-1} z^q). \end{aligned}$$

$\square$

Theorem 2.3. *The S -algebra $\mathcal{F}(E_S)$ is not finitely generated.*

Proof. It is enough to show that for all $e \geq 1$, $\mathcal{F}(E_S)$ is not in $\mathcal{F}_{<e}$, and we establish this by showing that the generator $x^{p^e} y^{p^e-1}$ of K_e is not in L_e .

Since L_e is a sum of monomial ideals, $x^{p^e} y^{p^e-1} \in L_e$ if and only if $x^{p^e} y^{p^e-1}$ is in one of the summands. So we now fix $e \geq 1$ and $1 \leq \beta_1, \dots, \beta_s < e$ such that $\beta_1 + \cdots + \beta_s = e$, and we show that the ideal

$$K_{\beta_1} K_{\beta_2}^{[p^{\beta_1}]} K_{\beta_3}^{[p^{\beta_1+\beta_2}]} \cdots K_{\beta_s}^{[p^{\beta_1+\cdots+\beta_{s-1}}]}$$

does not contain $x^{p^e} y^{p^e-1}$.

Since z does not occur in $x^{p^e} y^{p^e-1}$, it is enough to show that with $J_e = x^{p^e} y^{p^e-1} R$,

$$J_{\beta_1} J_{\beta_2}^{[p^{\beta_1}]} J_{\beta_3}^{[p^{\beta_1+\beta_2}]} \cdots J_{\beta_s}^{[p^{\beta_1+\cdots+\beta_{s-1}}]}$$

does not contain $x^{p^e} y^{p^e-1}$. The exponent of x in the generator of the product above is

$$p^{\beta_1+(\beta_1+\beta_2)+\cdots+(\beta_1+\cdots+\beta_s)} > p^{\beta_1+\cdots+\beta_s} = p^e,$$

where the inequality follows from the fact that we must have $s > 1$. $\square$

3. A CONJECTURE

Although the example in section 2 settles the question raised in [LS], one might still raise the question of whether such examples exist over “nice” rings, e.g., normal domains.

Let $\mathbb{K}$ be a field of prime characteristic p , let $R = \mathbb{K}[[x, y, z, u, v, w]]$ and let I be the ideal generated by the 2×2 minors of the matrix $\begin{pmatrix} x & y & z \\ u & v & w \end{pmatrix}$. The ring $S = R/I$ is a normal, Cohen-Macaulay domain (cf. Theorem 7.3.1 in [BH]). Let E_S be the injective hull of the residue field of S and, as before, for all $e \geq 1$ let $\mathcal{F}_{<e}$ be the S -subalgebra of $\mathcal{F}^e(E_S)$ generated by $\mathcal{F}^1(E_S), \dots, \mathcal{F}^{e-1}(E_S)$. Note that $\mathcal{F}^0(E_S) = S$.

Conjecture 3.1. *For all $e \geq 1$, $\mathcal{F}^e(E_S)$ is not contained in $\mathcal{F}_{<e}$, and hence $\mathcal{F}^e(E_S)$ is not a finitely generated S -algebra.*

I have tested this conjecture using the computer system Macaulay 2 ([GS]), and, for example, in characteristic 2, it holds for $1 \leq e \leq 6$.

REFERENCES

- [BH] W. Bruns and J. Herzog. *Cohen-Macaulay rings*. Cambridge Studies in Advanced Mathematics, **39**. Cambridge University Press, Cambridge, 1993. MR1251956 (95h:13020)
- [GS] D. Grayson and M. Stillman. Macaulay 2 – a software system for algebraic geometry and commutative algebra, available at <http://www.math.uiuc.edu/Macaulay2>.
- [K] M. Katzman. *Parameter-test-ideals of Cohen-Macaulay rings*. Compositio Mathematica, **144** (2008), pp. 933–948. MR2441251 (2009d:13030)
- [LS] G. Lyubeznik and K. E. Smith. *On the commutation of the test ideal with localization and completion*. Transactions of the AMS **353** (2001), no. 8, pp. 3149–3180. MR1828602 (2002f:13010)

DEPARTMENT OF PURE MATHEMATICS, UNIVERSITY OF SHEFFIELD, HICKS BUILDING, SHEFFIELD S3 7RH, UNITED KINGDOM

E-mail address: M.Katzman@sheffield.ac.uk