POINT-COFINITE COVERS IN THE LAVER MODEL

ARNOLD W. MILLER AND BOAZ TSABAN

(Communicated by Julia Knight)

ABSTRACT. Let $S_1(\Gamma, \Gamma)$ be the statement: For each sequence of point-cofinite open covers, one can pick one element from each cover and obtain a point-cofinite cover. $\mathfrak{b}$ is the minimal cardinality of a set of reals not satisfying $S_1(\Gamma, \Gamma)$. We prove the following assertions:

- (1) If there is an unbounded tower, then there are sets of reals of cardinality $\mathfrak{b}$ satisfying $S_1(\Gamma, \Gamma)$.
- (2) It is consistent that all sets of reals satisfying $S_1(\Gamma, \Gamma)$ have cardinality smaller than $\mathfrak{b}$.

These results can also be formulated as dealing with Arhangel'skii's property α_2 for spaces of continuous real-valued functions.

The main technical result is that in Laver's model, each set of reals of cardinality $\mathfrak{b}$ has an unbounded Borel image in the Baire space ω^ω .

1. BACKGROUND

Let P be a nontrivial property of sets of reals. The *critical cardinality* of P , denoted $\text{non}(P)$, is the minimal cardinality of a set of reals not satisfying P . A natural question is whether there is a set of reals of cardinality at least $\text{non}(P)$, which satisfies P , i.e., a *nontrivial* example.

We consider the following property. Let X be a set of reals. $\mathcal{U}$ is a *point-cofinite* cover of X if $\mathcal{U}$ is infinite, and for each $x \in X$, $\{U \in \mathcal{U} : x \in U\}$ is a cofinite subset of $\mathcal{U}$.¹ Having X fixed in the background, let Γ be the family of all point-cofinite *open* covers of X . The following properties were introduced by Hurewicz [8], Tsaban [19], and Scheepers [15], respectively.

$\mathcal{U}_{\text{fin}}(\Gamma, \Gamma)$: For all $\mathcal{U}_0, \mathcal{U}_1, \dots \in \Gamma$, none containing a finite subcover, there are finite $\mathcal{F}_0 \subseteq \mathcal{U}_0, \mathcal{F}_1 \subseteq \mathcal{U}_1, \dots$ such that $\{\bigcup \mathcal{F}_n : n \in \omega\} \in \Gamma$.

$\mathcal{U}_2(\Gamma, \Gamma)$: For all $\mathcal{U}_0, \mathcal{U}_1, \dots \in \Gamma$, there are $\mathcal{F}_0 \subseteq \mathcal{U}_0, \mathcal{F}_1 \subseteq \mathcal{U}_1, \dots$ such that $|\mathcal{F}_n| = 2$ for all n , and $\{\bigcup \mathcal{F}_n : n \in \omega\} \in \Gamma$.

$S_1(\Gamma, \Gamma)$: For all $\mathcal{U}_0, \mathcal{U}_1, \dots \in \Gamma$, there are $U_0 \in \mathcal{U}_0, U_1 \in \mathcal{U}_1, \dots$ such that $\{U_n : n \in \omega\} \in \Gamma$.

Received by the editors October 21, 2009.

2010 *Mathematics Subject Classification*. Primary 03E35, 26A03; Secondary 03E17.

¹Historically, point-cofinite covers were named γ -covers, since they are related to a property numbered γ in a list from α to ϵ in the seminal paper [7] of Gerlits and Nagy.

Clearly, $S_1(\Gamma, \Gamma)$ implies $U_2(\Gamma, \Gamma)$, which in turn implies $U_{\text{fin}}(\Gamma, \Gamma)$. None of these implications is reversible in ZFC [19]. The critical cardinality of all three properties is $\mathfrak{b}$ [9].²

Bartoszyński and Shelah [1] proved that there are, provably in ZFC, totally imperfect sets of reals of cardinality $\mathfrak{b}$ satisfying the Hurewicz property $U_{\text{fin}}(\Gamma, \Gamma)$. Tsaban proved the same assertion for $U_2(\Gamma, \Gamma)$ [19]. These sets satisfy $U_{\text{fin}}(\Gamma, \Gamma)$ in all finite powers [2].

We show that in order to obtain similar results for $S_1(\Gamma, \Gamma)$, hypotheses beyond ZFC are necessary.

2. CONSTRUCTIONS

We show that certain weak (but not provable in ZFC) hypotheses suffice to have nontrivial $S_1(\Gamma, \Gamma)$ sets, even ones which possess this property in all finite powers.

Definition 2.1. A *tower* of cardinality κ is a set $T \subseteq [\omega]^\omega$ which can be enumerated bijectively as $\{x_\alpha : \alpha < \kappa\}$, such that for all $\alpha < \beta < \kappa$, $x_\beta \subseteq^* x_\alpha$.

A set $T \subseteq [\omega]^\omega$ is *unbounded* if the set of its enumeration functions is unbounded; i.e., for any $g \in \omega^\omega$ there is an $x \in T$ such that for infinitely many n , $g(n)$ is less than the n -th element of x .

Scheepers [16] proved that if $\mathfrak{t} = \mathfrak{b}$, then there is a set of reals of cardinality $\mathfrak{b}$ satisfying $S_1(\Gamma, \Gamma)$. If $\mathfrak{t} = \mathfrak{b}$, then there is an unbounded tower of cardinality $\mathfrak{b}$, but the latter assumption is weaker.

Lemma 2.2 (folklore). *If $\mathfrak{b} < \mathfrak{d}$, then there is an unbounded tower of cardinality $\mathfrak{b}$.*

Proof. Let $B = \{b_\alpha : \alpha < \mathfrak{b}\} \subseteq \omega^\omega$ be a $\mathfrak{b}$ -scale; that is, each b_α is increasing, $b_\alpha \leq^* b_\beta$ for all $\alpha < \beta < \mathfrak{b}$, and B is unbounded.

As $|B| < \mathfrak{d}$, B is not dominating. Let $g \in \omega^\omega$ exemplify that. For each $\alpha < \mathfrak{b}$, let $x_\alpha = \{n : b_\alpha(n) \leq g(n)\}$. Then $T = \{x_\alpha : \alpha < \mathfrak{b}\}$ is an unbounded tower. Clearly, $x_\beta \subseteq^* x_\alpha$ for $\alpha < \beta$. Assume that T is bounded, and let $f \in \omega^\omega$ exemplify that. For each α , writing $x_\alpha(n)$ for the n -th element of x_α :

$$b_\alpha(n) \leq b_\alpha(x_\alpha(n)) \leq g(x_\alpha(n)) \leq g(f(n))$$

for all but finitely many n . Thus, $g \circ f$ shows that B is bounded, a contradiction. $\square$

Theorem 2.3. *If there is an unbounded tower (of any cardinality), then there is a set of reals X of cardinality $\mathfrak{b}$ that satisfies $S_1(\Gamma, \Gamma)$.*

Theorem 2.3 follows from Propositions 2.4 and 2.5.

Proposition 2.4. *If there is an unbounded tower, then there is one of cardinality $\mathfrak{b}$.*

Proof. By Lemma 2.2, it remains to consider the case $\mathfrak{b} = \mathfrak{d}$. Let T be an unbounded tower of cardinality κ . Let $\{f_\alpha : \alpha < \mathfrak{b}\} \subseteq \omega^\omega$ be dominating. For each $\alpha < \mathfrak{b}$, pick $x_\alpha \in T$ which is not bounded by f_α . $\{x_\alpha : \alpha < \mathfrak{b}\}$ is unbounded, being unbounded in a dominating family. $\square$

²Blass's survey [4] is a good reference for the definitions and details of the special cardinals mentioned in this paper.

Define a topology on $P(\omega)$ by identifying $P(\omega)$ with the Cantor space 2^ω , via characteristic functions. Scheepers's mentioned proof actually establishes the following result, to which we give an alternative proof.

Proposition 2.5 (essentially, Scheepers [16]). *For each unbounded tower T of cardinality $\mathfrak{b}$, $T \cup [\omega]^{<\omega}$ satisfies $\mathsf{S}_1(\Gamma, \Gamma)$.*

Proof. Let $T = \{x_\alpha : \alpha < \mathfrak{b}\}$ be an unbounded tower of cardinality $\mathfrak{b}$. For each α , let $X_\alpha = \{x_\beta : \beta < \alpha\} \cup [\omega]^{<\omega}$. Let $\mathcal{U}_0, \mathcal{U}_1, \dots$ be point-cofinite open covers of $X_\mathfrak{b} = T \cup [\omega]^{<\omega}$. We may assume that each $\mathcal{U}_n$ is countable and that $\mathcal{U}_i \cap \mathcal{U}_j = \emptyset$ whenever $i \neq j$.

By the proof of Lemma 1.2 of [6], for each k there are distinct $U_0^k, U_1^k, \dots \in \mathcal{U}_k$, and an increasing sequence $m_0^k < m_1^k < \dots$, such that for each n and k ,

$$\{x \subseteq \omega : x \cap (m_n^k, m_{n+1}^k) = \emptyset\} \subseteq U_n^k.$$

As T is unbounded, there is $\alpha < \mathfrak{b}$ such that for each k , $I_k = \{n : x_\alpha \cap (m_n^k, m_{n+1}^k) = \emptyset\}$ is infinite.

For each k , $\{U_n^k : n \in \omega\}$ is an infinite subset of $\mathcal{U}_k$, and thus a point-cofinite cover of X_α . As $|X_\alpha| < \mathfrak{b}$, there is $f \in {}^\omega\omega$ such that

$$\forall x \in X_\alpha \exists k_0 \forall k \geq k_0 \forall n > f(k) \ x \in U_n^k.$$

For each k , pick $n_k \in I_k$ such that $n_k > f(k)$.

We claim that $\{U_{n_k}^k : k \in \omega\}$ is a point-cofinite cover of $X_\mathfrak{b}$: If $x \in X_\alpha$, then $x \in U_{n_k}^k$ for all but finitely many k , because $n_k > f(k)$ for all k . If $x = x_\beta$, $\beta \geq \alpha$, then $x \subseteq^* x_\alpha$. For each large enough k , $m_{n_k}^k$ is large enough, so that $x \cap (m_{n_k}^k, m_{n_k+1}^k) \subseteq x_\alpha \cap (m_{n_k}^k, m_{n_k+1}^k) = \emptyset$, and thus $x \in U_{n_k}^k$. $\square$

Remark 2.6. Zdomsky points out that for the proof to go through, it suffices that $\{x_\alpha : \alpha < \mathfrak{b}\}$ is such that there is an unbounded $\{y_\alpha : \alpha < \mathfrak{b}\} \subseteq [\omega]^\omega$ such that for each α , x_α is a pseudointersection of $\{y_\beta : \beta < \alpha\}$. We do not know whether the assertion mentioned here is weaker than the existence of an unbounded tower.

We now turn to nontrivial examples of sets satisfying $\mathsf{S}_1(\Gamma, \Gamma)$ in all finite powers. In general, $\mathsf{S}_1(\Gamma, \Gamma)$ is not preserved by taking finite powers [9], and we use a slightly stronger hypothesis in our construction.

Definition 2.7. Let $\mathfrak{b}_0$ be the additivity number of $\mathsf{S}_1(\Gamma, \Gamma)$, that is, the minimum cardinality of a family $\mathcal{F}$ of sets of reals, each satisfying $\mathsf{S}_1(\Gamma, \Gamma)$, such that the union of all members of $\mathcal{F}$ does not satisfy $\mathsf{S}_1(\Gamma, \Gamma)$.

$\mathfrak{t} \leq \mathfrak{h}$, and Scheepers proved that $\mathfrak{h} \leq \mathfrak{b}_0 \leq \mathfrak{b}$ [17]. It follows from Theorem 3.6 that, consistently, $\mathfrak{h} < \mathfrak{b}_0 = \mathfrak{b}$. It is open whether $\mathfrak{b}_0 = \mathfrak{b}$ is provable. If $\mathfrak{t} = \mathfrak{b}$ or $\mathfrak{h} = \mathfrak{b} < \mathfrak{d}$, then there is an unbounded tower of cardinality $\mathfrak{b}_0$.

Theorem 2.8. *For each unbounded tower T of cardinality $\mathfrak{b}_0$, all finite powers of $T \cup [\omega]^{<\omega}$ satisfy $\mathsf{S}_1(\Gamma, \Gamma)$.*

Proof. We say that $\mathcal{U}$ is an ω -cover of X if no member of $\mathcal{U}$ contains X as a subset, but each finite subset of X is contained in some member of $\mathcal{U}$. We need a multidimensional version of Lemma 1.2 of [6].

Lemma 2.9. *Assume that $[\omega]^{<\omega} \subseteq X \subseteq P(\omega)$, and let $e \in \omega$. For each open ω -cover $\mathcal{U}$ of X^e , there are $m_0 < m_1 < \dots$ and $U_0, U_1, \dots \in \mathcal{U}$, such that for all $x_0, \dots, x_{e-1} \subseteq \omega$, $(x_0, \dots, x_{e-1}) \in U_n$ whenever $x_i \cap (m_n, m_{n+1}) = \emptyset$ for all $i < e$.*

Proof. As $\mathcal{U}$ is an open ω -cover of X^e , there is an open ω -cover $\mathcal{V}$ of X such that $\{V^e : V \in \mathcal{V}\}$ refines $\mathcal{U}$ [9].

Let $m_0 = 0$. For each $n \geq 0$: Assume that $V_0, \dots, V_{n-1} \in \mathcal{V}$ are given, and $U_0, \dots, U_{n-1} \in \mathcal{U}$ are such that $V_i^e \subseteq U_i$ for all $i < n$. Fix a finite $F \subseteq X$ such that F^e is not contained in any of the sets $U_0, \dots, U_{n-1}$. As $\mathcal{V}$ is an ω -cover of X , there is $V_n \in \mathcal{V}$ such that $F \cup P(\{0, \dots, m_n\}) \subseteq V_n$. Take $U_n \in \mathcal{U}$ such that $V_n^e \subseteq U_n$. Then $U_n \notin \{U_0, \dots, U_{n-1}\}$. As V_n is open, for each $s \subseteq \{0, \dots, m_n\}$ there is k_s such that for each $x \in P(\omega)$ with $x \cap \{0, \dots, k_s - 1\} = s$, $x \in V_n$. Let $m_{n+1} = \max\{k_s : s \subseteq \{0, \dots, m_n\}\}$.

If $x_i \cap (m_n, m_{n+1}) = \emptyset$ for all $i < e$, then $(x_0, \dots, x_{e-1}) \in V_n^e \subseteq U_n$. $\square$

The assumption in the theorem that there is an unbounded tower of cardinality $\mathfrak{b}_0$ implies that $\mathfrak{b}_0 = \mathfrak{b}$. The proof is by induction on the power e of $T \cup [\omega]^{<\omega}$. The case $e = 1$ follows from Theorem 2.5.

Let $\mathcal{U}_0, \mathcal{U}_1, \dots \in \Gamma((T \cup [\omega]^{<\omega})^e)$. We may assume that these covers are countable. As in the proof of Theorem 2.5 (this time using Lemma 2.9), there are for each k , $m_0^k < m_1^k < \dots$ and $U_0^k, U_1^k, \dots \in \mathcal{U}_k$ (so that $\{U_n^k : n \in \omega\} \in \Gamma((T \cup [\omega]^{<\omega})^e)$), such that for all $y_0, \dots, y_{e-1} \subseteq \omega$, $(y_0, \dots, y_{e-1}) \in U_n^k$ whenever $y_i \cap (m_n^k, m_{n+1}^k) = \emptyset$ for all $i < e$.

Let α_0 be such that $X_{\alpha_0}^e$ is not contained in any member of $\bigcup_n \mathcal{U}_n$. As T is unbounded, there is α such that $\alpha_0 \leq \alpha < \mathfrak{b}$, and for each k , $I_k = \{n : x_\alpha \cap (m_n^k, m_{n+1}^k) = \emptyset\}$ is infinite.

Let $Y = \{x_\beta : \beta \geq \alpha\}$. $(T \cup [\omega]^{<\omega})^e \setminus Y^e$ is a union of fewer than $\mathfrak{b}_0$ homeomorphic copies of $(T \cup [\omega]^{<\omega})^{e-1}$. By the induction hypothesis, $(T \cup [\omega]^{<\omega})^{e-1}$ satisfies $S_1(\Gamma, \Gamma)$, and therefore so does $(T \cup [\omega]^{<\omega})^e \setminus Y^e$. For each k , $\{U_n^k : n \in I_k\}$ is a point-cofinite cover of $(T \cup [\omega]^{<\omega})^e \setminus Y^e$, and thus there are infinite $J_0 \subseteq I_0, J_1 \subseteq I_1, \dots$, such that $\{\bigcap_{n \in J_k} U_n^k : k \in \omega\}$ is a point-cofinite cover of $(T \cup [\omega]^{<\omega})^e \setminus Y^e$.³ For each k , pick $n_k \in J_k$ such that: $m_{n_k}^k > m_{n_{k-1}+1}^{k-1}$, $x_\alpha \cap (m_{n_k}^k, m_{n_k+1}^k) = \emptyset$, and $U_{n_k}^k \notin \{U_{n_0}^0, \dots, U_{n_{k-1}}^{k-1}\}$.

$\{U_{n_k}^k : k \in \omega\} \in \Gamma(T \cup [\omega]^{<\omega})$: If $x \in (T \cup [\omega]^{<\omega})^e \setminus Y^e$, then $x \in U_{n_k}^k$ for all but finitely many k . If $x = (x_{\beta_0}, \dots, x_{\beta_{e-1}}) \in Y$, then $\beta_0, \dots, \beta_{e-1} \geq \alpha$, and thus $x_{\beta_0}, \dots, x_{\beta_{e-1}} \subseteq^* x_\alpha$. For each large enough k , $m_{n_k}^k$ is large enough, so that $x_{\beta_i} \cap (m_{n_k}^k, m_{n_k+1}^k) \subseteq x_\alpha \cap (m_{n_k}^k, m_{n_k+1}^k) = \emptyset$ for all $i < e$, and thus $x \in U_{n_k}^k$. $\square$

There is an additional way to obtain nontrivial $S_1(\Gamma, \Gamma)$ sets: The hypothesis $\mathfrak{b} = \text{cov}(\mathcal{N}) = \text{cof}(\mathcal{N})$ provides $\mathfrak{b}$ -Sierpiński sets, and $\mathfrak{b}$ -Sierpiński sets satisfy $S_1(\Gamma, \Gamma)$, even for Borel point-cofinite covers. Details are available in [18].

We record the following consequence of Theorem 2.3 for later use.

Corollary 2.10. *For each unbounded tower T of cardinality $\mathfrak{b}$, $T \cup [\omega]^{<\omega}$ satisfies $S_1(\Gamma, \Gamma)$ for open covers, but not for Borel covers.*

Proof. The latter property is hereditary for subsets [18]. By a theorem of Hurewicz, a set of reals satisfies $U_{\text{fin}}(\Gamma, \Gamma)$ if and only if each continuous image of X in ω^ω is bounded. It follows that the set $T \subseteq T \cup [\omega]^{<\omega}$ does not even satisfy $U_{\text{fin}}(\Gamma, \Gamma)$. $\square$

³Choosing infinitely many elements from each cover, instead of one, can be done by adding to the given sequence of covers all cofinite subsets of the given covers.

3. A CONSISTENCY RESULT

By the results of the previous section, we have the following.

Lemma 3.1. *Assume that every set of reals with property $S_1(\Gamma, \Gamma)$ has cardinality $< \mathfrak{b}$, and $\mathfrak{c} = \aleph_2$. Then $\aleph_1 = \mathfrak{t} = \text{cov}(\mathcal{N}) < \mathfrak{b} = \aleph_2$.*

Proof. As there is no unbounded tower, we have that $\mathfrak{t} < \mathfrak{b} = \mathfrak{d}$. As $\mathfrak{c} = \aleph_2$, $\aleph_1 = \mathfrak{t} < \mathfrak{b} = \aleph_2$. Since there are no $\mathfrak{b}$ -Sierpiński sets and $\mathfrak{b} = \text{cof}(\mathcal{N}) = \mathfrak{c}$, $\text{cov}(\mathcal{N}) < \mathfrak{b}$. $\square$

In Laver's model [11], $\aleph_1 = \mathfrak{t} = \text{cov}(\mathcal{N}) < \mathfrak{b} = \aleph_2$. We will show that, indeed, $S_1(\Gamma, \Gamma)$ is trivial there. Laver's model was constructed to realize Borel's Conjecture, asserting that “strong measure zero” is trivial. In some sense, $S_1(\Gamma, \Gamma)$ is a dual of strong measure zero. For example, the canonical examples of $S_1(\Gamma, \Gamma)$ sets are Sierpiński sets, a measure-theoretic object, whereas the canonical examples of strong measure zero sets are Luzin sets, a Baire category theoretic object. More about that can be seen in [18].

The main technical result of this paper is the following.

Theorem 3.2. *In the Laver model, if $X \subseteq 2^\omega$ has cardinality $\mathfrak{b}$, then there is a Borel map $f : 2^\omega \rightarrow \omega^\omega$ such that $f[X]$ is unbounded.*

Proof. The notation in this proof is as in Laver [11]. We will use the following slightly simplified version of Lemma 14 of [11].

Lemma 3.3 (Laver). *Let $\mathbb{P}_{\omega_2}$ be the countable support iteration of Laver forcing, $p \in \mathbb{P}_{\omega_2}$, and let $\dot{a}$ be a $\mathbb{P}_{\omega_2}$ -name such that*

$$p \Vdash \dot{a} \in 2^\omega.$$

Then there are a condition q stronger than p and finite $U_s \subseteq 2^\omega$ for each $s \in q(0)$ extending the root of $q(0)$ such that for all such s and all n :

$$q(0)_t \wedge q \restriction [1, \omega_2) \Vdash “\exists u \in \check{U}_s \ u \restriction n = \dot{a} \restriction n”$$

for all but finitely many immediate successors t of s in $q(0)$.

Assume that $X \subseteq 2^\omega$ has no unbounded Borel image in $\mathcal{M}[G_{\omega_2}]$, i.e., Laver's model. For every code $u \in 2^\omega$ for a Borel function $f : 2^\omega \rightarrow \omega^\omega$ there exists $g \in \omega^\omega$ such that for every $x \in X$ we have that $f(x) \leq^* g$.

By a standard Löwenheim-Skolem argument (see Theorem 4.5 on page 281 of [3] or section 4 on page 580 of [12]), we may find $\alpha < \omega_2$ such that for every code $u \in \mathcal{M}[G_\alpha]$ there is an upper bound $g \in \mathcal{M}[G_\alpha]$. By the arguments employed by Laver [11, Lemmata 10 and 11], we may assume that $\mathcal{M}[G_\alpha]$ is the ground model $\mathcal{M}$.

Since the continuum hypothesis holds in $\mathcal{M}$ and $|X| = \mathfrak{b} = \aleph_2$, there are $p \in G_{\omega_2}$ and $\dot{a}$ such that

$$p \Vdash \dot{a} \in \check{X} \text{ and } \dot{a} \notin \mathcal{M}.$$

Work in the ground model $\mathcal{M}$.

Let $q \leq p$ be as in Lemma 3.3. Define

$$Q = \{s \in q(0) : \text{root}(q(0)) \subseteq s\}$$

and let U_s , $s \in Q$, be the finite sets from the lemma. Let $U = \bigcup_{s \in Q} U_s$. Define a Borel map $f : 2^\omega \rightarrow \omega^Q$ so that for every $x \in 2^\omega \setminus U$ and for each $s \in Q$: If

$f(x)(s) = n$, then $x \restriction n \neq u \restriction n$ for each $u \in U_s$. For $x \in U$, $f(x)$ may be arbitrary. There must be a $g \in \omega^Q \cap \mathcal{M}$ and $r \leq q$ such that

$$r \Vdash f(\check{a}) \leq^* \check{g}.$$

Since p forced that a is not in the ground model, it cannot be that a is in U . We may extend $r(0)$ if necessary so that if $s = \text{root}(r(0))$, then

$$r \Vdash f(\check{a})(s) \leq \check{g}(s).$$

But this is a contradiction to Lemma 3.3, since for all but finitely many $t \in r(0)$ which are immediate extensions of s :

$$r(0)_t \wedge q \restriction [1, \omega_2) \Vdash f(\check{a})(s) > \check{g}(s). \quad \square$$

In [20], Tsaban and Zdomskyy prove that $S_1(\Gamma, \Gamma)$ for Borel covers is equivalent to the Kočinac property $S_{\text{cof}}(\Gamma, \Gamma)$ [10], asserting that for all $\mathcal{U}_0, \mathcal{U}_1, \dots \in \Gamma$, there are cofinite subsets $\mathcal{V}_0 \subseteq \mathcal{U}_0, \mathcal{V}_1 \subseteq \mathcal{U}_1, \dots$ such that $\bigcup_n \mathcal{V}_n \in \Gamma$. The main result of [5] can be reformulated as follows.

Theorem 3.4 (Dow [5]). *In Laver's model, $S_1(\Gamma, \Gamma)$ implies $S_{\text{cof}}(\Gamma, \Gamma)$.*

For the reader's convenience, we give Dow's proof, adapted to the present notation.

Proof. A family $\mathcal{H} \subseteq [\omega]^\omega$ is ω -splitting if for each countable $\mathcal{A} \subseteq [\omega]^\omega$, there is $H \in \mathcal{H}$ which splits each element of $\mathcal{A}$, i.e.,

$$|A \cap H| = |A \setminus H| = \omega \text{ for all } A \in \mathcal{A}.$$

The main technical result in [5] is the following.

Lemma 3.5 (Dow). *In Laver's model, each ω -splitting family contains an ω -splitting family of cardinality $< \mathfrak{b}$.*

Assume that X satisfies $S_1(\Gamma, \Gamma)$. Let $\mathcal{U}_0, \mathcal{U}_1, \dots$ be open point-cofinite countable covers of X . We may assume⁴ that $\mathcal{U}_i \cap \mathcal{U}_j = \emptyset$ whenever $i \neq j$. Put $\mathcal{U} = \bigcup_{n < \omega} \mathcal{U}_n$. We identify $\mathcal{U}$ with ω , its cardinality.

Define $\mathcal{H} \subseteq [\mathcal{U}]^\omega$ as follows. For $H \in [\mathcal{U}]^\omega$, put $H \in \mathcal{H}$ if and only if there exists $\mathcal{V} \in [\mathcal{U}]^\omega$, a point-cofinite cover of X , such that $H \cap \mathcal{U}_n \subseteq^* \mathcal{V}$ for all n . We claim that $\mathcal{H}$ is an ω -splitting family. As $\mathcal{H}$ is closed under taking infinite subsets, it suffices to show that it is ω -hitting; i.e., for any countable $\mathcal{A} \subseteq [\mathcal{U}]^\omega$ there exists $H \in \mathcal{H}$ which intersects each $A \in \mathcal{A}$. (It is enough to intersect each $A \in \mathcal{A}$, since we may assume that $\mathcal{A}$ is closed under taking cofinite subsets.)

Let $\mathcal{A} \subseteq [\mathcal{U}]^\omega$ be countable. For each n , choose sets $\mathcal{U}_{n,m} \in [\mathcal{U}_n]^\omega$, $m \in \omega$, such that for each $A \in \mathcal{A}$, if $A \cap \mathcal{U}_n$ is infinite, then $\mathcal{U}_{n,m} \subseteq A$ for some m . Apply the $S_1(\Gamma, \Gamma)$ to the family $\{\mathcal{U}_{n,m} : n, m \in \omega\}$ to obtain a point-cofinite $\mathcal{V} \subseteq \mathcal{U}$ such that $\mathcal{V} \cap \mathcal{U}_{n,m}$ is nonempty for all n, m .

Next, choose finite subsets $\mathcal{F}_n \subseteq \mathcal{U}_n$, $n \in \omega$, such that for each $A \in \mathcal{A}$ with $A \cap \mathcal{U}_n$ finite for all n , then $A \subseteq^* \bigcup_n \mathcal{F}_n$. Take $H = \mathcal{V} \cup \bigcup_n \mathcal{F}_n$. Then H is in $\mathcal{H}$ and meets each $A \in \mathcal{A}$. This shows that $\mathcal{H}$ is an ω -splitting family.

By Lemma 3.5, there is an ω -splitting $\mathcal{H}' \subseteq \mathcal{H}$ of cardinality $< \mathfrak{b}$. For each $H \in \mathcal{H}'$, let $\mathcal{V}_H$ witness that H is in $\mathcal{H}$; i.e., $\mathcal{V}_H \subseteq \mathcal{U}$ is a point-cofinite cover of X and $H \cap \mathcal{U}_n \subseteq^* \mathcal{V}_H$ for all n .

⁴To see why, replace each $\mathcal{U}_n$ by $\mathcal{U}_n \setminus \bigcup_{i < n} \mathcal{U}_i$ and discard the finite ones. It suffices to show that $S_{\text{cof}}(\Gamma, \Gamma)$ applies to those that are left.

By the definition of $\mathfrak{b}$, we may find finite $\mathcal{F}_n \subseteq \mathcal{U}_n$, $n \in \omega$, such that for each $H \in \mathcal{H}'$,

$$H \cap \mathcal{U}_n \subseteq \mathcal{V}_H \cup \mathcal{F}_n$$

for all but finitely many n . We claim that $\mathcal{W} = \bigcup_n \mathcal{U}_n \setminus \mathcal{F}_n$ is point-cofinite. Suppose it is not. Then there is $x \in X$ such that for infinitely many n , there is $U_n \in \mathcal{U}_n \setminus \mathcal{F}_n$ with $x \notin U_n$. Let $H \in \mathcal{H}'$ contain infinitely many of these U_n . By the above inclusion, all but finitely many of these U_n are in $\mathcal{V}_H$. This contradicts the fact that $\mathcal{V}_H$ is point-cofinite. $\square$

We therefore have the following.

Theorem 3.6. *In Laver's model, each set of reals X satisfying $\mathbf{S}_1(\Gamma, \Gamma)$ has cardinality less than $\mathfrak{b}$.*

Proof. By Dow's Theorem, $\mathbf{S}_1(\Gamma, \Gamma)$ implies $\mathbf{S}_{\text{cof}}(\Gamma, \Gamma)$, which in turn implies $\mathbf{S}_1(\Gamma, \Gamma)$ for Borel covers [20]. The latter property is equivalent to having all Borel images in ω^ω bounded [18]. Apply Theorem 3.2. $\square$

Thus, it is consistent that strong measure zero and $\mathbf{S}_1(\Gamma, \Gamma)$ are both trivial.

The proof of Dow's Theorem 3.4 becomes more natural after replacing, in Lemma 3.5 “ ω -splitting” by “ ω -hitting”. This is possible, due to the following fact (cf. Remark 4 of [5]).

Proposition 3.7. *For each infinite cardinal κ , the following are equivalent:*

- (1) *Each ω -splitting family contains an ω -splitting family of cardinality $< \kappa$.*
- (2) *Each ω -hitting family contains an ω -hitting family of cardinality $< \kappa$.*

Proof. (1 $\Rightarrow$ 2) Suppose $\mathcal{A}$ is an ω -hitting family. Let $\mathcal{B} = \bigcup_{A \in \mathcal{A}} [A]^\omega$. Then $\mathcal{B}$ is ω -splitting. By (1) there exists $\mathcal{C} \subseteq \mathcal{B}$ of size $< \kappa$ which is ω -splitting. Choose $\mathcal{D} \subseteq \mathcal{A}$ of size $< \kappa$ such that for every $C \in \mathcal{C}$ there exists $D \in \mathcal{D}$ with $C \subseteq D$. Then $\mathcal{D}$ is ω -hitting.

(2 $\Rightarrow$ 1) Suppose $\mathcal{A}$ is an ω -splitting family. For each $A \subseteq \omega$ define

$$A^* = \{2n : n \in A\} \cup \{2n+1 : n \in \overline{A}\}.$$

Then the family $\mathcal{A}^* = \{A^* : A \in \mathcal{A}\}$ is ω -hitting. To see this, suppose that $\mathcal{B}$ is countable. Without loss we may assume that $\mathcal{B} = \mathcal{B}_0 \cup \mathcal{B}_1$, where each element of $\mathcal{B}_0$ is a subset of the evens and each element of $\mathcal{B}_1$ is a subset of the odds. For $B \in \mathcal{B}_0$ let $C_B = \{n : 2n \in B\}$ and for $B \in \mathcal{B}_1$ let $C_B = \{n : 2n+1 \in B\}$. Now put

$$\mathcal{C} = \{C_B : B \in \mathcal{B}\}.$$

Since $\mathcal{A}$ is ω -splitting there is $A \in \mathcal{A}$ which splits $\mathcal{C}$. If $B \in \mathcal{B}_0$, then $A \cap C_B$ infinite implies $B \cap A^*$ infinite. If $B \in \mathcal{B}_1$, then $\overline{A} \cap C_B$ infinite implies $B \cap A^*$ infinite.

By (2) there exists $\mathcal{A}_0 \subseteq \mathcal{A}$ of cardinality $< \kappa$ such that $\mathcal{A}_0^*$ is ω -hitting. We claim that $\mathcal{A}_0$ is ω -splitting. Given any $B \subseteq \omega$ let $B' = \{2n : n \in B\}$ and let $B'' = \{2n+1 : n \in B\}$. Given $\mathcal{B} \subseteq [\omega]^\omega$ countable, there exists $A \in \mathcal{A}_0$ such that A^* hits each B' and B'' for $B \in \mathcal{B}$. But this implies that A splits $\mathcal{B}$. $\square$

4. APPLICATIONS TO ARHANGEL'SKII'S α_i SPACES

Let Y be a general (not necessarily metrizable) topological space. We say that a countably infinite set $A \subseteq Y$ converges to a point $y \in Y$ if each (equivalently, some) bijective enumeration of A converges to y . The following concepts are due

to Arhangel'skiĭ. Y is an α_1 space if for each $y \in Y$ and each sequence $A_0, A_1, \dots$ of countably infinite sets, each converging to y , there are cofinite $B_0 \subseteq A_0, B_1 \subseteq A_1, \dots$, such that $\bigcup_n B_n$ converges to y . Replacing “cofinite” by “singletons” (or equivalently, by “infinite”), we obtain the definition of an α_2 space.

We first consider countable spaces.

Definition 4.1. Let X be a set of reals, and let $\mathcal{U}_0, \mathcal{U}_1, \dots$ be countable point-cofinite covers of X . For each n , enumerate bijectively $\mathcal{U}_n = \{U_m^n : m \in \omega\}$. We associate to X a (new) topology τ on the fan $S_\omega = \omega \times \omega \cup \{\infty\}$ as follows: ∞ is the only nonisolated point of S_ω , and a neighborhood base at ∞ is given by the sets

$$[\infty]_F = \{(n, m) : F \subseteq U_m^n\}$$

for each finite $F \subseteq X$.

Lemma 4.2. *In the notation of Definition 4.1: A converges to ∞ in τ if and only if $\mathcal{U}(A) = \{U_m^n : (n, m) \in A\}$ is a point-cofinite cover of X .* $\square$

Assume that there is an unbounded tower. By Corollary 2.10, there is a set of reals X satisfying $S_1(\Gamma, \Gamma)$ but not $S_{\text{cof}}(\Gamma, \Gamma)$. Let $\mathcal{U}_0, \mathcal{U}_1, \dots$ be countable open point-cofinite covers of X witnessing the failure of $S_{\text{cof}}(\Gamma, \Gamma)$. Then, by Lemma 4.2, (S_ω, τ) is α_2 but not α_1 . In particular, we reproduce the following.

Corollary 4.3 (Nyikos [13]). *If there is an unbounded tower of cardinality $\mathfrak{b}$, then there is a countable α_2 space which is not an α_1 space.* $\square$

Recall that by Proposition 2.4, it suffices to assume in Corollary 4.3 the existence of any unbounded tower.

Next, we consider spaces of continuous functions. Consider $C(X)$, the family of continuous real-valued functions, as a subspace of the Tychonoff product $\mathbb{R}^X$, i.e., with the topology of pointwise convergence. Sakai [14] proved that X satisfies $S_1(\Gamma, \Gamma)$ for clopen covers if and only if $C(X)$ is an α_2 space. The main result of [20] is that $C(X)$ is α_1 if and only if X satisfies $S_1(\Gamma, \Gamma)$ for Borel covers (equivalently, each Borel image of X in ω^ω is bounded).

The *Scheepers Conjecture* is that for subsets of $\mathbb{R} \setminus \mathbb{Q}$, $S_1(\Gamma, \Gamma)$ for clopen covers implies $S_1(\Gamma, \Gamma)$ for open covers. Dow [5] proved that in Laver's model, every α_2 space is α_1 . By Theorem 3.2, we can add the last item in the following list.

Corollary 4.4. *In Laver's model, the following are equivalent for sets of reals X :*

- (1) $C(X)$ is an α_2 space;
- (2) $C(X)$ is an α_1 space;
- (3) X satisfies $S_1(\Gamma, \Gamma)$ for clopen covers;
- (4) X satisfies $S_1(\Gamma, \Gamma)$ for open covers;
- (5) X satisfies $S_1(\Gamma, \Gamma)$ for Borel covers;
- (6) $|X| < \mathfrak{b}$.

$\square$

On the other hand, Corollary 2.10 implies the following.

Corollary 4.5. *If there is an unbounded tower, then there is a set of reals X such that $C(X)$ is α_2 but not α_1 .* $\square$

Essentially, Corollary 4.3 is a special case of Corollary 2.10, whereas Corollary 4.5 is equivalent to Corollary 2.10.

ACKNOWLEDGMENT

The authors thank Lyubomyr Zdomskyy for his useful comments.

REFERENCES

- [1] T. Bartoszyński and S. Shelah, *Continuous images of sets of reals*, Topology and its Applications **116** (2001), 243–253. MR1855966 (2002f:03090)
- [2] T. Bartoszyński and B. Tsaban, *Hereditary topological diagonalizations and the Menger-Hurewicz Conjectures*, Proceedings of the American Mathematical Society, **134** (2006), 605–615. MR2176030 (2006f:54038)
- [3] J. Baumgartner and R. Laver, *Iterated perfect-set forcing*, Annals of Mathematical Logic **17** (1979), 271–288. MR556894 (81a:03050)
- [4] A. Blass, *Combinatorial cardinal characteristics of the continuum*, in: **Handbook of Set Theory** (M. Foreman, A. Kanamori, and M. Magidor, eds.), Kluwer Academic Publishers, Dordrecht, to appear. <http://www.math.lsa.umich.edu/~ablass/hbk.pdf>
- [5] A. Dow, *Two classes of Fréchet-Urysohn spaces*, Proceedings of the American Mathematical Society **108** (1990), 241–247. MR975638 (90j:54028)
- [6] F. Galvin and A. Miller, γ -sets and other singular sets of real numbers, Topology and its Applications **17** (1984), 145–155. MR738943 (85f:54011)
- [7] J. Gerlits and Zs. Nagy, *Some properties of $C(X)$. I*, Topology and its Applications **14** (1982), 151–161. MR667661 (84f:54021)
- [8] W. Hurewicz, *Über eine Verallgemeinerung des Borelschen Theorems*, Mathematische Zeitschrift **24** (1925), 401–421.
- [9] W. Just, A. Miller, M. Scheepers, and P. Szeptycki, *The combinatorics of open covers. II*, Topology and its Applications **73** (1996), 241–266. MR1419798 (98g:03115a)
- [10] L. Kočinac, *Selection principles related to α_i -properties*, Taiwanese Journal of Mathematics **12** (2008), 561–572. MR2417134 (2009h:91049)
- [11] R. Laver, *On the consistency of Borel’s conjecture*, Acta Mathematica **137** (1976), 151–169. MR0422027 (54:10019)
- [12] A. Miller, *Mapping a set of reals onto the reals*, Journal of Symbolic Logic **48** (1983), 575–584. MR716618 (84k:03125)
- [13] P. Nyikos, *Subsets of ω^ω and the Fréchet-Urysohn and α_i -properties*, Topology and its Applications **48** (1992), 91–116. MR1195504 (93k:54011)
- [14] M. Sakai, *The sequence selection properties of $C_p(X)$* , Topology and its Applications **154** (2007), 552–560. MR2280899 (2007k:54007)
- [15] M. Scheepers, *Combinatorics of open covers. I: Ramsey theory*, Topology and its Applications **69** (1996), 31–62. MR1378387 (97h:90123)
- [16] M. Scheepers, *$C_p(X)$ and Arhangel’skiĭ’s α_i spaces*, Topology and its Applications **89** (1998), 265–275. MR1645184 (99g:54018)
- [17] M. Scheepers, *Sequential convergence in $C_p(X)$ and a covering property*, East-West Journal of Mathematics **1** (1999), 207–214. MR1727383 (2000i:54014)
- [18] M. Scheepers and B. Tsaban, *The combinatorics of Borel covers*, Topology and its Applications **121** (2002), 357–382. MR1908999 (2003e:03091)
- [19] B. Tsaban, *Menger’s and Hurewicz’s Problems: Solutions from “The Book” and refinements*, Contemporary Mathematics, American Mathematical Society, to appear.
- [20] B. Tsaban and L. Zdomskyy, *Hereditarily Hurewicz spaces and Arhangel’skiĭ sheaf amalgamations*, Journal of the European Mathematical Society, to appear.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WISCONSIN-MADISON, VAN VLECK HALL, 480 LINCOLN DRIVE, MADISON, WISCONSIN 53706-1388

E-mail address: miller@math.wisc.edu

URL: <http://www.math.wisc.edu/~miller/>

DEPARTMENT OF MATHEMATICS, BAR-ILAN UNIVERSITY, RAMAT-GAN 52900, ISRAEL

E-mail address: tsaban@math.biu.ac.il

URL: <http://www.cs.biu.ac.il/~tsaban>