THE TORSION INDEX OF A p -COMPACT GROUP

JAUME AGUADÉ

(Communicated by Brooke Shipley)

ABSTRACT. We extend the theory of torsion indices of compact connected Lie groups to p -compact groups and compute these indices in all cases.

1. INTRODUCTION AND STATEMENT OF RESULTS

The torsion index of a compact connected Lie group was defined by Grothendieck in 1958 ([10]) and has been investigated by several authors ([14], [6], [15], etc.). Recently, the computation of the torsion indices of all simply connected compact Lie groups has been completed (see [16]). Since we are going to work at a single prime p , instead of the torsion index of a Lie group G , we want to consider its p -primary part $t_p(G)$. We summarize the properties of $t_p(G)$ which are relevant to the present work in the following proposition ($\mathbb{Z}_p$ denotes the ring of p -adic integers).

Theorem 1.1. *Let p be a prime and let G be a compact connected Lie group with a maximal torus T and corresponding Weyl group W . The positive integer $t_p(G)$ has the following properties:*

- (TI1) *If A is a finite abelian p -subgroup of G , then A has a subgroup of index dividing $t_p(G)$ which is contained in a conjugate of T .*
- (TI2) *$t_p(G)$ kills the kernel and the cokernel of the homomorphism*

$$H^*(BG; \mathbb{Z}_p) \rightarrow H^*(BT; \mathbb{Z}_p)^W.$$

- (TI3) *$H^*(G/T; \mathbb{Z}_p)$ is torsion free and concentrated in even degrees $\leq N = \dim(G) - \text{rank}(G)$, with $H^N(G/T; \mathbb{Z}_p) \cong \mathbb{Z}_p$. Then, $t_p(G)$ is the order of the cokernel of $H^N(BT; \mathbb{Z}_p) \rightarrow H^N(G/T; \mathbb{Z}_p)$.*
- (TI4) *If p is not a torsion prime for G , then $t_p(G) = 1$.*

Notice that the property (TI3) can be taken as a *definition* of the (p -primary) torsion index $t_p(G)$. The other properties are well known and can be found in [15], which provides proofs or references for all of them. Actually, the properties above are usually stated using $H^*(-; \mathbb{Z})$ and $t(G) = \prod_p t_p(G)$ instead of $H^*(-; \mathbb{Z}_p)$ and $t_p(G)$, but it is easy to see that both formulations are indeed equivalent. For property (TI2) one should notice that $H^*(BT; \mathbb{Z}_p)^W = H^*(BT; \mathbb{Z})^W \otimes \mathbb{Z}_p$. This follows from exactness of $- \otimes \mathbb{Z}_p$ and the fact that the elements invariant under W can be viewed as the kernel of the homomorphism $\bigoplus_{g \in W} (1 - g)$.

Received by the editors February 5, 2009.

2010 *Mathematics Subject Classification*. Primary 55P35, 57T15.

The author is partially supported by grants MTM2007-61545 and SGR2005-00606.

©2010 American Mathematical Society
 Reverts to public domain 28 years from publication

The purpose of this paper is to extend the theorem above to connected p -compact groups ([8]) and to compute the torsion indices in all cases. We prove:

Theorem 1.2. *Let p be a prime and let X be a connected p -compact group with maximal torus T and corresponding Weyl group W . There is an integer $t_p(X)$ such that:*

- (1) *The properties (TI1), (TI2), (TI3), (TI4) in Theorem 1.1 hold after replacing G with X .*
- (2) *If X is exotic, then $t_p(X) = 1$ for p odd and $t_2(X) = 2$.*

Here we use the work *exotic* with the same meaning as in [1]: A p -compact group X is exotic if the associated pseudoreflexion representation of the Weyl group of X over the p -adic field is irreducible and does not come from a reflection group over $\mathbb{Z}$.

Section 2 deals with the (easier) odd prime case, and we show that if we define $t_p(X) = 1$ for any exotic X , then properties (TI1), (TI2), (TI3), (TI4) hold true. The hardest part consists of computing the torsion index of the only exotic 2-compact group, which we (following [12]) denote G_3 (other authors denote it as $DI(4)$). We need a comprehensive review of the cohomology of G_3 and BG_3 (section 3) and some computations on the cohomology of the exotic homogeneous space $G_3/\text{Spin}(7)$ (section 4) before we can prove that $t_2(G_3) = 2$. Finally, we prove Theorem 1.2 in section 6.

2. THE ODD PRIME CASE

The classification theorem for p -compact groups ([2]) tells us that any connected p -compact group X splits uniquely as a product $X \cong G_p^\wedge \times X_1$, where G is a compact connected Lie group and X_1 is a product of exotic p -compact groups. Notice that the splitting is as p -compact groups and not just as spaces. This splitting implies that it is enough to prove Theorem 1.2 for each exotic p -compact group, since it is already known to be true for the (p -completions of) compact connected Lie groups. Let us discuss this in some more detail. If Theorem 1.2 holds for the p -compact groups X_1 and X_2 , let $X = X_1 \times X_2$ and let us define $t_p(X) = t_p(X_1)t_p(X_2)$. We need to check that properties (TI1) to (TI4) hold for X if they hold for X_1 and X_2 . (TI4) is trivial and (TI3) is straightforward. To prove (TI2) let us observe that the kernel of $\gamma : H^*(BX; \mathbb{Z}_p) \rightarrow H^*(BT; \mathbb{Z}_p)^W$ is equal to the torsion elements in $H^*(BX; \mathbb{Z}_p)$. If X is of Lie type, this is well known (cf. [9]). If X is exotic and $p = 2$ (i.e. $X = G_3$), then this is assertion 4 in [12]; and if p is odd, this is proven in [1]. Then, it is clear that $t_p(X_1)t_p(X_2)$ kills the kernel of γ . It is obvious that $t_p(X_1)t_p(X_2)$ kills the cokernel of γ as well. Finally, (TI1) follows easily since we can use the theory of kernels of homomorphisms between p -compact groups which is developed in [8], section 7.

Let us assume now that p is odd and let X be an exotic p -compact group. These objects are very well understood. In particular, they satisfy the following properties (see [1]). Let T and W denote a maximal torus of X and the corresponding Weyl group, respectively. Then:

- (1) X is simply connected and center free and $H^*(X; \mathbb{Z}_p)$ is torsion free.
- (2) The natural map $BT \rightarrow BX$ induces an isomorphism

$$H^*(BX; \mathbb{Z}_p) \cong H^*(BT; \mathbb{Z}_p)^W.$$

In particular, $H^*(BX; \mathbb{Z}_p)$ is concentrated in even degrees.

- (3) $H^*(X/T; \mathbb{Z}_p)$ is a free $\mathbb{Z}_p$ -module concentrated in even degrees. Moreover (see [13], th. 7.5.1) $H^*(X/T; \mathbb{Z}_p) \otimes \mathbb{Q}$ is a Poincaré duality algebra with fundamental class in degree $\dim(X) - \text{rank}(X)$. Actually, as a W -module, $H^*(X/T; \mathbb{Z}_p) \otimes \mathbb{Q}$ coincides with the regular representation of W .

We also need another property of p -compact groups (which holds also for $p = 2$) that follows from the work in [5].

- (4) If X is any p -compact group such that $H^*(BX; \mathbb{F}_p)$ is concentrated in even degrees, then any finite abelian p -subgroup of X is conjugated to a subgroup of the maximal torus of X . In particular, this holds for any product of exotic p -compact groups for p odd.

Theorem 1.2 for p odd follows immediately from all these properties of p -compact groups. $\square$

3. THE 2-COMPACT GROUP G_3 AND ITS MAXIMAL TORUS

In this section we recollect several properties of G_3 that we need in the forthcoming sections. We state these properties without proof because either they can be found in the papers [7], [12], [4], [11] or they follow from straightforward computations that are left to the reader.

As is well known, G_3 is an exotic connected 2-compact group of rank three whose Weyl group W is the reflection group number 24 in the Shephard-Todd list of finite complex reflection groups. Its existence was established by Dwyer and Wilkerson in [7]. We remind the reader that some authors call this 2-compact group $DI(4)$, but we follow the notation used in [12]. As an abstract group, W is isomorphic to $\mathbb{Z}/2\mathbb{Z} \times GL_3(\mathbb{F}_2)$ and for a maximal torus T of G_3 , there is a basis $\{\epsilon_1, \epsilon_2, \epsilon_3\}$ of $H^2(BT; \mathbb{Z}_2)$ such that the action of W on $H^*(BT; \mathbb{Z}_2)$ is given by the pseudoreflections

$$s_1 = \begin{pmatrix} -1 & -\bar{\alpha} & 1 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad s_2 = \begin{pmatrix} 1 & 0 & 0 \\ -\alpha & -1 & 1 \\ 0 & 0 & 1 \end{pmatrix}, \quad s_3 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 1 & 1 & -1 \end{pmatrix},$$

where $\alpha, \bar{\alpha} \in \mathbb{Z}_2$ are the roots of $x^2 - x + 2$ chosen in such a way that α is odd and $\bar{\alpha}$ is even.

G_3 has $\text{Spin}(7)$ as a 2-compact subgroup of maximal rank. This means that there is a map $\phi : B\text{Spin}(7)_2^\wedge \rightarrow BG_3$ whose homotopical fibre is $\mathbb{F}_2$ -finite. It is natural to denote this fibre by $G_3/\text{Spin}(7)$. The restriction of ϕ to a maximal torus of $\text{Spin}(7)$ is a maximal torus of G_3 .

There is a subgroup $V \subset \text{Spin}(7)$ (explicitly described in [7]) which is an elementary abelian 2-group of rank four and such that the homomorphisms

$$H^*(BG_3; \mathbb{F}_2) \xrightarrow{\phi^*} H^*(B\text{Spin}(7); \mathbb{F}_2) \xrightarrow{k^*} H^*(BV; \mathbb{F}_2) \cong \mathbb{F}_2[V^*]$$

are monomorphisms (k^* is induced by the inclusion $V \subset \text{Spin}(7)$). Moreover, the image of $(\phi k)^*$ coincides with the rank four Dickson algebra which is the algebra of invariants of $H^*(BV; \mathbb{F}_2)$ under the action of the full linear group $GL(V^*)$, and the image of k^* coincides with the algebra of invariants $H^*(BV; \mathbb{F}_2)^H$ where $H \subset GL(V^*)$ can be described, in some appropriate basis of V^* , as the set of matrices with first row equal to $(1, 0, 0, 0)$. These algebras of invariants are well

known (also as algebras over the Steenrod algebra) and we have isomorphisms (subscripts denote degrees)

$$\begin{aligned} H^*(BG_3; \mathbb{F}_2) &\cong \mathbb{F}_2[c_8, c_{12}, c_{14}, c_{15}], \\ H^*(B\mathrm{Spin}(7); \mathbb{F}_2) &\cong \mathbb{F}_2[d_4, d_6, d_7, d_8], \end{aligned}$$

where the generators c_i and d_i can be explicitly described. In particular, we can see that ϕ^* is given by $\phi^*(c_8) = d_4^2 + d_8$, $\phi^*(c_{12}) = d_6^2 + d_4d_8$, $\phi^*(c_{14}) = d_7^2 + d_6d_8$, $\phi^*(c_{15}) = d_7d_8$. Sq^1 vanishes on d_4, d_7, d_8 , while $Sq^1(d_6) = d_7$.

As was said before, a maximal torus T of $\mathrm{Spin}(7)$ is also a maximal torus of G_3 . We have maps

$$BT_2^\wedge \xrightarrow{i} B\mathrm{Spin}(7)_2^\wedge \xrightarrow{\phi} BG_3$$

and we can view the Weyl group W_1 of $\mathrm{Spin}(7)$ as a subgroup of W , namely $W_1 = \langle s_1, s_2, s_1s_3s_2s_1s_2s_3s_1 \rangle$. It is known that the homomorphism

$$i^* : H^*(B\mathrm{Spin}(7); \mathbb{Z}_2) \rightarrow H^*(BT; \mathbb{Z}_2)^{W_1}$$

is surjective and its kernel coincides with the ideal of torsion elements. The integral invariants of W_1 are computed in [4]. They turn out to form a polynomial algebra on generators of degrees 4, 8, 12:

$$H^*(BT; \mathbb{Z}_2)^{W_1} \cong \mathbb{Z}_2[u_4, u_8, u_{12}].$$

Choosing an appropriate basis $\{x_1, x_2, A\}$ of $H^2(BT; \mathbb{Z}_2)$, these generators are

$$\begin{aligned} u_4 &= (1/2)(x_1^2 + x_2^2 + x_3^2), \\ u_8 &= (1/16)(x_1^4 + x_2^4 + x_3^4 - 2x_1^2x_2^2 - 2x_1^2x_3^2 - 2x_2^2x_3^2), \\ u_{12} &= x_1^2x_2^2x_3^2, \end{aligned}$$

where we have used the notation $x_3 = 2A - x_1 - x_2$, and one can check that in spite of the denominators, these polynomials belong to $\mathbb{Z}_2[x_1, x_2, A]$.

The generators u_4, u_8 and u_{12} have a rather simple form as polynomials on x_1, x_2, A , but this basis of $H^2(BT; \mathbb{Z}_2)$ does not coincide with the basis $\{\epsilon_1, \epsilon_2, \epsilon_3\}$ that we have used to describe the action of W on $H^*(BT; \mathbb{Z}_2)$. The matrix that expresses $\{\epsilon_1, \epsilon_2, \epsilon_3\}$ in terms of $\{x_1, x_2, A\}$ is

$$\begin{pmatrix} 0 & -\bar{\alpha}/2 & -(1+\alpha)/2 \\ 1 & 0 & -(1+\alpha)/2 \\ 0 & \bar{\alpha} & \alpha \end{pmatrix} \in GL_3(\mathbb{Z}_2).$$

Using this matrix we can express the generators u_4, u_8, u_{12} as polynomials in $\epsilon_1, \epsilon_2, \epsilon_3$ and so we have an explicit description of the homomorphism

$$\mathbb{Z}_2[u_4, u_8, u_{12}] = H^*(B\mathrm{Spin}(7); \mathbb{Z}_2)/\mathrm{Torsion} \rightarrow H^*(BT; \mathbb{Z}_2) = \mathbb{Z}_2[\epsilon_1, \epsilon_2, \epsilon_3].$$

Finally, we want to use this to describe the homomorphism

$$\mathbb{F}_2[d_4, d_6, d_7, d_8] = H^*(B\mathrm{Spin}(7); \mathbb{F}_2) \xrightarrow{i^*} H^*(BT; \mathbb{F}_2) = \mathbb{F}_2[\epsilon_1, \epsilon_2, \epsilon_3].$$

In the Bockstein spectral sequence for $B\mathrm{Spin}(7)$ we have $E_2 = E_\infty = \mathbb{F}_2[\bar{d}_4, \bar{d}_8, \bar{d}_6^2]$, and the surjection

$$j : H^*(B\mathrm{Spin}(7); \mathbb{Z}_2)/\mathrm{Torsion} \rightarrow E_\infty$$

is given by $j(u_4) = \bar{d}_4$, $j(u_8) = \bar{d}_8$, $j(u_{12}) = \bar{d}_6^2$. From this it is straightforward to perform the computations that yield

$$\begin{aligned} i^*(d_4) &= \epsilon_1^2 + \epsilon_1\epsilon_2 + \epsilon_2^2, \\ i^*(d_6) &= Sq^2 i^*(d_4) = \epsilon_1^2\epsilon_2 + \epsilon_1\epsilon_2^2, \\ i^*(d_7) &= 0, \\ i^*(d_8) &= \epsilon_1\epsilon_2\epsilon_3(\epsilon_1 + \epsilon_2 + \epsilon_3) + \epsilon_3^2(\epsilon_1 + \epsilon_2 + \epsilon_3)^2. \end{aligned}$$

4. THE EXOTIC HOMOGENEOUS SPACE $G_3/\text{Spin}(7)$

In this section we want to investigate the cohomology of the exotic homogeneous space $G_3/\text{Spin}(7)$. The computations presented here are probably known to experts, but it may be worthwhile to work them out here in some detail.

Let us consider the fibration $G_3/\text{Spin}(7) \xrightarrow{p} B\text{Spin}(7)_2^\wedge \rightarrow BG_3$ and let $V \subset \text{Spin}(7)$ denote the elementary abelian 2-group of rank 4 considered in the preceding section. To simplify the notation, let us write $S = H^*(BV; \mathbb{F}_2)$. Then, we have $H^*(B\text{Spin}(7); \mathbb{F}_2) = S^H$ and $H^*(BG_3; \mathbb{F}_2) = S^G$ for $G = GL_4(\mathbb{F}_2)$.

The computation of $H^*(G_3/\text{Spin}(7); \mathbb{F}_2)$ is best worked out with the Eilenberg-Moore spectral sequence

$$\text{Tor}_{H^*(BG_3; \mathbb{F}_2)}(H^*(B\text{Spin}(7); \mathbb{F}_2), \mathbb{F}_2) \Rightarrow H^*(G_3/\text{Spin}(7); \mathbb{F}_2).$$

Here the key observation is that $H^*(B\text{Spin}(7); \mathbb{F}_2) = S^H$ is a free module over $H^*(BG_3; \mathbb{F}_2) = S^G$ because of the following classic argument. S is an integral extension of S^G ; hence S^H is also an integral extension of S^G and, since S^H is a finitely generated algebra, we obtain that S^H is a finitely generated S^G -module. But both S^H and S^G are polynomial algebras, and we can apply [3], Chap. V, 5.5, or [13], 6.7.1, to conclude that S^H is S^G -free.

Hence the Eilenberg-Moore spectral sequence collapses to an isomorphism

$$H^*(G_3/\text{Spin}(7); \mathbb{F}_2) \cong \mathbb{F}_2[\bar{d}_4, \bar{d}_6, \bar{d}_7]/(\bar{d}_6^2 + \bar{d}_4^3, \bar{d}_7^2 + \bar{d}_4^2\bar{d}_6, \bar{d}_4^2\bar{d}_7),$$

where $\bar{d}_4, \bar{d}_6, \bar{d}_7$ are the images of $d_4, d_6, d_7 \in H^*(B\text{Spin}(7); \mathbb{F}_2)$, respectively. It is rather easy to completely work out the algebra structure of $H^*(G_3/\text{Spin}(7); \mathbb{F}_2)$. We obtain the following:

- (1) The Poincaré series of $H^*(G_3/\text{Spin}(7); \mathbb{F}_2)$ is

$$1 + t^4 + t^6 + t^7 + t^8 + t^{10} + t^{11} + t^{12} + t^{13} + t^{14} + t^{16} + t^{17} + t^{18} + t^{20} + t^{24}$$

and the Euler characteristic is $7 = [W : H]$.

- (2) An additive basis for $H^*(G_3/\text{Spin}(7); \mathbb{F}_2)$ is given by

$$\left\{ \bar{d}_4^i, i = 0, \dots, 6, \bar{d}_6, \bar{d}_7, \bar{d}_4\bar{d}_6, \bar{d}_4\bar{d}_7, \bar{d}_6\bar{d}_7, \bar{d}_4^2\bar{d}_6, \bar{d}_4\bar{d}_6\bar{d}_7, \bar{d}_4^3\bar{d}_6 \right\}.$$

- (3) $H^*(G_3/\text{Spin}(7); \mathbb{F}_2)$ is a Poincaré duality algebra with top class $\bar{d}_4^6$ (see [13], 6.5).

- (4) The Bockstein spectral sequence of $H^*(G_3/\text{Spin}(7); \mathbb{F}_2)$ collapses after the second term; i.e. $H^*(G_3/\text{Spin}(7); \mathbb{Z}_2)$ has only torsion of order 2. We have

$$H^*(G_3/\text{Spin}(7); \mathbb{Z}_2)/\text{Torsion} \cong \mathbb{Z}_2[\bar{a}]/\bar{a}^7$$

and

$$H^*(G_3/\text{Spin}(7); \mathbb{Z}_2) \cong \mathbb{Z}_2[\bar{a}, \bar{c}]/(\bar{a}^7, \bar{c}^3, \bar{a}^2\bar{c}, 2\bar{c}).$$

In particular, the top class in $H^*(G_3/\mathrm{Spin}(7); \mathbb{Z}_2)$ is $\bar{d}_4^6$ in dimension 24, and it is in the image of

$$\phi^* : H^*(B\mathrm{Spin}(7); \mathbb{Z}_2) \rightarrow H^*(G_3/\mathrm{Spin}(7); \mathbb{Z}_2).$$

5. THE TORSION INDEX OF G_3

To compute the torsion index of the 2-compact group G_3 we need a lemma on Poincaré duality in fibrations. I'm grateful to Aniceto Murillo for some helpful conversations on this subject. For this lemma we use the following notation. Let $\mathcal{O}$ denote the ring of integers or the ring of p -adic integers. Cohomology is taken with coefficients in $\mathcal{O}$, and we assume that all spaces are of finite type over $\mathcal{O}$. We say that $\eta \in H^n(X)$ is an *orientation class* if $H^i(X) = 0$ for $i > n$, $H^n(X) \cong \mathcal{O}$, and η is a generator of $H^n(X)$.

Lemma 5.1. *Let $F \xrightarrow{j} E \xrightarrow{\pi} B$ be a fibration of 1-connected spaces and assume that $\eta^F \in H^m(F)$ and $\eta^B \in H^n(B)$ are orientation classes. Assume $\alpha \in H^m(E)$ is such that $j^*(\alpha) = \lambda \eta^F$ for some $\lambda \neq 0$. Then there is an orientation class η^E for E such that $\alpha \cdot \pi^*(\eta^B) = \lambda \eta^E$.*

Proof. This follows easily from the cohomology spectral sequence of the fibration $F \xrightarrow{j} E \xrightarrow{\pi} B$. First of all, it is clear that $H^i(E) = 0$ for $i > n + m$ while $H^{n+m}(E) = E_\infty^{n,m} = E_2^{n,m} \cong \mathcal{O}$. Recall that the cohomology spectral sequence is multiplicative in the sense that (up to some signs which would not play any role here) the product in E_2 induced by the products in $H^*(B)$ and $H^*(F)$ yields a product in each E_r , $2 \leq r \leq \infty$, in such a way that the product in E_∞ is compatible with the product in $H^*(E)$.

At the E_2 level we have that $\eta^E := \eta^F \cdot \eta^B$ is a generator of $E_2^{n,m} = E_\infty^{n,m} = H^{n+m}(E)$. The hypothesis $j^*(\alpha) = \lambda \eta^F$, $\lambda \neq 0$ implies that α has filtration zero in $H^m(E)$ and its image in $E_\infty^{0,m}$ is $\lambda \eta^F$. Then, $\lambda \eta^E = (\lambda \eta^F) \cdot [\eta^B]$ holds in E_∞ where $[\eta^B]$ denotes the image of η^B in $E_\infty^{0,0}$. Since $E_\infty^{i,m+n-i} = 0$ for $i \neq n$, we deduce $\lambda \eta^E = \alpha \cdot \pi^*(\eta^B)$, as desired. $\square$

Now we can proceed to the computation of the torsion index of G_3 or, to be more precise, to the computation of the order of the cokernel of $k^* : H^{42}(BT; \mathbb{Z}_2) \rightarrow H^{42}(G_3/T; \mathbb{Z}_2)$. We consider the diagram

$$\begin{array}{ccccc} (\mathrm{Spin}(7)/T)_2^\wedge & \xrightarrow{j} & G_3/T & \xrightarrow{\pi} & G_3/\mathrm{Spin}(7) \\ & & k \downarrow & & \phi \downarrow \\ & & (BT)_2^\wedge & \xrightarrow{i} & (B\mathrm{Spin}(7))_2^\wedge \end{array}$$

$\mathrm{Spin}(7)/T$ is a compact orientable differentiable manifold of dimension 18, and we can choose an orientation class $\eta \in H^{18}(\mathrm{Spin}(7)/T; \mathbb{Z}_2)$. The torsion indices of the Lie groups $\mathrm{Spin}(n)$ have been computed by Totaro for all values of n ([15]), and it turns out that the torsion index of $\mathrm{Spin}(7)$ is equal to 2. This means that there is $\omega \in H^*(BT; \mathbb{Z})$ such that $f^*(\omega) = 2\eta$ for the natural map $f : \mathrm{Spin}(7)/T \rightarrow BT$.

The computations in the preceding section show that there is an orientation class $\rho \in H^{24}(G_3/\mathrm{Spin}(7); \mathbb{Z}_2)$ which is in the image of ϕ^* . Let $\rho = \phi^*(\gamma)$. We can now apply the lemma above to the fibration $\mathrm{Spin}(7)/T \rightarrow G_3/T \rightarrow G_3/\mathrm{Spin}(7)$ with $\alpha = k^*(\omega)$ and deduce that there is an orientation class $\theta \in H^{42}(G_3/T; \mathbb{Z}_2)$ such that $k^*(\omega \cdot i^*(\gamma)) = 2\theta$. This implies that the torsion index of G_3 divides 2.

Next, we prove that the torsion index of G_3 cannot be equal to 1. It is enough to prove that the homomorphism $H^{42}(BT; \mathbb{F}_2) \rightarrow H^{42}(G_3/T; \mathbb{F}_2)$ is equal to zero. Let us consider the $\mathbb{F}_2$ -spectral sequence of the fibration $G_3 \rightarrow G_3/T \rightarrow BT_2^\wedge$. We have that

$$\begin{aligned} H^*(G_3; \mathbb{F}_2) &\cong \mathbb{F}_2[x_7]/x_7^4 \otimes E(x_{11}, x_{13}), \\ Sq^4(x_7) &= x_{11}, \quad Sq^2(x_{11}) = x_{13}, \quad Sq^1(x_{13}) = x_7^2. \end{aligned}$$

Hence, the generators $x_7, x_{11}, x_{13}, x_7^2$ are transgressive to $c_8, c_{12}, c_{14}, 0$, respectively. Here we denote by c_8, c_{12}, c_{14} the images in $H^*(BT; \mathbb{F}_2)$ of the generators $c_8, c_{12}, c_{14} \in H^*(BG_3; \mathbb{F}_2)$. Recall that in section 3 we have computed these elements as explicit polynomials in some basis $\{\epsilon_1, \epsilon_2, \epsilon_3\}$ of $H^2(BT; \mathbb{F}_2)$.

In the E_2 -term of the spectral sequence of $G_3 \rightarrow G_3/T \rightarrow BT_2^\wedge$, let us consider the row containing x_7^2 . All elements in this row are permanent cycles, and the only boundaries are the elements of the form $x_7^2 q$ with q in the ideal of $\mathbb{F}_2[\epsilon_1, \epsilon_2, \epsilon_3]$ generated by c_8, c_{12}, c_{14} . If we compute the quotient algebra $\mathbb{F}_2[\epsilon_1, \epsilon_2, \epsilon_3]/(c_8, c_{12}, c_{14})$ (using any choice of computer algebra software), we see that it is a graded algebra with Poincaré series equal to

$$\begin{aligned} 1 + 3t^2 + 6t^4 + 10t^6 + 14t^8 + 18t^{10} + 21t^{12} + 22t^{14} \\ + 21t^{16} + 18t^{18} + 14t^{20} + 10t^{22} + 6t^{24} + 3t^{26} + t^{28}, \end{aligned}$$

and so in particular there is an element $q \in H^{28}(BT; \mathbb{F}_2)$ which does not belong to the ideal (c_8, c_{12}, c_{14}) . Hence, the element $x_7^2 q$ in the E_2 -term of the spectral sequence survives to a nontrivial element in $H^{42}(G_3/T; \mathbb{F}_2)$ which cannot be in the image of $H^*(BT; \mathbb{F}_2)$. This finishes the proof of

Theorem 5.2. *The cokernel of $H^{42}(BT; \mathbb{Z}_2) \rightarrow H^{42}(G_3/T; \mathbb{Z}_2)$ has order two.*

6. PROOF OF THEOREM 1.2

In section 2 we saw that it is enough to prove Theorem 1.2 for each exotic p -compact group, and we also saw that Theorem 1.2 is true for all odd primes. Since it is known ([2]) that the only exotic 2-compact group is G_3 , the only thing that remains to be proved is that G_3 satisfies the properties (TI1) to (TI4) with $t_2(G_3) = 2$.

(TI4) is void, and (TI3) is just Theorem 5.2 plus some facts about G_3/T which were proven in [2]. In [12] it is proven that the torsion elements in $H^*(BG_3; \mathbb{Z}_2)$ are of order two and the homomorphism $H^*(BG_3; \mathbb{Z}_2) \rightarrow H^*(BT; \mathbb{Z}_2)^W$ is surjective. This implies immediately that (TI2) holds. Let A be a nontrivial finite abelian 2-subgroup of G_3 and let E be a subgroup of A of order 2. Then, A factors through the centralizer of E in G_3 which is $\text{Spin}(7)$. Since $\text{Spin}(7)$ has 2-torsion index equal to 2, we deduce that A has a subgroup of index at most 2 which is included in a maximal torus of G_3 . So, we have (TI1), and the proof is complete. $\square$

REFERENCES

1. K.K.S. Andersen, J. Grodal, J.M. Møller, A. Viruel, *The classification of p -compact groups for p odd*, Ann. of Math. (2) 167 (2008), 95–210. MR2373153 (2009a:55012)
2. K.K.S. Andersen, J. Grodal, *The classification of 2-compact groups*, J. Amer. Math. Soc. 22 (2009), no. 2, 387–436. MR2476779
3. N. Bourbaki, *Groupes et algèbres de Lie*, Ch. 4, 5 et 6. Act. Sci. Ind. 1337, Paris, 1968. MR0240238 (39:1590)

4. D.J. Benson, J.A. Wood, *Integral invariants and cohomology of $B\text{Spin}(n)$* , *Topology* 34 (1995), 13–28. MR1308487 (95m:55023)
5. F.-X. Dehon, J. Lannes, *Sur les espaces fonctionnels dont la source est le classifiant d'un groupe de Lie compact commutatif*, *Inst. Hautes Études Sci. Publ. Math.*, No. 89 (1999), 127–177. MR1793415 (2001m:55038)
6. M. Demazure, *Invariants symétriques entiers des groupes de Weyl et torsion*, *Invent. Math.* 21 (1973), 287–301. MR0342522 (49:7268)
7. W. G. Dwyer, C. W. Wilkerson, *A new finite loop space at the prime 2*, *J. Amer. Math. Soc.* 6 (1993), 37–64. MR1161306 (93d:55011)
8. W. G. Dwyer, C. W. Wilkerson, *Homotopy fixed point methods for Lie groups and finite loop spaces*, *Ann. of Math. (2)* 139 (1994), 395–442. MR1274096 (95e:55019)
9. M. Feshbach, *The image of $H^*(BG; \mathbb{Z})$ in $H^*(BT; \mathbb{Z})$ for G a compact Lie group with maximal torus T* , *Topology* 20 (1981), 93–95. MR592571 (82g:55019)
10. A. Grothendieck, *La torsion homologique et les sections rationnelles*, in *Anneaux de Chow et applications*, Séminaire C. Chevalley, 1958, 2nd year, Secrétariat math., Paris, exp. 5.
11. D. Notbohm, *On the 2-compact group $DI(4)$* , *J. Reine Angew. Math.* 555 (2003), 163–185. MR1956596 (2003k:55018)
12. A. Osse, U. Suter, *Invariant theory and the K-theory of the Dwyer-Wilkerson space*, *Contemp. Math.*, 265, Amer. Math. Soc., Providence, RI, 2000, 17–185. MR1803957 (2001j:55005)
13. L. Smith, *Polynomial Invariants of Finite Groups*. A K Peters, Wellesley, Mass., 1995. MR1328644 (96f:13008)
14. J. Tits, *Sur les degrés des extensions de corps déployant les groupes algébriques simples*, *C. R. Acad. Sci. Paris Ser. I. Math.* 315 (1992), 1131–1138. MR1194504 (93m:20059)
15. B. Totaro, *The torsion index of the spin groups*, *Duke Math. J.* 129 (2005), 249–290. MR2165543 (2006f:57039b)
16. B. Totaro, *The torsion index of E_8 and other groups*, *Duke Math. J.* 129 (2005), 219–248. MR2165542 (2006f:57039a)

DEPARTAMENT DE MATEMÀTIQUES, UNIVERSITAT AUTÒNOMA DE BARCELONA, 08193 CERDANYOLA DEL VALLÈS, SPAIN

E-mail address: aguade@mat.uab.cat