ON THE SPATIAL ASYMPTOTICS OF SOLUTIONS OF THE ABLOWITZ–LADIK HIERARCHY

JOHANNA MICHOR

(Communicated by Peter A. Clarkson)

Dedicated with great pleasure to Peter W. Michor on the occasion of his 60th birthday

ABSTRACT. We show that for decaying solutions of the Ablowitz–Ladik system, the leading asymptotic term is time independent. In addition, two arbitrary bounded solutions of the Ablowitz–Ladik system which are asymptotically close at the initial time stay close. All results are also derived for the associated hierarchy.

1. INTRODUCTION

When solving completely integrable wave equations via the inverse scattering transform, a method developed by Gardner et al. [12] in 1967 for the Korteweg–de Vries (KdV) equation, one intends to prove the existence of solutions within the respective class. In particular, short-range perturbations of the background solution should remain short-range during the time evolution. So to what extent are spatial asymptotical properties time independent?

For the KdV equation, this question was answered by Bondareva and Shubin [9], [10], who considered the Cauchy problem for initial conditions which have a prescribed asymptotic expansion in terms of powers of the spatial variable and showed that the leading term of the expansion is time independent. Teschl [18] considered the initial value problem for the Toda lattice in the class of decaying solutions and obtained time independence of the leading term.

In this note we want to address the same question for the Ablowitz–Ladik (AL) system, an integrable discretization of the AKNS-ZS system derived by Ablowitz and Ladik ([3]–[6]) in the mid seventies. The AL system is given by

$$(1.1) \quad \begin{aligned} -i\alpha_t - (1 - \alpha\beta)(\alpha^- + \alpha^+) + 2\alpha &= 0, \\ -i\beta_t + (1 - \alpha\beta)(\beta^- + \beta^+) - 2\beta &= 0, \end{aligned}$$

where $\alpha = \alpha(n, t)$, $\beta = \beta(n, t)$, $(n, t) \in \mathbb{Z} \times \mathbb{R}$, are complex-valued sequences and $f^\pm(n, t) = f(n \pm 1, t)$. In the defocusing ($\beta = \bar{\alpha}$) and focusing case ($\beta = -\bar{\alpha}$), (1.1) is a discrete analog of the nonlinear Schrödinger (NLS) equation

$$iq_t + q_{xx} \pm 2q|q|^2 = 0.$$

Received by the editors September 16, 2009.

2010 *Mathematics Subject Classification.* Primary 37K40, 37K15; Secondary 35Q55, 37K10.

Key words and phrases. Spatial asymptotics, Ablowitz–Ladik hierarchy.

This research was supported by the Austrian Science Fund (FWF) under Grant No. V120.

©2010 American Mathematical Society
 Reverts to public domain 28 years from publication

We refer to the monographs [2], [7], or [13] for further information.

Our main result in Theorem 2.4 yields that the dominant term of suitably decaying solutions $\alpha(n, t)$, $\beta(n, t)$ of (1.1), for instance weighted ℓ^{2p} sequences whose spatial difference is in ℓ^p , $1 \leq p < \infty$, is time independent. For example,

$$(1.2) \quad \alpha(n, t) = \frac{a}{n^\delta} + O\left(\frac{1}{n^{\min(2\delta, \delta+1)}}\right), \quad \beta(n, t) = \frac{b}{n^\delta} + O\left(\frac{1}{n^{\min(2\delta, \delta+1)}}\right), \quad n \rightarrow \infty,$$

holds for fixed t , provided it holds at the initial time $t = t_0$. Here $a, b \in \mathbb{C}$ and $\delta \geq 0$. A similar expression is valid for $n \rightarrow -\infty$.

The inverse scattering transform for the AL system with vanishing boundary conditions was studied in [4]. Ablowitz, Biondini, and Prinari [1] (compare also Vekslerchik and Konotop [19]) considered nonvanishing steplike boundary conditions $\alpha(n) \rightarrow \alpha_0 e^{i\theta_\pm}$ as $|n| \rightarrow \infty$, $\alpha_0 > 0$, in the class

$$(1.3) \quad \sum_{j=n}^{\pm\infty} (\alpha(j) - \alpha_0 e^{i\theta_\pm}) < \infty$$

for the defocusing discrete NLS equation. Quasi-periodic boundary conditions for the AL hierarchy will be considered in Michor [17]. As mentioned, a crucial step is to show that short-range perturbations such as (1.3) of solutions stay short-range. Here we show in general that arbitrary bounded solutions of the AL system which are asymptotically close at the initial time stay close.

In Section 2 we derive our results for the AL system and extend them in Section 3 to the AL hierarchy, a completely integrable hierarchy of nonlinear evolution equations whose first nonlinear member is (1.1).

2. THE INITIAL VALUE PROBLEM FOR THE ABLOWITZ–LADIK SYSTEM

Let us begin by recalling some basic facts of the system (1.1). We will only consider bounded solutions and hence require

Hypothesis H.2.1. *Suppose that $\alpha, \beta: \mathbb{Z} \times \mathbb{R} \rightarrow \mathbb{C}$ satisfy*

$$(2.1) \quad \sup_{(n,t) \in \mathbb{Z} \times \mathbb{R}} (|\alpha(n, t)| + |\beta(n, t)|) < \infty, \\ \alpha(n, \cdot), \beta(n, \cdot) \in C^1(\mathbb{R}), \quad n \in \mathbb{Z}, \quad \alpha(n, t)\beta(n, t) \notin \{0, 1\}, \quad (n, t) \in \mathbb{Z} \times \mathbb{R}.$$

The AL system (1.1) is equivalent to the zero-curvature equation

$$(2.2) \quad U_t + UV - V^+U = 0,$$

where

$$(2.3) \quad U(z) = \begin{pmatrix} z & \alpha \\ \beta z & 1 \end{pmatrix}, \quad V(z) = i \begin{pmatrix} z - 1 - \alpha\beta^- & \alpha - \alpha^- z^{-1} \\ \beta^- z - \beta & 1 + \alpha^- \beta - z^{-1} \end{pmatrix}$$

for the spectral parameter $z \in \mathbb{C} \setminus \{0\}$. The AL system can also be formulated in terms of Lax pairs; see [15]. Then (1.1) is equivalent to the Lax equation

$$(2.4) \quad \frac{d}{dt} L(t) - [P(t), L(t)] = 0, \quad t \in \mathbb{R},$$

$$(2.5) \quad L = \begin{pmatrix} \ddots & \ddots & & \ddots & \ddots & & & & & \\ & 0 & -\alpha(0)\rho(-1) & -\beta(-1)\alpha(0) & -\alpha(1)\rho(0) & \rho(0)\rho(1) & & & & 0 \\ & \rho(-1)\rho(0) & \beta(-1)\rho(0) & -\beta(0)\alpha(1) & \beta(0)\rho(1) & 0 & & & & \\ & & 0 & -\alpha(2)\rho(1) & -\beta(1)\alpha(2) & -\alpha(3)\rho(2) & \rho(2)\rho(3) & & & \\ 0 & & & \rho(1)\rho(2) & \beta(1)\rho(2) & -\beta(2)\alpha(3) & \beta(2)\rho(3) & 0 & & \\ & & & & \ddots & \ddots & \ddots & \ddots & \ddots & \ddots \end{pmatrix}$$
$$P = \frac{i}{2}(L_+ - L_- + (L^{-1})_- - (L^{-1})_+ + 2Q_d).$$
$$(2.6) \quad L_{\pm} = (L_{\pm}(m, n))_{(m, n) \in \mathbb{Z}^2}, \quad L_{\pm}(m, n) = \begin{cases} L(m, n), & \pm(n - m) > 0, \\ 0, & \text{otherwise.} \end{cases}$$
$$L(s) = W(s, t)L(t)W(s, t)^{-1}, \quad s, t \in \mathbb{R}.$$

Theorem 2.2. *Let $t_0 \in \mathbb{R}$ and suppose $(\alpha_0, \beta_0) \in M = \ell^p(\mathbb{Z}) \oplus \ell^p(\mathbb{Z})$ for some $p \in [1, \infty) \cup \{\infty\}$. Then there exists $T > 0$ and a unique local integral curve $t \mapsto (\alpha(t), \beta(t))$ in $C^\infty((t_0 - T, t_0 + T), M)$ of the Ablowitz–Ladik system (1.1) such that $(\alpha, \beta)|_{t=t_0} = (\alpha_0, \beta_0)$.*

$$(2.7) \quad \|(\alpha, \beta)\|_{w,p} = \begin{cases} \left(\sum_{n \in \mathbb{Z}} w(n) (|\alpha(n)|^p + |\beta(n)|^p) \right)^{1/p}, & 1 \leq p < \infty, \\ \sup_{n \in \mathbb{Z}} w(n) (|\alpha(n)| + |\beta(n)|), & p = \infty. \end{cases}$$
$$(2.8) \quad \|(\alpha(t) - \tilde{\alpha}(t), \beta(t) - \tilde{\beta}(t))\|_{w,p} < \infty$$

But even the leading term is preserved by the time evolution.

Theorem 2.4. *Let $w(n) \geq 1$ be some weight with $\sup_n (|\frac{w(n+1)}{w(n)}| + |\frac{w(n)}{w(n+1)}|) < \infty$. Fix $1 \leq p \leq \infty$ and suppose α_0, β_0 and $\tilde{\alpha}_0, \tilde{\beta}_0$ are bounded sequences such that*

$$\begin{aligned} & \|(\alpha_0, \beta_0)\|_{w,2p} < \infty, \quad \|(\alpha_0 - \alpha_0^+, \beta_0 - \beta_0^+)\|_{w,p} < \infty, & \text{if } 1 \leq p < \infty, \\ & \|(\tilde{\alpha}_0, \tilde{\beta}_0)\|_{w,p} < \infty, \\ & \|(\alpha_0, \beta_0)\|_{w,\infty} < \infty, \quad \|(\alpha_0 - \alpha_0^+, \beta_0 - \beta_0^+)\|_{w^2,\infty} < \infty, & \text{if } p = \infty. \\ & \|(\tilde{\alpha}_0, \tilde{\beta}_0)\|_{w^2,\infty} < \infty, \end{aligned}$$

Let $(\alpha(t), \beta(t))$, $t \in (-T, T)$, be the unique solution of the Ablowitz–Ladik system (1.1) corresponding to the initial conditions

$$(2.9) \quad \alpha(0) = \alpha_0 + \tilde{\alpha}_0, \quad \beta(0) = \beta_0 + \tilde{\beta}_0.$$

Then this solution is of the form

$$(2.10) \quad \alpha(t) = \alpha_0 + \tilde{\alpha}(t), \quad \beta(t) = \beta_0 + \tilde{\beta}(t),$$

where $\|(\tilde{\alpha}(t), \tilde{\beta}(t))\|_{w,p} < \infty$, respectively, $\|(\tilde{\alpha}(t), \tilde{\beta}(t))\|_{w^2,\infty} < \infty$.

Proof. The proof relies on the idea to consider our differential equation in two nested spaces of sequences, the Banach space of all $(\alpha(n), \beta(n))$ with sup norm, and the Banach space with norm $\|\cdot\|_{w,p}$, as follows. Plugging $(\alpha_0 + \tilde{\alpha}(t), \beta_0 + \tilde{\beta}(t))$ into the AL equations (1.1) yields a differential equation for $(\tilde{\alpha}(t), \tilde{\beta}(t))$:

$$\begin{aligned} (2.11) \quad i\tilde{\alpha}_t(t) &= -(1 - (\alpha_0 + \tilde{\alpha}(t))(\beta_0 + \tilde{\beta}(t)))(\tilde{\alpha}^+(t) + \alpha_0^+ + \tilde{\alpha}^-(t) + \alpha_0^-) + 2\tilde{\alpha}(t) + 2\alpha_0 \\ &= \alpha_0 - \alpha_0^- + \alpha_0 - \alpha_0^+ + \alpha_0\beta_0(\alpha_0^+ + \alpha_0^-) \\ &\quad + \tilde{\alpha}(t)(2 + (\beta_0 + \tilde{\beta}(t))(\tilde{\alpha}^+(t) + \alpha_0^+ + \tilde{\alpha}^-(t) + \alpha_0^-)) \\ &\quad + \tilde{\beta}(t)\alpha_0(\tilde{\alpha}^+(t) + \alpha_0^+ + \tilde{\alpha}^-(t) + \alpha_0^-) \\ &\quad + \tilde{\alpha}^+(t)(\alpha_0\beta_0 - 1) + \tilde{\alpha}^-(t)(\alpha_0\beta_0 - 1), \\ i\tilde{\beta}_t(t) &= (1 - (\alpha_0 + \tilde{\alpha}(t))(\beta_0 + \tilde{\beta}(t)))(\tilde{\beta}^+(t) + \beta_0^+ + \tilde{\beta}^-(t) + \beta_0^-) - 2\tilde{\beta}(t) - 2\beta_0 \\ &= \beta_0^+ - \beta_0 + \beta_0^- - \beta_0 - \alpha_0\beta_0(\beta_0^+ + \beta_0^-) \\ &\quad - \tilde{\beta}(t)(2 + (\alpha_0 + \tilde{\alpha}(t))(\tilde{\beta}^+(t) + \beta_0^+ + \tilde{\beta}^-(t) + \beta_0^-)) \\ &\quad - \tilde{\alpha}(t)\beta_0(\tilde{\beta}^+(t) + \beta_0^+ + \tilde{\beta}^-(t) + \beta_0^-) \\ &\quad - \tilde{\beta}^+(t)(\alpha_0\beta_0 - 1) - \tilde{\beta}^-(t)(\alpha_0\beta_0 - 1). \end{aligned}$$

The requirement on $w(n)$ implies that the shift operators are continuous with respect to the norm $\|\cdot\|_{w,p}$ and the same is true for the multiplication operator with a bounded sequence. Therefore, using the generalized Hölder inequality yields that (2.11) is a system of inhomogeneous linear differential equations in the Banach space with norm $\|\cdot\|_{w,p}$ and has a local solution with respect to this norm (see e.g. [11] for the theory of ordinary differential equations in Banach spaces). Since $w(n) \geq 1$, this solution is bounded and the corresponding coefficients $(\tilde{\alpha}, \tilde{\beta})$ coincide with the solution (α, β) of the AL system (1.1) from Theorem 2.2.

Moreover, $(\tilde{\alpha}(t), \tilde{\beta}(t))$ are uniformly bounded for $t \in (-T, T)$, as writing (2.11) in integral form yields

$$\|(\tilde{\alpha}(t), \tilde{\beta}(t))\|_{w,p} \leq \|(\tilde{\alpha}(0), \tilde{\beta}(0))\|_{w,p} + tC\|(\alpha_0, \beta_0)\|_{w,2p} + C \int_0^t \|(\tilde{\alpha}(s), \tilde{\beta}(s))\|_{w,p} ds$$

for some constants C . $\square$

Example (1.2) in the introduction follows if we let $\tilde{\alpha}_0 = \tilde{\beta}_0 \equiv 0$ and

$$\alpha_0(n) = \frac{a}{n^\delta}, \quad \beta_0(n) = \frac{b}{n^\delta}, \quad a, b \in \mathbb{C}, \quad \delta \geq 0,$$

for $n > 0$, $\alpha_0(n) = \beta_0(n) = 0$ for $n \leq 0$. Now choose $p = \infty$ with

$$w(n) = \begin{cases} (1+n)^{\min(\delta, (\delta+1)/2)}, & n > 0, \\ 1, & n \leq 0, \end{cases}$$

and apply Theorem 2.4.

Finally, we remark that if a solution $(\alpha(n, t), \beta(n, t))$ vanishes at two consecutive points $n = n_0, n = n_0 + 1$ in an arbitrarily small time interval $t \in (t_1, t_2)$, then it vanishes identically for all (n, t) in $\mathbb{Z} \times \mathbb{R}$; see [16]. In particular, a compact support of the solution is not preserved. The corresponding result for the AL hierarchy is derived in [16] as well.

3. EXTENSION TO THE ABLOWITZ–LADIK HIERARCHY

In this section we show how our results extend to the AL hierarchy. The hierarchy can be constructed by generalizing the matrix $V(z)$ in the zero-curvature equation (2.2) to a 2×2 matrix $V_{\underline{r}}(z)$, $\underline{r} = (r_-, r_+) \in \mathbb{N}_0^2$, with Laurent polynomial entries; see [13, Sec. 3.2] or [14]. Suppose that $U(z)$ and $V_{\underline{r}}(z)$ satisfy the zero-curvature equation

$$(3.1) \quad U_t + UV_{\underline{r}} - V_{\underline{r}}^+ U = 0.$$

Then the coefficients $\{f_{\ell, \pm}\}_{\ell=0, \dots, r_{\pm}-1}$, $\{g_{\ell, \pm}\}_{\ell=0, \dots, r_{\pm}}$, and $\{h_{\ell, \pm}\}_{\ell=0, \dots, r_{\pm}-1}$ of the Laurent polynomials in the entries of $V_{\underline{r}}(z)$ are recursively defined by

$$(3.2) \quad \begin{aligned} g_{0,+} &= \frac{1}{2}c_{0,+}, \quad f_{0,+} = -c_{0,+}\alpha^+, \quad h_{0,+} = c_{0,+}\beta, \\ g_{\ell+1,+} - g_{\ell+1,+}^- &= \alpha h_{\ell,+} + \beta f_{\ell,+}, \quad 0 \leq \ell \leq r_+ - 1, \\ f_{\ell+1,+}^- &= f_{\ell,+} - \alpha(g_{\ell+1,+} + g_{\ell+1,+}^-), \quad 0 \leq \ell \leq r_+ - 2, \\ h_{\ell+1,+} &= h_{\ell,+} + \beta(g_{\ell+1,+} + g_{\ell+1,+}^-), \quad 0 \leq \ell \leq r_+ - 2, \end{aligned}$$

and

$$(3.3) \quad \begin{aligned} g_{0,-} &= \frac{1}{2}c_{0,-}, \quad f_{0,-} = c_{0,-}\alpha, \quad h_{0,-} = -c_{0,-}\beta^+, \\ g_{\ell+1,-} - g_{\ell+1,-}^- &= \alpha h_{\ell,-} + \beta f_{\ell,-}^-, \quad 0 \leq \ell \leq r_- - 1, \\ f_{\ell+1,-} &= f_{\ell,-}^- + \alpha(g_{\ell+1,-} + g_{\ell+1,-}^-), \quad 0 \leq \ell \leq r_- - 2, \\ h_{\ell+1,-}^- &= h_{\ell,-} - \beta(g_{\ell+1,-} + g_{\ell+1,-}^-), \quad 0 \leq \ell \leq r_- - 2. \end{aligned}$$

Note that $g_{\ell, \pm}$ are only defined up to summation constants $\{c_{\ell, \pm}\}_{\ell=0, \dots, r_{\pm}}$ by the difference equations in (3.2), (3.3). In addition, the zero-curvature equation (3.1)

is equivalent to

$$0 = i \begin{pmatrix} 0 & -i\alpha_t - \alpha(g_{r_+,+} + g_{r_-,-}^-) \\ z(-i\beta_t + \beta(g_{r_+,+}^- + g_{r_-,-}) & +f_{r_+-1,+} - f_{r_- -1,-}^- \\ -h_{r_- -1,-} + h_{r_+-1,+}^- & 0 \end{pmatrix}.$$

Varying $\underline{r} \in \mathbb{N}_0^2$, the collection of evolution equations

$$(3.4) \quad \text{AL}_{\underline{r}}(\alpha, \beta) = \begin{pmatrix} -i\alpha_t - \alpha(g_{r_+,+} + g_{r_-,-}^-) + f_{r_+-1,+} - f_{r_- -1,-}^- \\ -i\beta_t + \beta(g_{r_+,+}^- + g_{r_-,-}) - h_{r_- -1,-} + h_{r_+-1,+}^- \end{pmatrix} = 0, \\ t \in \mathbb{R}, \underline{r} = (r_-, r_+) \in \mathbb{N}_0^2,$$

then defines the time-dependent Ablowitz–Ladik hierarchy. Explicitly, taking $r_- = r_+$ for simplicity, the first few equations are:

$$(3.5) \quad \begin{aligned} \text{AL}_{(0,0)}(\alpha, \beta) &= \begin{pmatrix} -i\alpha_t - c_{(0,0)}\alpha \\ -i\beta_t + c_{(0,0)}\beta \end{pmatrix} = 0, \\ \text{AL}_{(1,1)}(\alpha, \beta) &= \begin{pmatrix} -i\alpha_t - \gamma(c_{0,-}\alpha^- + c_{0,+}\alpha^+) - c_{(1,1)}\alpha \\ -i\beta_t + \gamma(c_{0,+}\beta^- + c_{0,-}\beta^+) + c_{(1,1)}\beta \end{pmatrix} = 0, \\ \text{AL}_{(2,2)}(\alpha, \beta) &= \begin{pmatrix} -i\alpha_t - \gamma(c_{0,+}\alpha^{++}\gamma^+ + c_{0,-}\alpha^{--}\gamma^- \\ -\alpha(c_{0,+}\alpha^+\beta^- + c_{0,-}\alpha^-\beta^+) - \beta(c_{0,-}(\alpha^-)^2 + c_{0,+}(\alpha^+)^2)) \\ -i\beta_t + \gamma(c_{0,-}\beta^{++}\gamma^+ + c_{0,+}\beta^{--}\gamma^- \\ -\beta(c_{0,+}\alpha^+\beta^- + c_{0,-}\alpha^-\beta^+) - \alpha(c_{0,+}(\beta^-)^2 + c_{0,-}(\beta^+)^2)) \\ + \begin{pmatrix} -\gamma(c_{1,-}\alpha^- + c_{1,+}\alpha^+) - c_{(2,2)}\alpha \\ \gamma(c_{1,+}\beta^- + c_{1,-}\beta^+) + c_{(2,2)}\beta \end{pmatrix} \end{pmatrix} = 0, \text{ etc.,} \end{aligned}$$

where we abbreviated $c_{\underline{r}} = (c_{r,-} + c_{r,+})/2$ and $\gamma = 1 - \alpha\beta$. Different ratios of $c_{0,+}/c_{0,-}$ lead to different hierarchies. The AL system (1.1) corresponds to the case $\underline{r} = (1, 1)$, $c_{0,\pm} = 1$, and $c_{(1,1)} = -2$. The special choices $\beta = \pm\bar{\alpha}$, $c_{0,\pm} = 1$ lead to the discrete NLS hierarchy; the choices $\beta = \bar{\alpha}$, $c_{0,\pm} = \mp i$ yield the hierarchy of Schur flows. The AL hierarchy is invariant under the scaling transform

$$(3.6) \quad \{(\alpha(n), \beta(n))\}_{n \in \mathbb{Z}} \rightarrow \{(c\alpha(n), \beta(n)/c)\}_{n \in \mathbb{Z}}, \quad c \in \mathbb{C} \setminus \{0\}.$$

Hence choosing $c = e^{ic_{\underline{r}}t}$ it is no restriction to assume $c_{\underline{r}} = 0$.

By [15], the AL hierarchy is equivalent to the Lax equation

$$(3.7) \quad \frac{d}{dt}L(t) - [P_{\underline{r}}(t), L(t)] = 0, \quad t \in \mathbb{R}, \quad \underline{r} \in \mathbb{N}_0^2,$$

where L is the doubly infinite five-diagonal matrix (2.5) and (recall (2.6))

$$P_{\underline{r}} = \frac{i}{2} \sum_{\ell=1}^{r_+} c_{r_+-\ell,+} ((L^\ell)_+ - (L^\ell)_-) - \frac{i}{2} \sum_{\ell=1}^{r_-} c_{r_- - \ell,-} ((L^{-\ell})_+ - (L^{-\ell})_-) - \frac{i}{2} c_{\underline{r}} Q_d.$$

Since the AL flows are autonomous and $f_{r_{\pm}-1,\pm}$, $g_{r_{\pm},\pm}$, and $h_{r_{\pm}-1,\pm}$ depend polynomially on α, β and their shifts, [8, Thm. 4.1.5] implies local existence, uniqueness, and smoothness of the solution of the initial value problem of the hierarchy as well (see [13, Sec. 3.8], [15]).

Theorem 3.1. *Let $t_0 \in \mathbb{R}$ and suppose $\alpha_0, \beta_0 \in \ell^p(\mathbb{Z})$ for some $p \in [1, \infty) \cup \{\infty\}$. Then the $\underline{r}$ th Ablowitz–Ladik initial value problem*

$$(3.8) \quad \text{AL}_{\underline{r}}(\alpha, \beta) = 0, \quad (\alpha, \beta)|_{t=t_0} = (\alpha_0, \beta_0)$$

for some $\underline{r} \in \mathbb{N}_0^2$, has a unique, local, and smooth solution in time; that is, there exists a $T > 0$ such that $\alpha(\cdot), \beta(\cdot) \in C^\infty((t_0 - T, t_0 + T), \ell^p(\mathbb{Z}))$.

Next we show that short-range perturbations of bounded solutions remain short-range. In fact, we will be more general to include perturbations of steplike background solutions as for example (1.3).

Lemma 3.2. *Let $w(n) \geq 1$ be some weight with $\sup_n (|\frac{w(n+1)}{w(n)}| + |\frac{w(n)}{w(n+1)}|) < \infty$ and fix $1 \leq p \leq \infty$. Suppose $(\alpha(t), \beta(t))$ and $(\alpha_{\ell,r}(t), \beta_{\ell,r}(t))$ are arbitrary bounded solutions of some equation $\text{AL}_{\underline{r}}$ in the AL hierarchy and abbreviate*

$$(3.9) \quad \tilde{\alpha}(n, t) = \begin{cases} \alpha_r(n, t), & n \geq 0, \\ \alpha_\ell(n, t), & n < 0, \end{cases} \quad \tilde{\beta}(n, t) = \begin{cases} \beta_r(n, t), & n \geq 0, \\ \beta_\ell(n, t), & n < 0. \end{cases}$$

If

$$(3.10) \quad \|(\alpha(t) - \tilde{\alpha}(t), \beta(t) - \tilde{\beta}(t))\|_{w,p} < \infty$$

holds for one $t = t_0 \in \mathbb{R}$, then it holds for all $t \in (t_0 - T, t_0 + T)$.

Proof. Without loss we assume that $t_0 = 0$. First we derive the differential equation for the differences $\delta(n, t) = (\alpha(n, t) - \tilde{\alpha}(n, t), \beta(n, t) - \tilde{\beta}(n, t))$ in the Banach space of pairs of bounded sequences $\delta = (\delta_1, \delta_2)$ for which the norm $\|\delta\|_{w,p}$ is finite.

Let us show by induction on $r_\pm$ that $f_{r_\pm-1,\pm}(t) - \tilde{f}_{r_\pm-1,\pm}(t)$, $g_{r_\pm,\pm}(t) - \tilde{g}_{r_\pm,\pm}(t)$, and $h_{r_\pm-1,\pm}(t) - \tilde{h}_{r_\pm-1,\pm}(t)$ can be written as a linear combination of shifts of δ with the coefficients depending only on $(\alpha(t), \beta(t))$ and $(\alpha_{\ell,r}(t), \beta_{\ell,r}(t))$. It suffices to consider the homogeneous case where $c_{j,\pm} = 0$, $1 \leq j \leq r_\pm$, since all involved sums are finite. In this case, [14, Lemma A.3] yields that $f_{j,+}$, $g_{j,+}$, and $h_{j,+}$ can be recursively computed from $f_{0,+} = -\alpha^+$, $g_{0,+} = \frac{1}{2}$, and $h_{0,+} = \beta$ via

$$\begin{aligned} f_{\ell+1,+}^- &= f_{\ell,+} - \alpha(g_{\ell+1,+} + g_{\ell+1,+}^-), \\ h_{\ell+1,+} &= h_{\ell,+}^- + \beta(g_{\ell+1,+} + g_{\ell+1,+}^-), \\ g_{\ell+1,+} &= \sum_{k=0}^{\ell} f_{\ell-k,+} h_{k,+} - \sum_{k=1}^{\ell} g_{\ell+1-k,+} g_{k,+}, \end{aligned}$$

and similarly for the minus sign and $\tilde{f}_{j,\pm}$, $\tilde{g}_{j,\pm}$, and $\tilde{h}_{j,\pm}$. The fact that $(\tilde{\alpha}, \tilde{\beta})$ does not solve $\text{AL}_{\underline{r}}$ only affects finitely many terms and gives rise to an inhomogeneous term $B_{\underline{r}}(t)$ which is nonzero only for a finite number of terms.

Hence δ satisfies an inhomogeneous linear differential equation of the form

$$i \frac{d}{dt} \delta(t) = \sum_{|j| \leq \max(r_-, r_+)} A_{\underline{r},j}(t) (S^+)^j \delta(t) + B_{\underline{r}}(t).$$

Here $S^\pm(\delta_1(n, t), \delta_2(n, t)) = (\delta_1(n \pm 1, t), \delta_2(n \pm 1, t))$ are the shift operators,

$$A_{\underline{r},j}(n, t) = \begin{pmatrix} A_{\underline{r},j}^{11}(n, t) & A_{\underline{r},j}^{12}(n, t) \\ A_{\underline{r},j}^{21}(n, t) & A_{\underline{r},j}^{22}(n, t) \end{pmatrix}$$

are multiplication operators with bounded 2×2 matrix-valued sequences, and $B_{\underline{r}}(n, t) = (B_{\underline{r},1}(n, t), B_{\underline{r},2}(n, t))$ with $B_{\underline{r},i}(n, t) = 0$ for $|n| > \max(r_-, r_+)$. All entries of $A_{\underline{r},j}(t)$ and $B_{\underline{r}}(t)$ are polynomials with respect to $(\alpha(n+j, t), \beta(n+j, t))$, $(\alpha_{\ell,r}(n+j, t), \beta_{\ell,r}(n+j, t))$, $|j| \leq \max(r_-, r_+)$. Thus $\|B_{\underline{r}}(t)\|_{w,p} \leq D_{\underline{r}}$, where the constant depends only on the sup norms of $(\alpha(t), \beta(t))$ and $(\alpha_{\ell,r}(t), \beta_{\ell,r}(t))$. Moreover, the shift operators are continuous by our assumption since

$$\|S^{\pm}\| = \begin{cases} \sup_{n \in \mathbb{Z}} \left| \frac{w(n)}{w(n \pm 1)} \right|^{1/p}, & p \in [1, \infty), \\ \sup_{n \in \mathbb{Z}} \left| \frac{w(n)}{w(n \pm 1)} \right|, & p = \infty. \end{cases}$$

The same is true for the multiplication operators $A_{\underline{r},j}(t)$ whose norms depend only on the supremum of the entries by Hölder's inequality, that is, again on the sup norms of $(\alpha(t), \beta(t))$ and $(\alpha_{\ell,r}(t), \beta_{\ell,r}(t))$. Consequently, for $t \in (-T, T)$, there is a constant such that $\sum_{|j| \leq \max(r_-, r_+)} \|A_{\underline{r},j}(t)\| \|(S^+)^j\| \leq C_{\underline{r}}$. Hence

$$\|\delta(t)\|_{w,p} \leq \|\delta(0)\|_{w,p} + \int_0^t (C_{\underline{r}} \|\delta(s)\|_{w,p} + D_{\underline{r}}) ds,$$

and Gronwall's inequality implies that

$$\|\delta(t)\|_{w,p} \leq \|\delta(0)\|_{w,p} e^{C_{\underline{r}} t} + \frac{D_{\underline{r}}}{C_{\underline{r}}} (e^{C_{\underline{r}} t} - 1).$$

Since $w(n) \geq 1$, this solution is again bounded and hence coincides with the solution of the AL equation from Theorem 3.1. $\square$

For certain equations in the AL hierarchy, i.e. for certain configurations of summation coefficients $\{c_{j,\pm}\}$, our main result remains valid.

Theorem 3.3. *Let $\underline{r} = (r_-, r_+) \in \mathbb{N}_0^2 \setminus (0, 0)$ and assume that $c_{j,\pm} \in \mathbb{C}$, $j = 0, \dots, r_{\pm}$, satisfy*

$$(3.11) \quad \sum_{j=0}^{r_+-1} c_{j,+} + \sum_{j=0}^{r_--1} c_{j,-} = 0.$$

Let $w(n) \geq 1$ be some weight with $\sup_n (|\frac{w(n+1)}{w(n)}| + |\frac{w(n)}{w(n+1)}|) < \infty$. Fix $1 \leq p \leq \infty$ and suppose α_0, β_0 and $\tilde{\alpha}_0, \tilde{\beta}_0$ are bounded sequences such that

$$\begin{aligned} & \|(\alpha_0, \beta_0)\|_{w,2p} < \infty, \quad \|(\alpha_0 - \alpha_0^+, \beta_0 - \beta_0^+)\|_{w,p} < \infty, & \text{if } 1 \leq p < \infty, \\ & \|(\tilde{\alpha}_0, \tilde{\beta}_0)\|_{w,p} < \infty, \\ & \|(\alpha_0, \beta_0)\|_{w,\infty} < \infty, \quad \|(\alpha_0 - \alpha_0^+, \beta_0 - \beta_0^+)\|_{w^2,\infty} < \infty, & \text{if } p = \infty. \\ & \|(\tilde{\alpha}_0, \tilde{\beta}_0)\|_{w^2,\infty} < \infty, \end{aligned}$$

Let $(\alpha(t), \beta(t))$, $t \in (-T, T)$, be the unique solution of the equation $\text{AL}_{\underline{r}}(\alpha, \beta) = 0$ with summation coefficients $\{c_{j,\pm}\}_{j=0}^{r_{\pm}}$, corresponding to the initial conditions

$$(3.12) \quad \alpha(0) = \alpha_0 + \tilde{\alpha}_0, \quad \beta(0) = \beta_0 + \tilde{\beta}_0.$$

Then this solution is of the form

$$(3.13) \quad \alpha(t) = \alpha_0 + \tilde{\alpha}(t), \quad \beta(t) = \beta_0 + \tilde{\beta}(t),$$

where $\|(\tilde{\alpha}(t), \tilde{\beta}(t))\|_{w,p} < \infty$, respectively, $\|(\tilde{\alpha}(t), \tilde{\beta}(t))\|_{w^2,\infty} < \infty$.

Proof. The proof is similar to that of Theorem 2.4. From $\text{AL}_{\underline{r}}(\alpha, \beta) = 0$ we obtain an inhomogeneous differential equation for $(\tilde{\alpha}, \tilde{\beta})$. The homogeneous part is a finite sum over shifts of $(\tilde{\alpha}, \tilde{\beta})$. The inhomogeneous part consists of products of α_0 , β_0 and their shifts, whose $\|\cdot\|_{w,p}$ norm is finite by Hölder's inequality, and of sums of the form $c_{j,\pm}\alpha_0$, $c_{j,\pm}\beta_0$ and shifts thereof,

$$\begin{aligned} & -(c_{0,+}S^{+r_+} + c_{1,+}S^{+r_+-1} + \cdots + c_{r_+-1,+}S^{+1} + c_{\underline{r}} \\ & + c_{0,-}S^{-r_-} + c_{1,-}S^{-r_--1} + \cdots + c_{r_--1,-}S^{-1})\alpha_0 \end{aligned}$$

(and analogously for β_0), from which restriction (3.11) arises. Again $S^{\pm j}$ denote the shift operators $S^{\pm j}\alpha_0(n) = \alpha_0(n \pm j)$. The requirement $\|(\alpha_0 - \alpha_0^+, \beta_0 - \beta_0^+)\|_{w,p} < \infty$ yields the algebraic constraint (3.11) for $c_{j,\pm}$. Finally, note that it is no restriction to assume $c_{\underline{r}} = 0$ by (3.6). $\square$

For example, we obtain such decaying solutions for $\text{AL}_{(0,1)}(\alpha, \beta)$ if $c_{0,+} = 0$ and for $\text{AL}_{(1,1)}(\alpha, \beta)$ if $c_{0,+} = -c_{0,-}$ (or $c_{0,+} + c_{0,-} + c_{\underline{1}} = 0$ as in Theorem 2.4).

ACKNOWLEDGMENTS

The author thanks Armin Rainer and Gerald Teschl for valuable discussions on this topic.

REFERENCES

1. M. J. Ablowitz, G. Biondini, and B. Prinari, *Inverse scattering transform for the integrable discrete nonlinear Schrödinger equation with nonvanishing boundary conditions*, Inverse Problems **23** (2007), 1711–1758. MR2348731 (2008f:37154)
2. M. J. Ablowitz and P. A. Clarkson, *Solitons, Nonlinear Evolution Equations and Inverse Scattering*, Cambridge University Press, Cambridge, 1991. MR1149378 (93g:35108)
3. M. J. Ablowitz and J. F. Ladik, *Nonlinear differential-difference equations*, J. Math. Phys. **16** (1975), 598–603. MR0377223 (51:13396)
4. ———, *Nonlinear differential-difference equations and Fourier analysis*, J. Math. Phys. **17** (1976), 1011–1018. MR0427867 (55:897)
5. ———, *A nonlinear difference scheme and inverse scattering*, Studies Appl. Math. **55** (1976), 213–229. MR0471341 (57:11076)
6. ———, *On the solution of a class of nonlinear partial difference equations*, Studies Appl. Math. **57** (1977), 1–12. MR0492975 (58:12018)
7. M. J. Ablowitz, B. Prinari, and A. D. Trubatch, *Discrete and Continuous Nonlinear Schrödinger Systems*, London Mathematical Society Lecture Note Series, Vol. 302, Cambridge University Press, Cambridge, 2004. MR2040621 (2005c:37117)
8. R. Abraham, J.E. Marsden, and T. Ratiu, *Manifolds, Tensor Analysis, and Applications*, 2nd ed., Springer, New York, 1988. MR960687 (89f:58001)
9. I. N. Bondareva, *The Korteweg-de Vries equation in classes of increasing functions with prescribed asymptotic behaviour as $|x| \rightarrow \infty$* , Math. USSR Sb. **50** (1985), no. 1, 125–135. MR0717670 (85h:35180)
10. I. N. Bondareva and M. Shubin, *Increasing asymptotic solutions of the Korteweg-de Vries equation and its higher analogues*, Sov. Math. Dokl. **26** (1982), no. 3, 716–719. MR0685832 (84k:35121)
11. K. Deimling, *Ordinary Differential Equations on Banach Spaces*, Lecture Notes in Mathematics, 596, Springer, Berlin, 1977. MR0463601 (57:3546)
12. C. S. Gardner, J. M. Greene, M. D. Kruskal, and R. M. Miura, *A method for solving the Korteweg-de Vries equation*, Phys. Rev. Letters **19** (1967), 1095–1097.
13. F. Gesztesy, H. Holden, J. Michor, and G. Teschl, *Soliton Equations and Their Algebro-Geometric Solutions. Volume II: $(1+1)$ -Dimensional Discrete Models*, Cambridge Studies in Advanced Mathematics, 114, Cambridge University Press, Cambridge, 2008. MR2446594

14. ———, *The Ablowitz–Ladik hierarchy revisited*, in Methods of Spectral Analysis in Math. Physics, J. Janas et al. (eds.), 139–190, Oper. Theory Adv. Appl., 186, Birkhäuser, Basel, 2008.
15. ———, *Local conservation laws and the Hamiltonian formalism for the Ablowitz–Ladik hierarchy*, Stud. Appl. Math. **120** (2008), no. 4, 361–423. MR2416645 (2009i:37156)
16. H. Krüger and G. Teschl, *Unique continuation for discrete nonlinear wave equations*, arXiv:0904.0011.
17. J. Michor, *Inverse scattering transform for the Ablowitz–Ladik Hierarchy with quasi-periodic background*, in preparation.
18. G. Teschl, *On the spatial asymptotics of solutions of the Toda lattice*, Discrete Contin. Dyn. Sys. **273** (2010), no. 3, 1233–1239.
19. V. E. Vekslerchik and V. V. Konotop, *Discrete nonlinear Schrödinger equation under non-vanishing boundary conditions*, Inverse Problems **8** (1992), 889–909. MR1195946 (93j:35166)

FACULTY OF MATHEMATICS, UNIVERSITY OF VIENNA, NORDBERGSTRASSE 15, 1090 WIEN, AUSTRIA – AND – INTERNATIONAL ERWIN SCHRÖDINGER INSTITUTE FOR MATHEMATICAL PHYSICS, BOLTZMANNGASSE 9, 1090 WIEN, AUSTRIA

E-mail address: Johanna.Michor@univie.ac.at