SOME BASIC FACTS ON THE SYSTEM $\Delta u - W_u(u) = 0$

NICHOLAS D. ALIKAKOS

(Communicated by Yingfei Yi)

ABSTRACT. We rewrite the system $\Delta u - W_u(u) = 0$, for $u : \mathbb{R}^n \rightarrow \mathbb{R}^n$, in the form $\operatorname{div} T = 0$, where T is an appropriate stress-energy tensor, and derive certain *a priori* consequences on the solutions. In particular, we point out some differences between two paradigms: the phase-transition system, with target a finite set of points, and the Ginzburg–Landau system, with target a connected manifold.

1. INTRODUCTION

This paper is concerned with entire solutions to the system

$$(1.1) \quad \Delta u - W_u(u) = 0, \text{ for } u : \mathbb{R}^n \rightarrow \mathbb{R}^n,$$

where $W \in C^2(\mathbb{R}^n; \mathbb{R})$ and $W_u := (\partial W / \partial u_1, \dots, \partial W / \partial u_n)^\top$. Two distinguished examples are: (a) the phase-transition model or vector Allen–Cahn equation, where W has a finite number of global minima $a_1, a_2, \dots, a_N$ (see Baldo [4], Bronsard and Reitich [8]) and (b) the Ginzburg–Landau system $\Delta u - (|u|^2 - 1)u = 0$ (see Bethuel, Brezis, and Hélein [5]).

First, we introduce a stress-energy tensor T , that is, an $n \times n$ matrix $T = T(u, \nabla u)$, and show that (1.1) can be written in the form

$$(1.2) \quad \operatorname{div} T = 0,$$

for solutions $u \in W_{\operatorname{loc}}^{1,2} \cap L_{\operatorname{loc}}^\infty$. Next, by following the general procedure due to Schoen [26] for divergence-free tensors, we establish, under the hypothesis

$$(1.3) \quad W \geq 0,$$

the (weak) monotonicity formula

$$(1.4) \quad \frac{d}{dR} \left(\frac{1}{R^{n-2}} \int_{|x-x_0|<R} \left(\frac{1}{2} |\nabla u|^2 + W(u) \right) dx \right) \geq 0,$$

where ∇u is the matrix $(\partial u_i / \partial x_j)$, for $i, j = 1, \dots, n$, $|\cdot|$ is the Euclidean norm, $R > 0$, and x_0 is an arbitrary point in $\mathbb{R}^n$. Finally, as an application, we obtain a Liouville-type result: For a solution $u \in W_{\operatorname{loc}}^{1,2} \cap L_{\operatorname{loc}}^\infty$ to system (1.1) under the hypothesis (1.3) and for $n \geq 2$, we show that

$$(1.5) \quad \int_{\mathbb{R}^n} \left(\frac{1}{2} |\nabla u|^2 + W(u) \right) dx < \infty \quad \text{implies} \quad u \equiv \text{constant}.$$

Received by the editors September 23, 2009 and, in revised form, February 20, 2010.
 2000 *Mathematics Subject Classification*. Primary 35Jxx.

©2010 American Mathematical Society
 Reverts to public domain 28 years from publication

Formula (1.4) corresponds to the usual monotonicity formula for harmonic maps (see Schoen and Uhlenbeck [27]). The (strong) monotonicity formula

$$(1.6) \quad \frac{d}{dR} \left(\frac{1}{R^{n-1}} \int_{|x-x_0|<R} \left(\frac{1}{2} |\nabla u|^2 + W(u) \right) dx \right) \geq 0$$

implies (1.4). It can be motivated by the relationship between the solutions of the phase-transition model and minimal surfaces and thus by the monotonicity of the quantity

$$\frac{1}{R^{n-1}} \int_{|x-x_0|<R} |\nabla u| dx.$$

However, the validity of (1.6) depends on the estimate

$$(1.7) \quad |\nabla u|^2 \leq 2W(u).$$

This last estimate is a well-known result of Modica [20] for solutions to the scalar Allen–Cahn equation

$$(1.8) \quad \Delta u - W'(u) = 0, \text{ for } u : \mathbb{R}^n \rightarrow \mathbb{R},$$

but it is not known¹ for systems like (1.1). In the case of graphs $u : \mathbb{R}^n \rightarrow \mathbb{R}$, Modica [22] has also established (1.6). Caffarelli, Garofalo, and Segala [9] have extended both results of Modica to a wider class of scalar equations under the hypothesis that $W \geq 0$, with $W = 0$ on a finite set of points. Formula (1.4) generalizes analogous results of Bethuel, Brezis, and Orlandi [6], Rivière [24], and Farina [14] for the Ginzburg–Landau system.

Imanen [18] has introduced a predecessor of the stress-energy tensor in his work on the evolution scalar Allen–Cahn equation versus motion by mean curvature. In the context of the Ginzburg–Landau system, we note the book of Sandier and Serfaty [25], where this tensor has been utilized in a number of ways. Our source is Alikakos and Betelú [2], where the tensor was introduced specifically for writing the system as a divergence-free condition in order to derive the Plateau angle conditions in a transparent way.

In Section 2.4 we give a derivation of the well-known monotonicity formula for minimal surfaces (see Simon [28], Ecker [12]) under hypotheses of smoothness, based on the stress-energy tensor, by applying Schoen’s procedure [26].

2. THE WEAK MONOTONICITY FORMULA

2.1. The stress-energy tensor. We begin by introducing a *stress-energy tensor* for vector fields $u : \mathbb{R}^n \rightarrow \mathbb{R}^n$. Let

$$(2.1) \quad T := \frac{1}{2} \begin{pmatrix} |u_{x_1}|^2 - \sum_{i \neq 1}^n |u_{x_i}|^2 - 2W(u), & 2u_{x_1} \cdot u_{x_2}, & \dots, & 2u_{x_1} \cdot u_{x_n} \\ 2u_{x_2} \cdot u_{x_1}, & |u_{x_2}|^2 - \sum_{i \neq 2}^n |u_{x_i}|^2 - 2W(u), & \dots, & 2u_{x_2} \cdot u_{x_n} \\ & & \ddots & \\ 2u_{x_n} \cdot u_{x_1}, & 2u_{x_n} \cdot u_{x_2}, & \dots, & |u_{x_n}|^2 - \sum_{i \neq n}^n |u_{x_i}|^2 - 2W(u) \end{pmatrix}.$$

¹See Section 2.3 for useful information on this point, which we owe to Alberto Farina.

So, $T = (T_1, T_2, \dots, T_n)^\top$ and $\operatorname{div} T = (\operatorname{div} T_1, \operatorname{div} T_2, \dots, \operatorname{div} T_n)^\top$.

The relevance of T to the system (1.1) can be seen by calculating its divergence. We have

$$\begin{aligned} \operatorname{div} T_j &= \sum_{k \neq j} (u_{x_j} \cdot u_{x_k})_{,x_k} + \frac{1}{2} \left(|u_{x_j}|^2 - \sum_{i \neq j} |u_{x_i}|^2 - 2W(u) \right)_{,x_j} \\ &= \sum_{k \neq j} (u_{x_j x_k} \cdot u_{x_k} + u_{x_j} \cdot u_{x_k x_k}) + u_{x_j} \cdot u_{x_j x_j} - \sum_{i \neq j} u_{x_i} \cdot u_{x_i x_j} - W_u \cdot u_{x_j} \\ &= u_{x_j} \cdot (\Delta u - W_u(u)); \end{aligned}$$

thus,

$$(2.2) \quad \operatorname{div} T = \begin{pmatrix} u_{x_1} \cdot (\Delta u - W_u(u)) \\ u_{x_2} \cdot (\Delta u - W_u(u)) \\ \vdots \\ u_{x_n} \cdot (\Delta u - W_u(u)) \end{pmatrix} = (\nabla u)^\top (\Delta u - W_u(u)).$$

Therefore, in the class of smooth vector fields, equation (1.1) is almost equivalent to T being divergence-free since

$$(2.3) \quad \begin{cases} \Delta u - W_u(u) = 0 & \Rightarrow \operatorname{div} T = 0, \\ \Delta u - W_u(u) = 0 & \Leftarrow \operatorname{div} T = 0, \det \nabla u \neq 0. \end{cases}$$

Next, we calculate the trace of T :

$$\begin{aligned} \operatorname{tr} T &= \sum_{i=1}^n T_{ii} = \frac{1}{2} \left(\sum_{i=1}^n |u_{x_i}|^2 - 2nW(u) - \sum_{i=1}^n \left(\sum_{k \neq i} |u_{x_k}|^2 \right) \right) \\ &= \frac{1}{2} \left(\sum_{i=1}^n |u_{x_i}|^2 - 2nW(u) - \sum_{i=1}^n \left(\sum_{k=1}^n |u_{x_k}|^2 - |u_{x_i}|^2 \right) \right) \\ (2.4) \quad &= \frac{2-n}{2} \sum_{i=1}^n |u_{x_i}|^2 - nW(u). \end{aligned}$$

Finally, we introduce the *interface-energy density*

$$(2.5) \quad e(u) := \frac{1}{2} |\nabla u|^2 + W(u),$$

where ∇u is the matrix $(\partial u_i / \partial x_j)$, for $i, j = 1, \dots, n$ and $|\cdot|$ is the Euclidean norm.

Notice that

$$(2.6) \quad \operatorname{tr} T = -n e(u) + |\nabla u|^2,$$

$$(2.7) \quad \operatorname{tr} T \leq -(n-2) e(u).$$

2.2. The monotonicity formula.

Theorem 2.1. *Let u be a $W_{\text{loc}}^{1,2}(\mathbb{R}^n; \mathbb{R}^n) \cap L_{\text{loc}}^\infty(\mathbb{R}^n; \mathbb{R}^n)$ solution to the system*

$$\Delta u - W_u(u) = 0, \text{ for } u : \mathbb{R}^n \rightarrow \mathbb{R}^n,$$

under the hypothesis that W is $C^2(\mathbb{R}^n; \mathbb{R})$ and $W \geq 0$. Then, we have

$$(2.8) \quad \frac{d}{dR} \left(R^{-(n-2)} E_{B_R}(u) \right) \geq 0, \text{ for } R > 0,$$

where

$$(2.9) \quad E_{B_R}(u) = \int_{B_R} \left(\frac{1}{2} |\nabla u|^2 + W(u) \right) dx,$$

with $x_0 \in \mathbb{R}^n$ arbitrary and $B_R := B(x_0; R)$ the ball in $\mathbb{R}^n$.

Proof. We begin by noting the simple fact that

$$(2.10) \quad T + e(u) \text{Id} = (\nabla u)^\top (\nabla u) \geq 0,$$

where Id stands for the identity matrix on $\mathbb{R}^n$.

Take now $x_0 = 0$. Following Schoen [26], we have

$$\begin{aligned} \sum_{i,j} \int_{B_R} (x^i T_{ij})_{,j} &= \sum_{i,j} \int_{B_R} (\delta_{ij} T_{ij} + x^i T_{ij,j}) \\ &= \sum_i \int_{B_R} T_{ii} \quad (\text{from (2.3)}) \\ (2.11) \quad &\leq -(n-2) \int_{B_R} e(u) \quad (\text{from (2.7)}). \end{aligned}$$

On the other hand, by the divergence theorem and for $\nu = x/R$,

$$\begin{aligned} \sum_{i,j} \int_{B_R} (x^i T_{ij})_{,j} &= \sum_{i,j} \int_{\partial B_R} x^i T_{ij} \nu_j \\ &= R \sum_{i,j} \int_{\partial B_R} T_{ij} \nu_i \nu_j \\ &= R \int_{\partial B_R} \langle T\nu, \nu \rangle \\ &\geq -R \int_{\partial B_R} e(u) \quad (\text{from (2.10)}) \\ (2.12) \quad &= -R \frac{dE_{B_R}(u)}{dR}, \end{aligned}$$

where $\langle \cdot, \cdot \rangle$ is the Euclidean inner product.

Combining (2.11) and (2.12) we obtain

$$-(n-2) E_{B_R}(u) \geq -R \frac{dE_{B_R}(u)}{dR}$$

or, equivalently,

$$(2.13) \quad \frac{d}{dR} \left(R^{-(n-2)} E_{B_R}(u) \right) \geq 0.$$

By utilizing the translation invariance of (1.1), we conclude that x_0 can be arbitrary in (2.13) and the proof of the theorem is complete. $\square$

2.3. Remarks. Note that if the analog of the Modica gradient estimate

$$(2.14) \quad |\nabla u|^2 \leq 2W(u)$$

is assumed to hold for system (1.1), then (2.7) can be strengthened to

$$(2.15) \quad \text{tr } T \leq -(n-1) e(u),$$

from which the strong monotonicity formula

$$(2.16) \quad \frac{d}{dR} \left(R^{-(n-1)} E_{B_R}(u) \right) \geq 0, \text{ for } R > 0,$$

follows (by using in (2.11) formula (2.15) instead of (2.7)). Formula (2.16) has been derived already by Modica [20] (see also Caffarelli, Garofalo, and Segala [9]) for entire solutions to the scalar Allen–Cahn equation

$$(2.17) \quad \Delta u - W'(u) = 0, \text{ for } u : \mathbb{R}^n \rightarrow \mathbb{R}, \quad W : \mathbb{R} \rightarrow \mathbb{R}.$$

Theorem 2.1 is more general from the point of view of W . The estimate (2.14) does not hold in such generality. Actually neither (2.14) nor (2.16) hold for the Ginzburg–Landau system [14].

The stress-energy tensor T was introduced in joint unpublished work with Betelú [2] (see also the M.Sc. thesis of Dimitroula [11] for further elaboration), where it is utilized for deriving the Plateau angle conditions and was motivated by the work of Bronsard and Reitich [8], where the Plateau conditions were derived in two dimensions. See also the related work of Sandier and Serfaty [25] on the Ginzburg–Landau system. The predecessor of T for scalar fields $u : \mathbb{R}^n \rightarrow \mathbb{R}$ was introduced before by Ilmanen in [18].

The physical meaning of T can be seen via the scaled Allen–Cahn system

$$(2.18) \quad \varepsilon \Delta u - \frac{1}{\varepsilon} W_u(u) = 0, \text{ for } u : \mathbb{R}^n \rightarrow \mathbb{R}^n.$$

This requires a digression which we now take. As is well-known from the theory of Γ -convergence (see De Giorgi and Franzoni [10], Modica and Mortola [23], Modica [21], Sternberg [29], Baldo [4], and for expository work Alberti [1] and Braides [7]),

$$(2.19) \quad \Gamma\text{-}\lim_{\varepsilon \rightarrow 0} \int_{\Omega} \left(\frac{\varepsilon}{2} |\nabla u|^2 + \frac{1}{\varepsilon} W(u) \right) dx = F(u), \text{ for } \Omega \text{ bounded, open in } \mathbb{R}^n,$$

with

$$(2.20) \quad F(u) = \sum_{i < j} \sigma_{ij} \mathcal{H}^{n-1}(S_{ij}u),$$

where $S_{ij}u$ is the interface in Ω between the phases $\{u = a_i\}$ and $\{u = a_j\}$, $\mathcal{H}^{n-1}$ is the $(n-1)$ -dimensional Hausdorff measure, and σ_{ij} is the corresponding surface-tension coefficient (there holds, see below, $\sigma_{ij} + \sigma_{jk} \geq \sigma_{ik}$). For the vector case see Baldo [4], where $W \geq 0$ and vanishes precisely at $a_1, \dots, a_N$.

Without a constraint, the global minimizers u^ε are trivial. Γ -convergence methods treat the integral constraint (see [4]) and lead to surfaces of constant mean curvature. Local minimizers u^ε have been treated in Kohn and Sternberg [19] and, from a point of view closer to the present article, in Hutchinson and Tonegawa [17]. The blow-down limit in (2.19) relates global minimizers u^ε of

$$\int_{\Omega} \left(\frac{\varepsilon}{2} |\nabla u|^2 + \frac{1}{\varepsilon} W(u) \right) dx$$

to global minimizers of the perimeter functional (2.20), that is, to minimal surfaces S_{ij} . As is well-known, for minimal surfaces the following monotonicity formula holds (see Simon [28], Ecker [12]):

$$(2.21) \quad \frac{d}{dR} \left(R^{-(n-1)} \mathcal{H}^{n-1}(S_{ij} \cap B^n(p; R)) \right) \geq 0, \text{ for } 0 \leq R < \delta,$$

where $B^n(p; R)$ is any ball in $\mathbb{R}^n$ with center p , δ -away from the boundary $\partial\Omega$. Notice the relationship between (2.16) and (2.21).

2.4. The limiting stress-energy tensor and the monotonicity formula for minimal surfaces. After this background, we are ready to derive formally the limiting stress-energy tensor, which has a very clear physical meaning. Let U_{ij} be the connection between a_i and a_j (see Sternberg [30], Alikakos and Fusco [3]); that is, let

$$(2.22) \quad \begin{cases} \ddot{U}_{ij}(\eta) - W_u(U_{ij}(\eta)) = 0, \text{ where } U_{ij} : \mathbb{R} \rightarrow \mathbb{R}^n, \dot{U} := dU/d\eta, \\ U_{ij}(-\infty) = a_i, U_{ij}(+\infty) = a_j, \text{ for } i \neq j, U_{ij}(0) = 0. \end{cases}$$

It is well-known that U_{ij} satisfies the equipartition relation

$$\frac{1}{2}|\dot{U}_{ij}|^2 = W(U_{ij})$$

and also

$$\sigma_{ij} = \int_{\mathbb{R}} \left(\frac{1}{2}|\dot{U}_{ij}|^2 + W(U_{ij}) \right) d\eta.$$

For small $\varepsilon > 0$, the minimizer u^ε can be approximated by

$$u^\varepsilon \approx U\left(\frac{d(x, S)}{\varepsilon}\right),$$

where $d(\cdot, S)$ is the signed distance from the $(n-1)$ -manifold S , $S = S_{ij}$ is the part of the interface $I = \bigcup_{i < j} S_{ij}$ which realizes the distance of x from I , and $U = U_{ij}$ is the corresponding connection. We note that this approximation is appropriate away from the intersections of S_{ij} 's (for example, away from triple junctions in the plane). However, the contribution to the energy of the intersection set is proportional to the size of small neighborhoods surrounding the set, and so can be ignored. (This is supported by formal results in Bronsard and Reitich [8] and rigorous work in Baldo [4], plus a lot of accumulated evidence on diffuse-interface models.)

The scaled stress-energy tensor is given by

$$(2.23) \quad T_\varepsilon := \frac{1}{2} \begin{pmatrix} \varepsilon|u_{x_1}|^2 - \varepsilon \sum_{i \neq 1}^n |u_{x_i}|^2 - 2\frac{W(u)}{\varepsilon}, & 2\varepsilon u_{x_1} \cdot u_{x_2}, & \dots, & 2\varepsilon u_{x_1} \cdot u_{x_n} \\ 2\varepsilon u_{x_2} \cdot u_{x_1}, & \varepsilon|u_{x_2}|^2 - \varepsilon \sum_{i \neq 2}^n |u_{x_i}|^2 - 2\frac{W(u)}{\varepsilon}, & \dots, & 2\varepsilon u_{x_2} \cdot u_{x_n} \\ & & \ddots & \\ 2\varepsilon u_{x_n} \cdot u_{x_1}, & 2\varepsilon u_{x_n} \cdot u_{x_2}, & \dots, & \varepsilon|u_{x_n}|^2 - \varepsilon \sum_{i \neq n}^n |u_{x_i}|^2 - 2\frac{W(u)}{\varepsilon} \end{pmatrix}$$

and corresponds to (2.18). By evaluating T_ε on u^ε , by making use of the approximate relationship above, and by utilizing the canonical coordinates around S (see [16, Appendix]), we obtain

$$(2.24) \quad \lim_{\varepsilon \rightarrow 0} T_\varepsilon(u^\varepsilon) = T_0 := \sigma \mathcal{H}^{n-1} \llcorner S \begin{pmatrix} d_{x_1}^2 - 1 & d_{x_1} d_{x_2} & \dots & d_{x_1} d_{x_n} \\ d_{x_2} d_{x_1} & d_{x_2}^2 - 1 & \dots & d_{x_2} d_{x_n} \\ & & \ddots & \\ d_{x_n} d_{x_1} & d_{x_n} d_{x_2} & \dots & d_{x_n}^2 - 1 \end{pmatrix},$$

in the sense of Radon measures. For example, for the first entry in the matrix above, we have

$$\lim_{\varepsilon \rightarrow 0} \int_{\mathbb{R}^n} \left(\varepsilon |u_{x_1}|^2 - \varepsilon \sum_{i \neq 1} |u_{x_i}|^2 - 2 \frac{W(u^\varepsilon)}{\varepsilon} \right) \phi(x) dx = 2\sigma(d_{x_1}^2 - 1) \int_S \phi(y) dS_y,$$

for every $\phi \in C_c(\mathbb{R}^n; \mathbb{R})$. Thus,

$$(2.25) \quad T_0|_S = \sigma(\nabla d \otimes \nabla d - \text{Id});$$

that is, T_0 is an orthogonal projection in $\mathbb{R}^n$, on the tangent space of S . Therefore, T_0 is a symmetric matrix generating tangential forces and, as such, it is properly called a stress-energy tensor since the surface tension force is supposed to be tangential to the interface (see also Ilmanen [18]).

We write (2.25) by dropping the subscript for simplicity,

$$T|_S = \sigma(\nabla d \otimes \nabla d - \text{Id}), \quad T_{ij}|_S = \sigma(d_{x_i} d_{x_j} - \delta_{ij}).$$

Next we derive (2.21) under the hypothesis of smoothness.

A) Since $T = -\sigma \text{proj}_{T_x S}$, we have that

$$\text{div}_S T = -\sigma \mathbf{H},$$

where $\mathbf{H} := H\nu$ is the mean-curvature vector, with ν a unit normal. Thus, $\text{div}_S T = 0$ is equivalent to $H = 0$.

B) Utilizing that T is a projection gives

$$\text{div}_S T(X) = -\sigma \mathbf{H} \cdot X - \sigma \text{div}_S X,$$

for any vector field X in $\mathbb{R}^n$. Using that $H = 0$, we have

$$\int_{S \cap B_R} \text{div}_S T(X) = -\sigma \int_{S \cap B_R} \text{div}_S X,$$

and by selecting $X(x) = x - p$ and $B_R = B_R(p)$, for $p \in S$, we obtain

$$-\sigma \int_{S \cap B_R} \text{div}_S X = -\sigma(n-1) \mathcal{H}^{n-1}(S \cap B_R).$$

C) By the divergence theorem on surfaces (see Simon [28], Ecker [12, Appendix]),

$$\int_{S \cap B_R} \text{div}_S X = - \int_{S \cap B_R} H \cdot X + \int_{S \cap \partial B_R} X \cdot \gamma,$$

where γ is a unit co-normal. Hence,

$$\begin{aligned} \int_{S \cap B_R} \text{div}_S T(X) &= -\sigma \int_{S \cap B_R} \text{div}_S X \\ &= -\sigma \int_{S \cap \partial B_R} (x - p) \cdot \gamma \\ &\geq -\sigma R \mathcal{H}^{n-2}(S \cap \partial B_R) \\ &= -\sigma R \frac{d}{dR} \left(\mathcal{H}^{n-1}(S \cap B_R) \right). \end{aligned}$$

D) Combining B and C we obtain

$$-\sigma(n-1) \mathcal{H}^{n-1}(S \cap B_R) \geq -\sigma R \frac{d}{dR} \left(\mathcal{H}^{n-1}(S \cap B_R) \right)$$

or, equivalently,

$$\frac{d}{dR} \left(R^{-(n-1)} \mathcal{H}^{n-1}(S \cap B_R) \right) \geq 0.$$

The above gives an answer, under hypotheses of smoothness, to a question raised in Schoen [26] on the possibility of such an approach for minimal surfaces, that is, the existence of a physically meaningful stress-energy tensor. It is this question that motivated the present work.

The application of the divergence theorem on a vector field tangent to the surface as a means for deriving the monotonicity formula is well-known (see Simon [28, p. 83]). The point in Schoen [26] is that the stress-energy tensor, a physically motivated quantity, treats in a unified way a variety of setups. The calculation above can be extended to varifolds as in Simon [28, p. 235].

3. A LIOUVILLE THEOREM

Theorem 3.1. *Let u be a $W_{\text{loc}}^{1,2}(\mathbb{R}^n; \mathbb{R}^n) \cap L_{\text{loc}}^\infty(\mathbb{R}^n; \mathbb{R}^n)$ solution to the system*

$$\Delta u - W_u(u) = 0, \text{ for } u : \mathbb{R}^n \rightarrow \mathbb{R}^n,$$

under the hypothesis that W is $C^2(\mathbb{R}^n; \mathbb{R})$ and $W \geq 0$. Then, for $n \geq 2$ we have that

$$(3.1) \quad \int_{\mathbb{R}^n} \left(\frac{1}{2} |\nabla u|^2 + W(u) \right) dx < \infty \quad \text{implies} \quad u \equiv \text{constant}.$$

Proof. For $n \geq 3$, by (2.8), for $\sigma \geq R$,

$$\begin{aligned} \frac{1}{R^{n-2}} \int_{|x| < R} \left(\frac{1}{2} |\nabla u|^2 + W(u) \right) dx &\leq \frac{1}{\sigma^{n-2}} \int_{|x| < \sigma} \left(\frac{1}{2} |\nabla u|^2 + W(u) \right) dx \\ &\leq \frac{1}{\sigma^{n-2}} \int_{\mathbb{R}^n} \left(\frac{1}{2} |\nabla u|^2 + W(u) \right) dx. \end{aligned}$$

By taking the limit $\sigma \rightarrow \infty$ we are set.

For $n = 2$, (2.4) gives

$$\text{tr } T = -2W(u).$$

Following the proof of Theorem 2.1, we obtain

$$R \frac{dE_{B_R}(u)}{dR} \geq 2 \int_{B_R} W(u) dx \geq 2 \int_{B_{R_0}} W(u) dx,$$

for $R \geq R_0$. Hence, integrating we obtain

$$E_{B_R}(u) \geq E_{B_{R_0}}(u) + 2 \log \frac{R}{R_0} \int_{B_{R_0}} W(u) dx.$$

Thus, the hypothesis implies that $W(u) \equiv 0$. But then the components u_1, u_2 of u are harmonic, with

$$\int_{\mathbb{R}^2} |\nabla u_i|^2 dx < \infty, \text{ for } i = 1, 2.$$

Therefore, $u_i \equiv \text{constant}$ by the mean-value property of harmonic functions applied to $\partial u_i / \partial x_j$. The proof of the theorem is complete. $\square$

Note that an examination of the arguments above renders the estimates

$$\begin{cases} E_{B_R}(u) = o(R^{n-2}), \text{ as } R \rightarrow \infty, n \geq 3 & \text{implies } u \equiv \text{constant}, \\ E_{B_R}(u) = o(\log R), \text{ as } R \rightarrow \infty, n = 2 & \text{implies } u \equiv \text{constant}. \end{cases}$$

For related work on scalar equations see Modica [22] and Caffarelli, Garofalo, and Segala [9]. For Liouville theorems for the related Ginzburg–Landau systems see Farina [13], [14].

Note added after proof.

- (a) The stress-energy tensor is well defined for $u: \mathbb{R}^n \rightarrow \mathbb{R}^k$, and the statement and proof of Theorem 2.1 are valid in this more general setup.
- (b) The identities in the recent work of C. Gui, “Hamiltonian identities for elliptic partial differential equations”, *Journal of Functional Analysis*, 2008, 254:904–933, follow from the divergence free form of the system.

ACKNOWLEDGMENTS

This work was carried out during a visit to Stanford University in the spring of 2009. Thanks are due to Rafe Mazzeo and Rick Schoen for several stimulating conversations and to the Department of Mathematics of Stanford University as a whole for the great hospitality. Thanks also go to Peter Bates, Alex Freire, and Panagiotis Smyrnelis for their comments on a previous version of the paper.

REFERENCES

- [1] G. Alberti. Variational methods for phase transitions, an approach via Γ -convergence. In L. Ambrosio and N. Dancer, *Calculus of variations and partial differential equations*, edited by G. Buttazzo, A. Marino, M. K. V. Murthy. Springer, Berlin, 2000, pp. 95–114. MR1757697
- [2] N. D. Alikakos and S. I. Betelú. Plateau conditions for the vector Allen–Cahn and Cahn–Hilliard equations. Unpublished preprint, 2005.
- [3] N. D. Alikakos and G. Fusco. On the connection problem for potentials with several global minima. *Ind. Univ. Math. J.* **57** No. 4 (2008), pp. 1871–1906. MR2440884 (2009h:37124)
- [4] S. Baldo. Minimal interface criterion for phase transitions in mixtures of Cahn–Hilliard fluids. *Ann. Inst. Henri Poincaré, Anal. Non Linéaire* **7** No. 2 (1990), pp. 67–90. MR1051228 (91j:76005)
- [5] F. Bethuel, H. Brezis, and F. Hélein. *Ginzburg–Landau vortices*. Birkhäuser, Boston, MA, 1994. MR1269538 (95c:58044)
- [6] F. Bethuel, H. Brezis, and G. Orlandi. Asymptotics of the Ginzburg–Landau equation in arbitrary dimensions. *J. Funct. Anal.* **186** No. 2 (2001), pp. 432–520. MR1864830 (2002k:35081a)
- [7] A. Braides. A handbook of Γ -convergence. In *Handbook of differential equations, Stationary partial differential equations*, Vol. 3, edited by M. Chipot and P. Quittner. North-Holland, Amsterdam, 2006, pp. 101–213.
- [8] L. Bronsard and F. Reitich. On three-phase boundary motion and the singular limit of a vector-valued Ginzburg–Landau equation. *Arch. Rat. Mech. Anal.* **124** No. 4 (1993), pp. 355–379. MR1240580 (94h:35122)
- [9] L. Caffarelli, N. Garofalo, and F. Segala. A gradient bound for entire solutions of quasi-linear equations and its consequences. *Comm. Pure Appl. Math.* **47** No. 11 (1994), pp. 1457–1473. MR1296785 (95k:35030)
- [10] E. De Giorgi and T. Franzoni. Su un tipo di convergenza variazionale. *Atti Accad. Naz. Lincei Rend. Cl. Sci. Fis. Mat. Natur.* **58** No. 8 (1975), pp. 842–850. MR0448194 (56:6503)
- [11] S. Dimitroula. *The vector quasi-static Stefan problem*. M.Sc. thesis, University of Athens, 2008.
- [12] K. Ecker. *Regularity theory for mean curvature flow*. Birkhäuser Verlag, Basel, 2004. MR2024995 (2005b:53108)

- [13] A. Farina. Finite-energy solutions, quantization effects and Liouville-type results for a variant of the Ginzburg–Landau systems in $\mathbb{R}^K$. *C. R. Acad. Sci. Paris, Sér. I, Math.* **325** No. 5 (1997), pp. 487–491. MR1692312 (2000d:35213)
- [14] A. Farina. Two results on entire solutions of Ginzburg–Landau system in higher dimensions. *J. Funct. Anal.* **214** No. 2 (2004), pp. 386–395. MR2083306 (2005g:35093)
- [15] A. Farina. Private communication.
- [16] D. Gilbarg and N. S. Trudinger. *Elliptic partial differential equations of second order*. Springer-Verlag, Berlin, second edition, 1983. MR737190 (86c:35035)
- [17] J. E. Hutchinson and Y. Tonegawa. Convergence of phase interfaces in the van der Waals–Cahn–Hilliard theory. *Calc. Var.* **10** No. 1 (2000), pp. 49–84. MR1803974 (2001m:49070)
- [18] T. Ilmanen. Convergence of the Allen–Cahn equation to Brakke’s motion by mean curvature. *J. Diff. Geom.* **38** No. 2 (1993), pp. 417–461. MR1237490 (94h:58051)
- [19] R. V. Kohn and P. Sternberg. Local minimizers and singular perturbations. *Proc. R. Soc. Edinburgh* **111A** No. 1–2 (1989), pp. 69–84. MR985990 (90c:49021)
- [20] L. Modica. A gradient bound and a Liouville theorem for nonlinear Poisson equations. *Comm. Pure Appl. Math.* **38** No. 5 (1985), pp. 679–684. MR803255 (87m:35088)
- [21] L. Modica. The gradient theory of phase transitions and the minimal interface criterion. *Arch. Rat. Mech. Anal.* **98** No. 1 (1987), pp. 123–142. MR866718 (88f:76038)
- [22] L. Modica. Monotonicity of the energy for entire solutions of semilinear elliptic equations. In *Partial differential equations and the calculus of variations*, Essays in honor of Ennio De Giorgi, Vol. 2, edited by F. Colombini, A. Marino, and L. Modica. Birkhäuser, Boston, MA, 1989, pp. 843–850. MR1034031 (91a:35015)
- [23] L. Modica and S. Mortola. Un esempio di Γ -convergenza. *Boll. Un. Mat. Ital.* **14B** No. 5 (1977), pp. 285–299. MR0445362 (56:3704)
- [24] T. Rivière. Line vortices in the $U(1)$ -Higgs model. *ESAIM Control Optim. Calc. Var.* **1** (1996), pp. 77–167. MR1394302 (97g:58043)
- [25] E. Sandier and S. Serfaty. *Vortices in the magnetic Ginzburg–Landau model*. Birkhäuser, Basel, 1997. MR2279839 (2008g:82149)
- [26] R. Schoen. *Lecture notes on general relativity*. Stanford University, 2009.
- [27] R. Schoen and K. Uhlenbeck. A regularity theory for harmonic maps. *J. Diff. Geom.* **17** No. 2 (1982), pp. 307–335. MR664498 (84b:58037a)
- [28] L. Simon. *Lectures on geometric measure theory*. Proceedings of the Centre for Mathematics and its Applications, Vol. 3, Australian National University, Canberra, 1984. MR756417 (87a:49001)
- [29] P. Sternberg. The effect of a singular perturbation of nonconvex variational problems. *Arch. Rat. Mech. Anal.* **101** No. 1 (1988), pp. 209–260. MR930124 (89h:49007)
- [30] P. Sternberg. Vector-valued local minimizers of nonconvex variational problems. *Rocky Mt. J. Math.* **21** No. 2 (1991), pp. 799–807. MR1121542 (92e:49016)

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ATHENS, PANEPISTEMIOPOLIS, 15784 ATHENS, GREECE

E-mail address: nalikako@math.uoa.gr