RANK-ONE FLOWS OF TRANSFORMATIONS WITH INFINITE ERGODIC INDEX

ALEXANDRE I. DANILENKO AND KYEWON K. PARK

(Communicated by Bryna Kra)

ABSTRACT. A rank-one infinite measure preserving flow $T = (T_t)_{t \in \mathbb{R}}$ is constructed such that for each $t \neq 0$, the Cartesian powers of the transformation T_t are all ergodic.

0. INTRODUCTION

In 1963 Kakutani and Parry discovered an interesting phenomenon in the theory of infinite measure preserving maps. They showed that for each $p > 0$ there exists a transformation whose p -th Cartesian power is ergodic but whose $(p+1)$ -th Cartesian power is not [KP]. Since then a number of other examples of transformations with exotic (from the point of view of the classical “probability preserving” ergodic theory) weak mixing properties were constructed. See surveys [Da2] and [DaS3] for a detailed discussion on that. In [Da1], [DaS1] these examples were extended to infinite measure preserving actions of discrete countable Abelian groups. Weak mixing properties of infinite measure preserving actions of *continuous* Abelian groups such as $\mathbb{R}$ and $\mathbb{R}^d$ were under consideration in [I–W]. In particular, a rank-one flow (i.e. $\mathbb{R}$ -action) whose Cartesian square is ergodic was constructed there. A rank-one infinite measure preserving flow $T = (T_t)_{t \in \mathbb{R}}$ with infinite ergodic index (i.e. the Cartesian powers of T are all ergodic) appeared in a recent paper [DaSo]. It can be deduced easily from [DaSo] that there is a residual subset D_T of $\mathbb{R}$ such that for each $t \in D_T$, the transformation T_t has infinite ergodic index. However the following more subtle question by C. Silva remains open so far:

Is there a rank-one infinite measure preserving flow T with $D_T = \mathbb{R} \setminus \{0\}$?

Our purpose in this paper is to answer his question in the affirmative.

Theorem 0.1. *There is a rank-one infinite σ -finite measure preserving flow $T = (T_t)_{t \in \mathbb{R}}$ such that for each $t \neq 0$, the transformation T_t has infinite ergodic index.*

The main idea of the proof is different from those that were used in [I–W] and [DaSo]. It is based on a technique of *forcing* a dynamical property. Originating from [Ry1], such techniques were utilized in [Ry2], [DaR], etc., to obtain mixing,

Received by the editors October 14, 2009 and, in revised form, February 23, 2010.

2010 *Mathematics Subject Classification.* Primary 37A40; Secondary 37A10, 37A25.

Key words and phrases. Infinite ergodic index, (C, F) -construction.

The first author thanks Ajou University and KIAS for supporting in part his visit to South Korea.

The second author is supported in part by KRF 2007-313-C00044.

power weak mixing, etc., of some systems. In this paper the desired flow appears as a certain limit of a sequence of weakly mixing finite measure preserving flows. We construct this sequence in such a way as to retain the property of infinite ergodic index in the limit. The construction is implemented in the language of (C, F) -actions (see [Da2]).

1. PRELIMINARIES: RANK-ONE ACTIONS AND (C, F) -ACTIONS OF $\mathbb{R}^d$

We first recall the definition of rank one. Let $S = (S_g)_{g \in \mathbb{R}^d}$ be a measure preserving action of $\mathbb{R}^d$ on a standard σ -finite measure space $(Y, \mathfrak{C}, \nu)$.

Definition 1.1. (i) A *Rokhlin tower or column* for S is a triple (A, f, F) , where $A \in \mathfrak{C}$, F is a cube in $\mathbb{R}^d$ and $f : A \rightarrow F$ is a measurable mapping such that for any Borel subset $H \subset F$ and an element $g \in \mathbb{R}^d$ with $g + H \subset F$, one has $f^{-1}(g + H) = S_g f^{-1}(H)$.
(ii) S is said to be of *rank one (by cubes)* if there exists a sequence of Rokhlin towers (A_n, f_n, F_n) such that the volume of F_n goes to infinity and for any subset $C \in \mathfrak{C}$ of finite measure, there is a sequence of Borel subsets $H_n \subset F_n$ such that

$$\lim_{n \rightarrow \infty} \nu(C \triangle f_n^{-1}(H_n)) = 0.$$

The (C, F) -construction of measure preserving actions for discrete countable groups was introduced in [dJ] and [Da1]. It was extended to the case of locally compact second countable Abelian groups in [DaS2]. (See also [Da2].) Here we outline it briefly for $\mathbb{R}^d$, $d \in \mathbb{N}$.

Given two subsets $E, F \subset \mathbb{R}^d$, by $E + F$ we mean their algebraic sum, i.e. $E + F = \{e + f \mid e \in E, f \in F\}$. The algebraic difference $E - F$ is defined in a similar way. If F is a singleton, say $F = \{f\}$, then we will write $E + f$ for $E + F$. Two subsets E and F of $\mathbb{R}^d$ are called *independent* if $(E - E) \cap (F - F) = \{0\}$; i.e. if $e + f = e' + f'$ for some $e, e' \in E, f, f' \in F$, then $e = e'$ and $f = f'$.

Fix $p \in \mathbb{N}$ and consider two sequences $(F_n)_{n=0}^\infty$ and $(C_n)_{n=1}^\infty$ of subsets in $\mathbb{R}^d$ such that F_n is a cube $[0, h_n) \times \cdots \times [0, h_n)$ (d times) for an $h_n \in \mathbb{R}$, $C_n \subset \mathbb{R}^d$ is a finite set, $\#C_n > 1$,

(1-1) F_n and C_{n+1} are independent, and

(1-2) $F_n + C_{n+1} \subset F_{n+1}$.

This means that $F_n + C_{n+1}$ consists of $\#C_{n+1}$ mutually disjoint ‘copies’ $F_n + c$ of F_n , $c \in C_{n+1}$, and all these copies are contained in F_{n+1} . We equip F_n with the measure $(\#C_1 \cdots \#C_n)^{-1}(\lambda_{\mathbb{R}^d} \upharpoonright F_n)$, where $\lambda_{\mathbb{R}^d}$ denotes Lebesgue measure on $\mathbb{R}^d$. Endow C_n with the equidistributed probability measure. Let $X_n := F_n \times \prod_{k>n} C_k$ stand for the product of measure spaces. Define an embedding $X_n \rightarrow X_{n+1}$ by setting

$$(f_n, c_{n+1}, c_{n+2}, \dots) \mapsto (f_n + c_{n+1}, c_{n+2}, \dots).$$

It is easy to see that this embedding is measure preserving. Then $X_0 \subset X_1 \subset \cdots$. Let $X := \bigcup_{n=0}^\infty X_n$ denote the inductive limit of the sequence of measure spaces X_n and let μ denote the corresponding measure on X . Then μ is σ -finite. It is infinite if and only if

$$(1-3) \quad \lim_{n \rightarrow \infty} \frac{h_n^d}{\#C_1 \cdots \#C_n} = \infty.$$

Given $g \in \mathbb{R}^d$ and $n \in \mathbb{N}$, we set

$$L_g^{(n)} := (F_n \cap (F_n - g)) \times \prod_{k>n} C_k \quad \text{and} \quad R_g^{(n)} := (F_n \cap (F_n + g)) \times \prod_{k>n} C_k.$$

Clearly, $L_g^{(n)} \subset L_g^{(n+1)}$ and $R_g^{(n)} \subset R_g^{(n+1)}$. Define a map $T_g^{(n)}: L_g^{(n)} \rightarrow R_g^{(n)}$ by setting

$$T_g^{(n)}(f_n, c_{n+1}, \dots) := (f_n + g, c_{n+1}, \dots).$$

Put

$$L_g := \bigcup_{n=1}^{\infty} L_g^{(n)} \subset X \quad \text{and} \quad R_g := \bigcup_{n=1}^{\infty} R_g^{(n)} \subset X.$$

Then a Borel one-to-one map $T_g: L_g \rightarrow R_g$ is well defined by $T_g \upharpoonright L_g^{(n)} = T_g^{(n)}$. Since $h_n \rightarrow \infty$, it follows that $\mu(X \setminus L_g) = \mu(X \setminus R_g) = 0$ for each $g \in \mathbb{R}^d$. It is easy to verify that $T := (T_g)_{g \in \mathbb{R}^d}$ is a Borel μ -preserving action of $\mathbb{R}^d$.

Definition 1.2. T is called the (C, F) -action of $\mathbb{R}^d$ associated with $(C_{n+1}, F_n)_{n \geq 0}$.

Each (C, F) -action is of rank one.

Given a Borel subset $A \subset F_n$, we set $[A]_n := \{x = (x_i)_{i=n}^{\infty} \in X_n \mid x_n \in A\}$ and call it an n -cylinder in X . Clearly,

$$[A]_n = \bigsqcup_{c \in C_{n+1}} [A + c]_{n+1}.$$

Notice also that

$$(1-4) \quad T_g[A]_n = [A + g]_n \text{ for all } g \in \mathbb{R}^d \text{ and } A \subset F_n \cap (F_n - g), \quad n \in \mathbb{N}.$$

The sequence of all n -cylinders approximates the entire Borel σ -algebra on X when $n \rightarrow \infty$.

We state without proof the following standard lemma (see, e.g., Lemma 2.4 from [Da1]).

Lemma 1.3. *Let $\mathcal{P}_n$ be a finite partition of F_n into parallelepipeds such that for each atom Δ of $\mathcal{P}_n$ and an element $c \in C_{n+1}$, the parallelepiped $\Delta + c$ is $\mathcal{P}_{n+1}$ -measurable and the maximal diameter of the atoms in $\mathcal{P}_n$ goes to zero. Let S be a measure preserving transformation of X . Then the following hold:*

- (i) *The sequence of collections of n -cylinders $\{[A]_n \mid A \subset F_n \text{ is } \mathcal{P}_n\text{-measurable}\}$ approximates the entire σ -algebra $\mathfrak{B}$ as $n \rightarrow \infty$.*
- (ii) *If for each pair of atoms $\Delta_1, \Delta_2 \in \mathcal{P}_n$, there are a subset $A \subset [\Delta_1]_n$ and a μ -preserving one-to-one map $\gamma: A \rightarrow [\Delta_2]_n$ such that $\mu(A) > 0.5\mu([\Delta_1]_n)$ and $\gamma x \in \{S^i x \mid i \in \mathbb{Z}\}$ for all $x \in A$, then S is ergodic.*

We will also use the following property of the (C, F) -actions. If T is associated with $(C_{n+1}, F_n)_{n \geq 0}$, then for each $p > 1$, the product action

$$(T_{t_1} \times \cdots \times T_{t_p})_{(t_1, \dots, t_p) \in (\mathbb{R}^d)^p}$$

is the (C, F) -action of $(\mathbb{R}^d)^p$ associated with $(C_{n+1}^p, F_n^p)_{n \geq 0}$. The upper index p means the p -th Cartesian power.

2. TWO AUXILIARY FACTS

Given a σ -finite measure space (X, μ) , we denote by $\text{Aut}(X, \mu)$ the group of all μ -preserving (invertible) transformations of X . It is a Polish group when endowed with the *weak topology* [Aa]. Recall that the weak topology is the weakest topology in which the maps

$$\text{Aut}(X, \mu) \ni T \mapsto \mu(TA \cap B) \in \mathbb{R}$$

are continuous for all subsets $A, B \subset X$ of finite measure.

Given $S \in \text{Aut}(X, \mu)$ and two subsets $A, B \subset X$ with $\mu(A) = \mu(B) < \infty$, we define subsets $A_0, A_1, \dots$ of A as follows:

$$(2-1) \quad \begin{aligned} A_0 &:= A \cap B, \\ A_i &:= \left(A \setminus \bigcup_{j=0}^{i-1} A_j \right) \cap S^{-i} \left(B \setminus \bigcup_{j=0}^{i-1} S^j A_j \right), \quad i > 0. \end{aligned}$$

We now let $\mathcal{N}_{S,A,B} := \min\{i \geq 0 \mid \mu(A_0 \sqcup \dots \sqcup A_i) > 0.5\mu(A)\}$. If S is ergodic, then $A = \bigcup_{i \geq 0} A_i$ and hence $\mathcal{N}_{S,A,B}$ is well defined. Denote by $\mathcal{E}$ the subset of all ergodic transformations in $\text{Aut}(X, \mu)$. It is well known that $\mathcal{E}$ is a dense G_δ in $\text{Aut}(X, \mu)$. Since for each $i \geq 0$ the map

$$\text{Aut}(X, \mu) \ni S \mapsto \mu(A_0 \sqcup \dots \sqcup A_i) \in \mathbb{R}$$

is continuous, we obtain the following lemma.

Lemma 2.1. *The map $\mathcal{E} \ni S \mapsto \mathcal{N}_{S,A,B} \in \mathbb{R}$ is upper semicontinuous for all subsets $A, B \subset X$ with $\mu(A) = \mu(B) < \infty$.*

In the case of (C, F) -actions we can say more about the “structure” of the sets A_i , $i = 0, \dots, \mathcal{N}_{S,A,B}$. For $q = (q_1, \dots, q_d) \in \mathbb{R}^d$, we let $\|q\| := \max_{1 \leq i \leq d} |q_i|$.

Lemma 2.2. *Let $(X, \mu, (T_t)_{t \in \mathbb{R}^d})$ be a (C, F) -action of $\mathbb{R}^d$ associated with a sequence $(C_{n+1}, F_n)_{n \geq 0}$ such that*

$$(2-2) \quad a + F_n + C_{n+1} \subset F_{n+1}$$

for each $a = (a_1, \dots, a_p)$ with $a_i \geq 0$, $i = 1, \dots, p$, and $\|a\| \leq 1$. Fix two n -cylinders A and B of equal measure and a transformation $S = T_q$ for some $q \in \mathbb{R}_+^d$. Then the subsets $A_0, A_1, \dots, A_{\mathcal{N}(S,A,B)}$ defined by (2-1) are $(n + Q \cdot \mathcal{N}(S, A, B))$ -cylinders, where Q is any integer greater than $\|q\|$.

Proof. We let $N := Q \cdot \mathcal{N}(S, A, B)$. If $A = [\tilde{A}]_n$ for some $\tilde{A} \subset F_n$, then $A = [\hat{A}]_{n+N}$, where $\hat{A} := \tilde{A} + C_{n+1} + \dots + C_{n+N} \subset F_{n+N}$. From (1-2) and (2-2) we deduce that the sets $\hat{A} + q, \dots, \hat{A} + \mathcal{N}(S, A, B)q$ are all contained in F_{n+N} . It remains to use (2-1) and (1-4). $\square$

We also note that $S^i A_i \subset B$ and $S^i A_i \cap S^j A_j = \emptyset$ for all $i, j = 0, \dots, \mathcal{N}(S, A, B)$.

3. PROOF OF THE MAIN RESULT

Theorem 3.1. *There exists a (C, F) -flow $T = (T_t)_{t \in \mathbb{R}}$ such that each transformation T_t , $t \neq 0$, has infinite ergodic index.*

Proof. We will construct this flow via an inductive procedure. Fix a sequence of integers $(p_n)_{n \geq 1}$ in which every integer greater than 1 occurs infinitely many times. Suppose that after $n - 1$ steps of the construction we have already defined

$$(3-1) \quad F_0, C_1, F_1, \dots, C_{m_{n-1}}, F_{m_{n-1}}.$$

Suppose also that for each $0 \leq i \leq m_{n-1}$, a finite partition $\mathcal{P}_i$ of F_i into intervals is chosen in such a way that

- the interval $\Delta + c$ is $\mathcal{P}_{i+1}$ -measurable for each atom Δ of $\mathcal{P}_i$, $0 \leq i < m_{n-1}$, and each $c \in C_{i+1}$ and
- the length of any atom of $\mathcal{P}_i$ is no more than i^{-1} , $1 \leq i \leq m_{n-1}$.

Step n. Consider a rank-one weakly mixing finite measure preserving (C, F) -flow $T^{(n)} = (T_t^{(n)})_{t \in \mathbb{R}}$ associated with a sequence $(C_{k+1,n}, F_{k,n})_{k \geq 0}$ such that $F_{0,n} := F_{m_{n-1}}$. Examples of weakly mixing rank-one finite measure preserving flows are well known—see, e.g., [dJP]. In [DaS2] one can find explicit (C, F) -construction of mixing finite measure preserving flows. Let $(X^{(n)}, \mu_n)$ be the space of this action. Since $T^{(n)}$ is weakly mixing, it follows that for each $t > 0$, the transformation

$$(3-2) \quad S_t := T_t^{(n)} \times \dots \times T_t^{(n)} \text{ (} p_n \text{ times)}$$

of the product space $(X^{(n)}, \mu_n)^{p_n}$ is ergodic. We note that this space is the space of the (C, F) -action of $\mathbb{R}^{p_n}$ associated with the sequence $(C_{k+1,n}^{p_n}, F_{k,n}^{p_n})_{k \geq 0}$ (see our remark at the end of §1). Given $k \geq 0$, let $\mathcal{P}_{k,n}$ be a finite partition of $F_{k,n}$ into intervals such that

- $\mathcal{P}_{0,n} = \mathcal{P}_{m_{n-1}}$,
- the interval $\Delta + c$ is $\mathcal{P}_{k+1,n}$ -measurable for each atom Δ of $\mathcal{P}_{k,n}$ and $c \in C_{k+1,n}$ and
- the length of any atom of $\mathcal{P}_{k,n}$ is no more than $(m_{n-1} + k)^{-1}$.

We now let $\mathcal{P}_{k,n}^{p_n} := \mathcal{P}_{k,n} \times \dots \times \mathcal{P}_{k,n}$ (p_n times). Then $\mathcal{P}_{k,n}^{p_n}$ is a finite partition of $F_{k,n}^{p_n}$ into parallelepipeds and

- the parallelepiped $\Delta + c$ is $\mathcal{P}_{k+1,n}^{p_n}$ -measurable for each atom Δ of $\mathcal{P}_{k,n}^{p_n}$ and $c \in C_{k+1,n}^{p_n}$ and
- the diameter of an atom of $\mathcal{P}_{k,n}^{p_n}$ is no more than $(m_{n-1} + k)^{-p_n}$.

Denote by D_n the maximum of $\mathcal{N}(S_t, [\Delta]_0, [\Delta']_0)$ when Δ and Δ' run independently over the atoms of $\mathcal{P}_{0,n}^{p_n}$ and t runs over the segment $[n^{-1}, n] \subset \mathbb{R}$. It exists by Lemma 2.1. It now follows from Lemma 2.2 that for any pair of parallelepipeds $\Delta, \Delta' \in \mathcal{P}_{0,n}^{p_n}$ and a real $t \in [n^{-1}, n]$, there exist nD_n -cylinders $A_1, \dots, A_{D_n} \subset [\Delta]_0$ such that

$$(3-3) \quad \mu_n^{p_n} \left(\bigcup_{i=1}^{D_n} A_i \right) > \frac{1}{2} \mu_n^{p_n}([\Delta]_0),$$

$$S_t^i A_i \subset [\Delta']_0 \text{ for each } 1 \leq i \leq D_n \text{ and}$$

$$S_t^i A_i \cap S_t^j A_j = \emptyset \text{ if } 1 \leq i \neq j \leq D_n.$$

We now “continue” the sequence (3-1) by setting

$$C_{m_{n-1}+1} := C_{1,n}, \quad F_{m_{n-1}+1} := F_{1,n}, \dots, \quad C_{m_{n-1}+nD_n} := C_{nD_n,n}.$$

Next, to define $F_{m_{n-1}+nD_n}$ we “double” the set $F_{nD_n,n}$, i.e.

$$(3-4) \quad F_{m_{n-1}+nD_n} := [0, 2a) \text{ if } F_{nD_n,n} = [0, a) \text{ for some } a > 0.$$

It remains to put $m_n := m_{n-1} + nD_n$. The n -th step is now completed.

Continuing this procedure infinitely many times, we obtain the entire sequence $(C_{i+1}, F_i)_{i=0}^\infty$. Denote by $T = (T_t)_{t \in \mathbb{R}}$ the associated (C, F) -flow. Let (X, μ) be the space of this flow. It follows from (3-4) that $\lambda_{\mathbb{R}}(F_i) > 2\lambda_{\mathbb{R}}(F_{i-1})\#C_i$ for infinitely many i . Hence $\mu(X) = \infty$. Moreover, a finite partition $\mathcal{P}_i$ of F_i into intervals is fixed such that the conditions of Lemma 1.3 are satisfied. Next, there are one-to-one correspondences (natural identifications) between

- the collection of 0-cylinders in $X^{(n)}$ and the collection of m_{n-1} -cylinders in X and
- the collection of nD_n -cylinders in $X^{(n)}$ and the collection of m_n -cylinders in X .

Moreover, the “dynamics” of $T^{(n)}$ on the nD_n -cylinders is the same as the dynamics of T on the m_n -cylinders. This means the following: if $A, B \subset F_{nD_n}^{(n)}$ and $[B]_{nD_n} = T_w^{(n)}[A]_{nD_n}$ for some $w \in \mathbb{R}$, then $[B]_{m_n} = T_w[A]_{m_n}$. Therefore we deduce from (3-3) that for any pair of parallelepipeds $\Delta, \Delta' \in \mathcal{P}_{m_{n-1}}^{p_n}$ and a real $t \in [n^{-1}, n]$, there exist m_n -cylinders $A_1, \dots, A_{D_n} \subset [\Delta]_{m_{n-1}}$ such that

$$\mu^{p_n} \left(\bigcup_{i=1}^{D_n} A_i \right) > \frac{1}{2} \mu^{p_n}([\Delta]_{m_{n-1}}),$$

$$V_t^i A_i \subset [\Delta']_{m_{n-1}} \text{ for each } 1 \leq i \leq D_n \text{ and}$$

$$V_t^i A_i \cap V_t^j A_j = \emptyset \text{ if } 1 \leq i \neq j \leq D_n,$$

where $V_t := T_t \times \dots \times T_t$ (p_n times). Fix $p > 0$. Passing to a subsequence where $p_n = p$ we now deduce from Lemma 1.3(ii) that $T_t \times \dots \times T_t$ (p times) is ergodic for each $t > 0$. Hence T_t has infinite ergodic index. $\square$

4. CONCLUDING REMARKS

4.1. When constructing T , we use only finitely many initial terms of the sequence $(C_{k+1}^{(n)}, F_k^{(n)})_{k \geq 0}$ for each $n > 0$. However to determine “where to stop” (i.e. to determine D_n) we use the weak mixing properties of the auxiliary flow $T^{(n)}$ which depends on the entire *infinite* sequence $(C_{k+1}^{(n)}, F_k^{(n)})_{k \geq 0}$. No upper bound on D_n is found. This means that the construction of T is not *effective*. In this connection we raise a question:

Is it possible to find an effective construction for the flow from Theorem 0.1?

We note that the construction in [DaSo] is effective.

4.2. It is possible to strengthen Theorem 0.1 by replacing the infinite ergodic index with a stronger property of power weak mixing. Recall that a measure preserving transformation S is called power weak mixing if for each finite sequence $n_1, \dots, n_k$ of nonzero integers the transformation $S^{n_1} \times \dots \times S^{n_k}$ is ergodic. Only a slight modification of our argument is needed to show the following theorem. (The idea is to define S_t in (3-2) as the product of *different* nonzero powers $T_t^{(n)}$. The product is ergodic since $T^{(n)}$ is finite measure preserving and weakly mixing.)

Theorem 4.1. *There exists a rank-one infinite σ -finite measure preserving flow $T = (T_t)_{t \in \mathbb{R}}$ such that the transformation T_t is power weakly mixing for each $t \neq 0$.*

Also, it is easy to extend Theorems 0.1 and 4.1 to actions of $\mathbb{R}^d$.

Theorem 4.2. *For each $d > 1$, there exists a rank-one infinite σ -finite measure preserving action $T = (T_g)_{g \in \mathbb{R}^d}$ of $\mathbb{R}^d$ such that the transformation T_g is power weakly mixing for each $g \neq 0$.*

We leave the proofs of Theorems 4.1 and 4.2 to the reader.

REFERENCES

- [Aa] J. Aaronson, *An introduction to infinite ergodic theory*, Amer. Math. Soc., Providence, R.I., 1997. MR1450400 (99d:28025)
- [Da1] A. I. Danilenko, *Funny rank one weak mixing for nonsingular Abelian actions*, Isr. J. Math. **121** (2001), 29–54. MR1818379 (2002a:37003)
- [Da2] ———, *(C, F)-actions in ergodic theory*, Geometry and Dynamics of Groups and Spaces, Progr. Math., vol. 265, 2008, pp. 325–351. MR2402408 (2009g:37003)
- [DaS1] A. I. Danilenko and C. E. Silva, *Multiple and polynomial recurrence for abelian actions in infinite measure*, J. London Math. Soc. **69** (2004), 183–200. MR2025335 (2005a:37008)
- [DaS2] ———, *Mixing rank-one actions of locally compact Abelian groups*, Ann. Inst. H. Poincaré, Probab. Statist. **43** (2007), 375–398. MR2329508 (2008j:37005)
- [DaS3] ———, *Ergodic theory: non-singular transformations*, Encyclopedia of complexity and systems science, Springer, 2009.
- [DaSo] A. I. Danilenko and A. Solomko, *Infinite measure preserving flows with infinite ergodic index*, Colloq. Math. **115** (2009), 13–19. MR2475836 (2009k:37014)
- [DaR] A. I. Danilenko and V. V. Ryzhikov, *Mixing constructions with infinite invariant measure and spectral multiplicities*, Ergod. Th. & Dynam. Syst. (to appear), doi:10.1017/S0143385710000052.
- [dJ] A. del Junco, *A simple map with no prime factors*, Isr. J. Math. **104** (1998), 301–320. MR1622315 (99d:28037)
- [dJP] A. del Junco and K. K. Park, *An example of a measure preserving flow with minimal self-joinings*, J. d'Analyse Mathématique **42** (1984), 199–209. MR729410 (85e:28029)
- [I–W] S. Iams, B. Katz, C. E. Silva, B. Street and K. Wickelgren, *On weakly mixing and doubly ergodic nonsingular actions*, Colloq. Math. **103** (2005), 247–264. MR2197854 (2007a:37006)
- [KP] S. Kakutani and W. Parry, *Infinite measure preserving transformations with “mixing”*, Bull. Amer. Math. Soc. **69** (1963), 752–756. MR0153815 (27:3776)
- [Ry1] V. V. Ryzhikov, *Homogeneous spectrum, disjointness of convolutions, and mixing properties of dynamical systems*, Selected Russian Mathematics **1** (1999), 13–24.
- [Ry2] ———, *Weak limits of powers, the simple spectrum of symmetric products and mixing constructions of rank 1*, Sb. Math. **198** (2007), 733–754. MR2354530 (2008m:37008)

INSTITUTE FOR LOW TEMPERATURE PHYSICS AND ENGINEERING OF NATIONAL ACADEMY OF SCIENCES OF UKRAINE, 47 LENIN AVENUE, KHARKOV, 61164, UKRAINE

E-mail address: alexandre.danilenko@gmail.com

DEPARTMENT OF MATHEMATICS, COLLEGE OF NATURAL SCIENCE, AJOU UNIVERSITY, SUWON 442-749, KOREA

E-mail address: kkpark@madang.ajou.ac.kr