ON ADDITIVE COMPLEMENTS. II

YONG-GAO CHEN AND JIN-HUI FANG

(Communicated by Matthew A. Papanikolas)

ABSTRACT. Two infinite sequences A and B of non-negative integers are called additive complements if their sum contains all sufficiently large integers. Let $A(x)$ and $B(x)$ be the counting functions of A and B and let $\limsup_{x \rightarrow \infty} A(x)B(x)/x = \alpha(A, B)$. Recently, the authors [Proceedings of the American Mathematical Society 138 (2010), 1923-1927] proved that for additive complements A and B , if $\alpha(A, B) < 5/4$ or $\alpha(A, B) > 2$, then $A(x)B(x) - x \rightarrow +\infty$ as $x \rightarrow \infty$. In this paper, we prove that for any $\varepsilon > 0$ there exist additive complements A and B with $2 - \varepsilon < \alpha(A, B) < 2$ and $A(x)B(x) - x = 1$ for infinitely many positive integers x .

1. INTRODUCTION

Two infinite sequences A and B of non-negative integers are called additive complements if their sum contains all sufficiently large integers.

Let $A(x)$ and $B(x)$ be the counting functions of A and B . Namely,

$$A(x) = \sum_{\substack{a \leq x \\ a \in A}} 1 \quad \text{and} \quad B(x) = \sum_{\substack{b \leq x \\ b \in B}} 1.$$

For the construction of additive complements A and B with $A(x)B(x) \sim x$, one may refer to [6].

In 1964, Danzer [1] conjectured that for additive complements A and B , if

$$\limsup_{x \rightarrow \infty} \frac{A(x)B(x)}{x} \leq 1,$$

then

$$A(x)B(x) - x \rightarrow +\infty \quad \text{as} \quad x \rightarrow +\infty.$$

(See also [2], [4, p. 75] and [5].) In [7], Sárközy and Szemerédi proved this conjecture. Recently, the authors [3] proved that

Received by the editors April 14, 2010.

2010 *Mathematics Subject Classification*. Primary 11B13, 11B34.

Key words and phrases. Additive complements, sequences, counting functions.

This work was supported by the National Natural Science Foundation of China, Grant No. 10771103.

©2010 American Mathematical Society
 Reverts to public domain 28 years from publication

Theorem A. For additive complements A and B , if

$$\limsup_{x \rightarrow \infty} \frac{A(x)B(x)}{x} > 2 \text{ or } \limsup_{x \rightarrow \infty} \frac{A(x)B(x)}{x} < \frac{5}{4},$$

then $A(x)B(x) - x \rightarrow +\infty$ as $x \rightarrow \infty$.

Once we conjectured that if A and B are additive complements, then $A(x)B(x) - x \rightarrow +\infty$ as $x \rightarrow +\infty$. After paper [3] went to print, we found a counterexample, so we removed the conjecture from [3] when we checked the proofs. In this paper, as a complement to [3], we prove the following result.

Theorem 1. For any integer a with $a \geq 2$, there exist additive complements A and B such that $A(x)B(x) - x = 1$ for infinitely many positive integers x and

$$\limsup_{x \rightarrow \infty} \frac{A(x)B(x)}{x} = \frac{2a+2}{a+2}.$$

2. PROOFS OF THEOREM 1

We will use the following lemmas. Their proofs are clear.

Lemma 1. For any integers a_1, a_2, b_1, b_2 and u , if $a_1b_2 - a_2b_1 \geq 0$ and $a_2x + b_2 > 0$ for all $0 \leq x \leq u$, then for all $0 \leq x \leq u$ we have

$$\frac{a_1x + b_1}{a_2x + b_2} \leq \frac{a_1u + b_1}{a_2u + b_2}.$$

Lemma 2. For any integers a_1, a_2, b_1, b_2 and v , if $a_2 > 0$, $a_1b_2 - a_2b_1 \geq 0$ and $a_2v + b_2 > 0$, then

$$\frac{a_1v + b_1}{a_2v + b_2} \leq \frac{a_1}{a_2}.$$

Proof of Theorem 1. For any integer a with $a \geq 2$, assume that

$$A_j = \left\{ \sum \varepsilon_i a^{2i+j} \mid \varepsilon_i = 0, 1, \dots, a-1 \right\}, \quad j = 0, 1,$$

where the summation is a finite sum. It is clear that A_0 and A_1 are additive complements.

For $x_k = a^{2k} - 1$ we have

$$A_0(x_k) = A_1(x_k) = a^k, \quad A_0(x_k)A_1(x_k) - x_k = 1.$$

Now we prove that

$$\limsup_{x \rightarrow \infty} \frac{A_0(x)A_1(x)}{x} = \frac{2a+2}{a+2}.$$

For $x_{l,j} = a^{2l+j} + (a-1) \sum_{i=0}^{l-1} a^{2i+j}$ ($j = 0, 1$), we have $A_j(x_{l,j}) = 2a^l$, $A_{1-j}(x_{l,j}) = a^{l+j}$. Then

$$\frac{A_0(x_{l,j})A_1(x_{l,j})}{x_{l,j}} = \frac{2}{(1 + \frac{1}{a+1}) - \frac{1}{a+1} \frac{1}{a^{2l}}} \rightarrow \frac{2a+2}{a+2}, \quad l \rightarrow \infty.$$

Now it suffices to prove that for any positive integers $x \neq x_{l,j}$ we have

$$\frac{A_0(x)A_1(x)}{x} \leq \frac{2a+2}{a+2}.$$

If a positive integer $x \notin A_0 \cup A_1$, then

$$\frac{A_0(x)A_1(x)}{x} = \frac{A_0(x-1)A_1(x-1)}{x} < \frac{A_0(x-1)A_1(x-1)}{x-1}.$$

So we may assume that $x \in A_0 \cup A_1$. Hence, there exist integers l, j with $j \in \{0, 1\}$ and $x = \epsilon_j a^j + \epsilon_{2+j} a^{2+j} + \cdots + \epsilon_{2l+j} a^{2l+j}$, where $\epsilon_{2l+j} > 0$ and $\epsilon_{2i+1+j} = 0$ for all i . Then $A_{1-j}(x) = a^{l+j}$. If $\epsilon_{2l+j} \geq 2$, then, by $A_j(x) \leq \epsilon_{2l+j} a^l + a^l$, we have

$$\frac{A_j(x)A_{1-j}(x)}{x} \leq \frac{a^{2l+j}\epsilon_{2l+j} + a^{2l+j}}{x} \leq \frac{a^{2l+j}\epsilon_{2l+j} + a^{2l+j}}{a^{2l+j}\epsilon_{2l+j}} \leq \frac{3}{2} \leq \frac{2a+2}{a+2}.$$

So we only need to consider $\epsilon_{2l+j} = 1$. Since $x \neq x_{l,j}$, we may assume that

$$x = \sum_{i=0}^{m-1} \epsilon_{2i+j} a^{2i+j} + (a-1) \sum_{i=m}^{l-1} a^{2i+j} + a^{2l+j},$$

where $0 < m \leq l$ and $\epsilon_{2m+j-2} < a-1$ (if $m = l$, then the second sum is zero). Thus

$$A_j(x) \leq a^l + (a-1) \sum_{i=m}^{l-1} a^i + \epsilon_{2m+j-2} a^{m-1} + a^{m-1} = 2a^l - a^m + (\epsilon_{2m+j-2} + 1)a^{m-1}.$$

Hence by Lemmas 1 ($x = \epsilon_{2m+j-2}$) and 2 ($v = a^{2l+j}$) we have

$$\begin{aligned} \frac{A_{1-j}(x)A_j(x)}{x} &\leq \frac{a^{l+j}(2a^l - a^m + (\epsilon_{2m+j-2} + 1)a^{m-1})}{a^{2l+j} + (a-1)(a^{2m+j} + \cdots + a^{2l+j-2}) + \epsilon_{2m+j-2}a^{2m+j-2}} \\ &= \frac{2a^{2l+j} - a^{m+l+j} + (\epsilon_{2m+j-2} + 1)a^{m+l+j-1}}{(1 + \frac{1}{a+1})a^{2l+j} - \frac{1}{a+1}a^{2m+j} + \epsilon_{2m+j-2}a^{2m+j-2}} \\ &\leq \frac{2a^{2l+j} - a^{m+l+j} + (a-2+1)a^{m+l+j-1}}{(1 + \frac{1}{a+1})a^{2l+j} - \frac{1}{a+1}a^{2m+j} + (a-2)a^{2m+j-2}} \\ &\leq \frac{2a+2}{a+2}. \end{aligned}$$

This completes the proof of Theorem 1. $\square$

REFERENCES

- [1] L. Danzer, Über eine Frage von G. Hanani aus der additiven Zahlentheorie, J. Reine Angew. Math. 214/215(1964), 392-394. MR0161830 (28:5034)
- [2] P. Erdős, R. L. Graham, Old and New Problems and Results in Combinatorial Number Theory, Monographies de L'Enseignement Mathématique, 28, Université de Genève, Geneva (1980). MR592420 (82j:10001)
- [3] J. H. Fang, Y. G. Chen, On additive complements, Proc. Amer. Math. Soc. 138(2010), 1923-1927. MR2596025
- [4] H. Halberstam, K. F. Roth, Sequences, 2nd edition, Springer-Verlag (1983). MR687978 (83m:10094)
- [5] W. Narkiewicz, Remarks on a conjecture of Hanani in additive number theory, Colloq. Math. 7(1960), 161-165. MR0112876 (22:3722)
- [6] I. Z. Ruzsa, Additive completion of lacunary sequences, Combinatorica 21(2)(2001), 279-291. MR1832452 (2002f:11008)
- [7] A. Sárközy, E. Szemerédi, On a problem in additive number theory, Acta Math. Hungar. 64(3)(1994), 237-245. MR1275641 (95c:11121)

SCHOOL OF MATHEMATICAL SCIENCES, NANJING NORMAL UNIVERSITY, NANJING 210046, PEOPLE'S REPUBLIC OF CHINA

E-mail address: ygchen@njnu.edu.cn

DEPARTMENT OF MATHEMATICS, NANJING UNIVERSITY OF INFORMATION SCIENCE & TECHNOLOGY, NANJING 210044, PEOPLE'S REPUBLIC OF CHINA

E-mail address: fangjinhui1114@163.com