CHARACTERIZATION OF SUBDIAGONAL ALGEBRAS

TURDEBEK N. BEKJAN

(Communicated by Marius Junge)

ABSTRACT. Let $\mathcal{M}$ be a finite von Neumann algebra with a faithful normal tracial state τ , and let $\mathcal{A}$ be a tracial subalgebra of $\mathcal{M}$. We show that $\mathcal{A}$ has L^p -factorization ($1 \leq p < \infty$) if and only if $\mathcal{A}$ is a subdiagonal algebra. Also, we obtain some characterizations of subdiagonal algebras.

0. INTRODUCTION

Let $\mathcal{M}$ be a finite von Neumann algebra with a faithful normal tracial state τ . In [1], Arveson introduced the notion of finite, maximal, subdiagonal algebras $\mathcal{A}$ of $\mathcal{M}$, as noncommutative analogues of weak-* Dirichlet algebras. Subsequently several authors studied the (noncommutative) $\mathcal{H}^p$ -spaces associated with such algebras ([9, 11, 12, 13, 14, 16, 17, 18]). Arveson [1] proved a Szegő type factorization theorem. Some extensions can be found in [8, 18], and [15]. Labuschagne [10] proved a non-commutative version of Szegő's theorem. In the recent articles [3, 4, 5], among other things, Blecher and Labuschagne studied tracial subalgebras of $\mathcal{M}$ and gave several characterizations of subdiagonal algebras. They proved that if a tracial subalgebra $\mathcal{A}$ has L^∞ -factorization, then $\mathcal{A}$ is a subdiagonal algebra. We will consider the L^p -factorization ($0 < p < \infty$) property of tracial subalgebras. This paper is organized as follows. Section 1 contains some preliminary definitions. In section 2, we prove that if a tracial subalgebra $\mathcal{A}$ has L^p -factorization ($1 \leq p < \infty$), then $\mathcal{A}$ is a subdiagonal algebra. In section 3, we consider tracial subalgebras, which satisfy L^2 -density. We show that if a tracial subalgebra $\mathcal{A}$ has L^p -factorization ($0 < p < 1$) and satisfies L^2 -density, then $\mathcal{A}$ is a subdiagonal algebra.

1. PRELIMINARIES

Throughout this paper, we denote by $\mathcal{M}$ a finite von Neumann algebra on a Hilbert space $\mathcal{H}$ with a faithful normal tracial state τ . For $0 < p < \infty$ we denote by $L^p(\mathcal{M})$ the usual noncommutative L^p -space associated with $(\mathcal{M}, \tau)$. Recall that $L^\infty(\mathcal{M}) = \mathcal{M}$, equipped with the operator norm. It is well-known that $L^p(\mathcal{M})$ is a Banach space under $\|\cdot\|_p$ ($1 \leq p \leq \infty$) satisfying all the expected properties such as duality of $L^p(\mathcal{M})$ and $L^q(\mathcal{M})$ for $\frac{1}{p} + \frac{1}{q} = 1$ (see [7, 20]).

Received by the editors November 30, 2009 and, in revised form, April 13, 2010.

2010 *Mathematics Subject Classification.* Primary 46L51, 46L52, 47L75.

Key words and phrases. von Neumann algebras, tracial subalgebra, subdiagonal algebra, L^2 -density, L^p -factorization.

The author was partially supported by NSFC grant No. 10761009.

©2010 American Mathematical Society
 Reverts to public domain 28 years from publication

For a subset K of $L^p(\mathcal{M})$, put $J(K) = \{x^* : x \in K\}$, $K^{-1} = \{x : x, x^{-1} \in K\}$, $K^+ = \{x : x \geq 0, x \in K\}$, and $[K]_p$ the closed linear span of K in $L^p(\mathcal{M})$. (Here $[K]_\infty$ is the weak* closure of K .)

Given a von Neumann subalgebra $\mathcal{N}$ of $\mathcal{M}$, an expectation $\mathcal{E} : \mathcal{M} \rightarrow \mathcal{N}$ is defined to be a positive linear map which preserves the identity and satisfies $\mathcal{E}(xy) = x\mathcal{E}(y)$ for all $x \in \mathcal{N}$ and $y \in \mathcal{M}$. Since $\mathcal{E}$ is positive it is Hermitian; i.e. $\mathcal{E}(x)^* = \mathcal{E}(x^*)$ for all $x \in \mathcal{M}$. Hence $\mathcal{E}(yx) = \mathcal{E}(y)x$ for all $x \in \mathcal{N}$ and $y \in \mathcal{M}$. For a complete study of $\mathcal{E}$, we refer to [1, 12].

Definition 1.1. Let $\mathcal{A}$ be a w^* -closed unital subalgebra of $\mathcal{M}$. If there exists a linear projection $\mathcal{E}$ from $\mathcal{A}$ onto $\mathcal{D} = \mathcal{A} \cap J(\mathcal{A})$ such that

- (i) $\mathcal{E}(xy) = \mathcal{E}(x)\mathcal{E}(y)$, $\forall x, y \in \mathcal{A}$;
- (ii) $\tau \circ \mathcal{E} = \tau$,

then $\mathcal{A}$ is called a tracial subalgebra of $\mathcal{M}$.

Definition 1.2. Let $\mathcal{A}$ be a w^* -closed unital subalgebra of $\mathcal{M}$, and let $\mathcal{E}$ be a faithful, normal expectation from $\mathcal{M}$ onto the diagonal von Neumann algebra $\mathcal{D} = \mathcal{A} \cap J(\mathcal{A})$. Then $\mathcal{A}$ is a finite subdiagonal subalgebra of $\mathcal{M}$ with respect to $\mathcal{E}$ if:

- (i) $\mathcal{A} + J(\mathcal{A})$ is w^* -dense in $\mathcal{M}$;
- (ii) $\mathcal{E}(xy) = \mathcal{E}(x)\mathcal{E}(y)$, $\forall x, y \in \mathcal{A}$;
- (iii) $\tau \circ \mathcal{E} = \tau$.

By Theorem 5.6 of [3], $\mathcal{E}$ is precisely the restriction to $\mathcal{A}$ of the unique faithful normal conditional expectation Φ from $\mathcal{M}$ onto $\mathcal{D}$ such that $\tau = \tau \circ \Phi$. Hence we may continue to write Φ as $\mathcal{E}$, and we call this extension the conditional expectation onto $\mathcal{D}$. It is well-known that $\mathcal{E}$ extends to a contractive projection from $L_p(\mathcal{M})$ onto $L_p(\mathcal{D})$ for every $1 \leq p \leq \infty$. The extension will still be denoted by $\mathcal{E}$.

Let $\mathcal{A}_0 = \mathcal{A} \cap \ker(\mathcal{E})$. We say that $\mathcal{A}$ is τ -maximal if

$$(1.1) \quad \mathcal{A} = \{x \in \mathcal{M} : \tau(xy) = 0, \forall y \in \mathcal{A}_0\}.$$

2. FACTORIZATION

Let $\mathcal{A}$ be a tracial subalgebra of $\mathcal{M}$. For $0 < p < \infty$, we write $\mathcal{A}_p$ for $[\mathcal{A}]_p \cap \mathcal{M}$.

Lemma 2.1. Let $\mathcal{A}$ be a tracial subalgebra of $\mathcal{M}$. Then $\mathcal{A}_p$ is a tracial subalgebra of $\mathcal{M}$ for $1 \leq p < \infty$.

Proof. The proof is the same as that of Theorem 4.4 in [4]. We only need to prove the following:

$$\mathcal{E}(ab) = \mathcal{E}(a)\mathcal{E}(b), \forall a, b \in \mathcal{A}_p.$$

Indeed, let $b \in \mathcal{A}_p$. Then there exists a sequence $\{b_n\}$ in $\mathcal{A}$ such that $b_n \rightarrow b$ in $L_p(\mathcal{M})$. So for all $a \in \mathcal{A}$ we have $ab_n \in \mathcal{A}$, and $ab_n \rightarrow ab$ in $L_p(\mathcal{M})$. Thus

$$\mathcal{E}(ab) = \lim_{n \rightarrow \infty} \mathcal{E}(ab_n) = \lim_{n \rightarrow \infty} \mathcal{E}(a)\mathcal{E}(b_n) = \mathcal{E}(a)\mathcal{E}(b).$$

Hence, by what we just proved,

$$\mathcal{E}(ab) = \lim_{n \rightarrow \infty} \mathcal{E}(ab_n) = \lim_{n \rightarrow \infty} \mathcal{E}(a)\mathcal{E}(b_n) = \mathcal{E}(a)\mathcal{E}(b), \forall a \in \mathcal{A}_p.$$

□

Definition 2.2. Let $0 < p \leq \infty$, and let $\mathcal{A}$ be a tracial subalgebra of $\mathcal{M}$. We say that $\mathcal{A}$ has L^p -factorization, if for all $x \in L^p(\mathcal{M})^{-1}$, there is a unitary $u \in \mathcal{M}$ and $a \in [\mathcal{A}]_p^{-1}$ such that $x = ua$.

Proposition 2.3. *Let $\mathcal{A}$ be a tracial subalgebra of $\mathcal{M}$. Consider the following conditions:*

- (i) *For some $1 \leq p_0 < \infty$, $\mathcal{A}$ has L^{p_0} -factorization.*
- (ii) *For some $0 < p_0 < 1$, $\mathcal{A}$ has L^{p_0} -factorization and $\mathcal{A}_{p_0}$ is a tracial subalgebra of $\mathcal{M}$.*

Then either one of the conditions (i) or (ii) implies that $\mathcal{A}$ has L^p -factorization for all $0 < p < p_0$.

Proof. Let $x \in \mathcal{M}^{-1} \subset L^{p_0}(\mathcal{M})^{-1}$. Then there is a unitary $u \in \mathcal{M}$ and $a \in [\mathcal{A}]_{p_0}^{-1}$ such that $x = ua$, since $\mathcal{A}$ has L^{p_0} -factorization. So $a \in \mathcal{A}_{p_0}^{-1}$, and therefore $\mathcal{A}_{p_0}$ has L^∞ -factorization. By Theorem 1.1 of [4], $\mathcal{A}_{p_0}$ is a subdiagonal subalgebra of $\mathcal{M}$. Let $x \in L^p(\mathcal{M})^{-1}$. By Theorem 3.1 of [2], there is a unitary $u \in \mathcal{M}$ and $a \in [\mathcal{A}_{p_0}]_p^{-1}$ such that $x = ua$. On the other hand, we have

$$[\mathcal{A}]_p \subset [\mathcal{A}_{p_0}]_p \subset [[\mathcal{A}]_{p_0}]_p = [\mathcal{A}]_p.$$

Thus $a \in [\mathcal{A}]_p^{-1}$, and so $\mathcal{A}$ has L^p -factorization. $\square$

Theorem 2.4. *Let $\mathcal{A}$ be a tracial subalgebra of $\mathcal{M}$. Then the following conditions are equivalent:*

- (i) *$\mathcal{A}$ is a subdiagonal subalgebra of $\mathcal{M}$.*
- (ii) *For all $0 < p \leq \infty$, $\mathcal{A}$ has L^p -factorization.*
- (iii) *For some $1 \leq p \leq \infty$, $\mathcal{A}$ has L^p -factorization.*

Proof. (i) $\Rightarrow$ (ii) follows by Theorem 3.1 of [2] (also see Theorem 4.2.1 of [1], Corollary 4.13 of [5] and the remark following Theorem 8.1 of [15]).

(ii) $\Rightarrow$ (iii) is clear.

(iii) $\Rightarrow$ (i). When $p = \infty$, it is (b) of Theorem 1.1 of [4]. From the proof of Proposition 2.3, we know that $\mathcal{A}_p$ is a subdiagonal subalgebra of $\mathcal{M}$. We claim that $\mathcal{A}_p = \mathcal{A}$. Indeed, it suffices to show that if $y \in L^1(\mathcal{M})$ and $\tau(ya) = 0$ for each $a \in \mathcal{A}$, then $\tau(ya) = 0$ for each $a \in \mathcal{A}_p$. Let $y = v|y|$ be the polar decomposition of y , and $v_1 \in \mathcal{M}$ such that $|y| = v_1(|y| + I)$. Notice that $\mathcal{A}_p$ is a subdiagonal subalgebra and $|y| + I \in L^1(\mathcal{M})$, $(|y| + I)^{-1} \in \mathcal{M}$. Hence, there is a unitary $u \in \mathcal{M}$ and $b \in [\mathcal{A}_p]_1$ such that $|y| + I = ub$, $b^{-1} \in \mathcal{A}_p$. So $y = wb$, where $w = vv_1u \in \mathcal{M}$. On the other hand,

$$[\mathcal{A}]_1 = [b^{-1}b\mathcal{A}]_1 \subset [b^{-1}[\mathcal{A}_p]_1]_1 \subset [\mathcal{A}_p]_1 = [\mathcal{A}]_1.$$

Since $\tau(wba) = \tau(ya) = 0$ for each $a \in \mathcal{A}$, it follows that $\tau(wa) = 0$ for each $a \in [\mathcal{A}_p]_1$. By τ -maximality of $\mathcal{A}_p$ we obtain $w \in (\mathcal{A}_p)_0$. Hence $y \in [(\mathcal{A}_p)_0]_1$, or $\tau(ya) = 0$ for each $a \in \mathcal{A}_p$. $\square$

Theorem 2.5. *Let $\mathcal{A}$ be a τ -maximal tracial subalgebra of $\mathcal{M}$. Then the following conditions are equivalent:*

- (i) *$\mathcal{A}$ is a subdiagonal subalgebra of $\mathcal{M}$.*
- (ii) *For some $0 < p < 1$, $\mathcal{A}$ has L^p -factorization and $\mathcal{A}_p$ is a tracial subalgebra of $\mathcal{M}$.*

Proof. We need only prove (ii) $\Rightarrow$ (i). It is clear that $\mathcal{A}_p$ is a subdiagonal subalgebra of $\mathcal{M}$ and $\mathcal{A} \subset \mathcal{A}_p$. If $b \in \mathcal{A}_p$, then $\tau(ba) = \tau(\mathcal{E}(ba)) = \tau(\mathcal{E}(b)\mathcal{E}(a)) = 0$ for each $a \in \mathcal{A}_0$. By τ -maximality of $\mathcal{A}$ we have $b \in \mathcal{A}$. Thus $\mathcal{A} = \mathcal{A}_p$. $\square$

Proposition 2.6. *Let $0 < p < 1$, and let $\mathcal{A}$ be a tracial subalgebra of $\mathcal{M}$. If $\mathcal{A}$ has L^{p_0} -factorization, and $\mathcal{A}_{p_0}$ is a tracial subalgebra of $\mathcal{M}$ for some $0 < p_0 < 1$, then*

$$(2.1) \quad \|\mathcal{E}(a)\|_p \leq \|a\|_p \quad \forall a \in \mathcal{A}.$$

Consequently, $\mathcal{E}$ extends to a contractive projection from $[\mathcal{A}]_p$ onto $L^p(\mathcal{D})$. The extension will still be denoted by $\mathcal{E}$.

Proof. Since $\mathcal{A}_{p_0}$ is a subdiagonal subalgebra of $\mathcal{M}$, Theorem 2.1 of [2] implies that

$$\|\mathcal{E}(a)\|_p \leq \|a\|_p \quad \forall a \in \mathcal{A}_{p_0}.$$

From this follows (2.1). $\square$

Corollary 2.7. *With the same assumption as in Proposition 2.6, we have that $\mathcal{E}(ab) = \mathcal{E}(a)\mathcal{E}(b)$ for $a \in [\mathcal{A}]_p$ and $b \in [\mathcal{A}]_q$ for $0 < p, q \leq \infty$.*

Proof. Note that $ab \in [\mathcal{A}]_r$ for any $a \in [\mathcal{A}]_p$ and $b \in [\mathcal{A}]_q$, where $1/r = 1/p + 1/q$. Thus $\mathcal{E}(ab)$ is well defined. Then the corollary follows immediately from the multiplicativity of $\mathcal{E}$ on $\mathcal{A}$ and (2.1). $\square$

3. L^2 -DENSITY

Definition 3.1. We say that a tracial subalgebra $\mathcal{A}$ of $\mathcal{M}$ satisfies L^2 -density if $\mathcal{A} + J(\mathcal{A})$ is dense in $L^2(\mathcal{M})$ in the usual Hilbert space norm on that space.

It is clear that $\mathcal{A} = \mathcal{A}_0 + \mathcal{D}$, $\mathcal{A} + J(\mathcal{A}) = \mathcal{A}_0 + \mathcal{D} + J(\mathcal{A}_0)$, $[\mathcal{A}]_2 = [\mathcal{A}_0]_2 + [\mathcal{D}]_2$, and

$$[\mathcal{A} + J(\mathcal{A})]_2 = [\mathcal{A}_0]_2 + [\mathcal{D}]_2 + [J(\mathcal{A}_0)]_2.$$

Hence, if $\mathcal{A}$ is an L^2 -dense tracial subalgebra of $\mathcal{M}$, then we have

$$(3.1) \quad L^2(\mathcal{M}) = [\mathcal{A}_0]_2 \oplus L^2(\mathcal{D}) \oplus [J(\mathcal{A}_0)]_2.$$

Lemma 3.2. *Let $\mathcal{A}$ be a τ -maximal tracial subalgebra of $\mathcal{M}$ that satisfies L^2 -density. Then we have*

- (i) $\mathcal{A} = ([J(\mathcal{A}_0)]_1)^\perp$, $\mathcal{A}_0 = ([J(\mathcal{A})]_1)^\perp$;
- (ii) $[\mathcal{A}]_1 = {}^\perp J(\mathcal{A}_0)$, $[\mathcal{A}_0]_1 = {}^\perp J(\mathcal{A})$;
- (iii) $[\mathcal{A}]_2 = L^2(\mathcal{M}) \cap [\mathcal{A}]_1$, $[\mathcal{A}_0]_2 = L^2(\mathcal{M}) \cap [\mathcal{A}_0]_1$.

Proof. (i) Assume $x \in \mathcal{A}$. If $y^* \in [J(\mathcal{A}_0)]_1$, then there exist $\{a_n\} \subset \mathcal{A}_0$ such that $a_n \rightarrow y$ in $L^1(\mathcal{M})$. Hence, $\tau(xy) = \lim_{n \rightarrow \infty} \tau(xa_n) = 0$, so $\mathcal{A} \subset ([J(\mathcal{A}_0)]_1)^\perp$. Conversely, we take $x \in \mathcal{M}$, $x \in ([J(\mathcal{A}_0)]_1)^\perp$; i.e., $\tau(xa) = 0$ for all a in $\mathcal{A}_0$. Since $\mathcal{A}$ is τ -maximal, $x \in \mathcal{A}$. Similarly, we have $\mathcal{A}_0 = ([J(\mathcal{A})]_1)^\perp$.

(ii) It is clear that $[\mathcal{A}]_1 \subset {}^\perp J(\mathcal{A}_0)$. Let $x \in L^1(\mathcal{M})$ and $x \in {}^\perp J(\mathcal{A}_0)$. Assume $x \notin [\mathcal{A}]_1$. Then there exists a $y \in \mathcal{M}$ such that $\tau(y^*x) = 1$ and $y \in ([\mathcal{A}]_1)^\perp$. By (i) we obtain that $y \in J(\mathcal{A}_0)$. Hence, $\tau(y^*x) = 0$, and this is a contradiction. Similarly we can prove $[\mathcal{A}_0]_1 = {}^\perp J(\mathcal{A})$.

(iii) It is obvious that $[\mathcal{A}]_2 \subseteq L^2(\mathcal{M}) \cap [\mathcal{A}]_1$. To prove the converse inclusion, let $a \in L^2(\mathcal{M}) \cap [\mathcal{A}]_1$. Then $a \perp J(\mathcal{A}_0)$. From this it follows that $a \perp [J(\mathcal{A})]_2$. By (3.1), $a \in [\mathcal{A}]_2$. $[\mathcal{A}_0]_2 = L^2(\mathcal{M}) \cap [\mathcal{A}_0]_1$ follows from the continuity of $\mathcal{E}$ on $[\mathcal{A}]_2$. $\square$

We say that $\xi \in L^2(\mathcal{M})$ is a right wandering vector if $\tau(\xi^*\xi b) = 0$ for all $b \in \mathcal{A}_0$.

Lemma 3.3. *Let $\mathcal{A}$ be a τ -maximal tracial subalgebra of $\mathcal{M}$ that satisfies L^2 -density. If $|\xi| \in L^2(\mathcal{D})$ for all right wandering vectors ξ , then $\mathcal{A}$ has L^2 -factorization.*

The proof of Lemma 3.3 is the same as that of Theorem 3.1 of [2]. We omit the details and refer the reader to [2].

Theorem 3.4. *Let $\mathcal{A}$ be a τ -maximal tracial subalgebra of $\mathcal{M}$ that satisfies L^2 -density. Then the following conditions are equivalent:*

- (i) *If $x \in [\mathcal{A}]_1^+$, then $x^{\frac{1}{2}} \in H^2(\mathcal{A})$.*
- (ii) *If ξ is right wandering vector, then $|\xi| \in L^2(\mathcal{D})$.*
- (iii) *$\mathcal{A}$ is a subdiagonal subalgebra of $\mathcal{M}$.*

Proof. (ii) $\Rightarrow$ (iii) follows from Lemma 3.3 and Theorem 2.4.

(iii) $\Rightarrow$ (i) is trivial.

(i) $\Rightarrow$ (ii). Let ξ be a right wandering vector. Then

$$\tau(\xi^* \xi b) = 0, \forall b \in \mathcal{A}_0.$$

By (ii) of Lemma 3.2 we have $\xi^* \xi \in [\mathcal{A}]_1$, and $|\xi| \in [\mathcal{A}]_2$. Hence,

$$\tau(|\xi|b) = 0, \forall b \in \mathcal{A}_0 + J(\mathcal{A}_0).$$

So by the L^2 -density of $\mathcal{A}$, it follows that $|\xi| \in L^2(\mathcal{D})$. $\square$

Theorem 3.5. *Let $\mathcal{A}$ be a tracial subalgebra of $\mathcal{M}$ that satisfies L^2 -density. Then the following conditions are equivalent:*

- (i) *$\mathcal{A}$ is a subdiagonal subalgebra of $\mathcal{M}$.*
- (ii) *For some $0 < p < 1$, $\mathcal{A}$ has L^p -factorization and $\mathcal{A}_p$ is a tracial subalgebra of $\mathcal{M}$.*

Proof. (ii) $\Rightarrow$ (i). Since $\mathcal{A}_p$ is a subdiagonal subalgebra of $\mathcal{M}$,

$$L^2(\mathcal{M}) = [\mathcal{A}_p]_2 \oplus [J((\mathcal{A}_p)_0)]_2.$$

So from

$$L^2(\mathcal{M}) = [\mathcal{A}]_2 \oplus [J(\mathcal{A}_0)]_2, [\mathcal{A}]_2 \subset [\mathcal{A}_p]_2, [J(\mathcal{A}_0)]_2 \subset [J((\mathcal{A}_p)_0)]_2,$$

it follows that $[\mathcal{A}]_2 = [\mathcal{A}_p]_2$. If $x \in L^2(\mathcal{M})^{-1}$, then there is a unitary $u \in \mathcal{M}$ and $a \in [\mathcal{A}_p]_2^{-1} = [\mathcal{A}]_2^{-1}$ such that $x = ua$, since $\mathcal{A}_p$ is a subdiagonal subalgebra. Hence, $\mathcal{A}$ has L^2 -factorization. By Theorem 2.4 we obtain the desired result. $\square$

REFERENCES

- [1] W. B. Arveson, Analyticity in operator algebras, *Amer. J. Math.* **89** (1967), 578-642. MR0223899 (36:6946)
- [2] T. N. Bekjan, Q. Xu, Riesz and Szegő type factorizations for noncommutative Hardy spaces, *J. Operator Theory* **62** (2009), 215-231. MR2520548
- [3] D. P. Blecher, L. E. Labuschagne, Logmodularity and isometries of operator algebras, *Trans. Amer. Math. Soc.* **355** (2003), 1621-1646. MR1946408 (2004c:46113)
- [4] D. P. Blecher, L. E. Labuschagne, Characterization of noncommutative $\mathcal{H}^\infty$, *Integr. Equ. Oper. Theory* **56** (2006), 301-321. MR2270840 (2007m:46101)
- [5] D. P. Blecher, L. E. Labuschagne, Applications of Fuglede-Kadison determinant: Szegő's theorem and outers for noncommutative H^p , *Trans. Amer. Math. Soc.* **360** (2008), 6131-6147. MR2425707 (2009m:46096)
- [6] R. Exel, Maximal subdiagonal algebras, *Amer. J. Math.* **110** (1988), 775-782. MR955297 (90b:46114)
- [7] T. Fack, H. Kosaki, Generalized s -numbers of τ -measurable operators, *Pac. J. Math.* **123** (1986), 269-300. MR840845 (87h:46122)
- [8] G. Ji, K.-S. Saito, Factorization in subdiagonal algebras, *J. Funct. Anal.* **159** (1998), 191-202. MR1654186 (99k:46106)

- [9] S. Kawamura, J. Tomiyana, On subdiagonal algebras associated with flows in operator algebras, *J. Math. Soc. Japan* **29** (1977), 73-90. MR0454650 (56:12899)
- [10] L. E. Labuschagne, A noncommutative Szegő theorem for subdiagonal subalgebras of von Neumann algebras. *Proc. Amer. Math. Soc.* **133** (2005), 3643-3646. MR2163602 (2006d:46081)
- [11] M. Marsalli, Noncommutative H^2 -spaces, *Proc. Amer. Math. Soc.* **125** (1997), 779-784. MR1350954 (97e:46089)
- [12] M. Marsalli, G. West, Noncommutative H^p -spaces, *J. Operator Theory* **40** (1997), 339-355. MR1660390 (2000b:46117)
- [13] M. Marsalli, G. West, The dual of noncommutative H^1 , *Indiana Univ. Math. J.* **47** (1998), 489-500. MR1647920 (99h:46122)
- [14] M. McAsey, P. S. Muhly, K.-S. Saito, Nonselfadjoint crossed products (invariant subspaces and maximality), *Trans. Amer. Math. Soc.* **248** (1979), 381-409. MR522266 (80j:46101b)
- [15] G. Pisier, Q. Xu, Non-commutative L^p -spaces, in *Handbook of the geometry of Banach spaces*, Vol. 2, North-Holland, Amsterdam, 2003, 1459-1517. MR1999201 (2004i:46095)
- [16] N. Randrianantoanina, Hilbert transform associated with finite maximal subdiagonal algebras, *J. Austral. Math. Soc. (Series A)* **65** (1999), 388-404. MR1660424 (2000a:46109)
- [17] K.-S. Saito, On non-commutative Hardy spaces associated with flows on finite von Neumann algebras, *Tohoku Math. J.* **29** (1977), 585-595. MR0463932 (57:3870)
- [18] K.-S. Saito, A note on invariant subspaces for maximal subdiagonal algebras, *Proc. Amer. Math. Soc.* **77** (1979), 349-352. MR545594 (81b:46078)
- [19] M. Takesaki, Duality for crossed products and the structure of von Neumann algebras of type III, *Acta Math.* **131** (1973), 249-310. MR0438149 (55:11068)
- [20] M. Terp, L^p -spaces associated with an arbitrary von Neumann algebra, Notes, Math. Institute, Copenhagen Univ., 1981.

COLLEGE OF MATHEMATICS AND SYSTEMS SCIENCES, XINJIANG UNIVERSITY, URUMQI 830046,
PEOPLE'S REPUBLIC OF CHINA