

## RELATIVELY POINTWISE RECURRENT GRAPH MAP

HATTAB HAWETE

(Communicated by Yingfei Yi)

ABSTRACT. Let  $f$  be a self-continuous map of a graph  $G$ . Let  $P(f)$  and  $R(f)$  denote the sets of periodic points and recurrent points respectively. We say that the map  $f$  is *relatively recurrent* if  $\overline{R(f)} = G$ . In this paper, it is shown that  $f$  is relatively recurrent if and only if one of the following statements holds:

- (a)  $G$  is a circle and  $f$  is a homeomorphism topologically conjugate to an irrational rotation of the unit circle  $\mathbb{S}^1$ ;
  - (b)  $\overline{P(f)} = G$ .
- Part (b) extends a result of Blokh.

### 1. INTRODUCTION

A *topological dynamical system* is a pair  $(X, f)$ , where  $X$  is a compact metric space and  $f$  is a continuous map from  $X$  to itself. Let  $\mathbb{N}$  be the set of positive integers. Let  $f^0$  be the identity map of  $X$ . Define, inductively,  $f^n = f \circ f^{n-1}$  for any non-zero positive integer  $n$ . For  $x \in X$ ,  $\{f^n(x) : n \in \mathbb{N}\}$  is called the *orbit* of  $x$  and is denoted by  $O(x, f)$ . Here  $x$  is *periodic* if  $f^n(x) = x$  for some non-zero positive integer  $n$ . Also,  $x$  is called a *recurrent point* of  $f$  if for any neighborhood  $U$  of  $x$  and any  $m \in \mathbb{N}$  there exists  $n > m$  such that  $f^n(x) \in U$ . It is easy to see that if  $x$  is recurrent, then every iterate of  $x$  is also recurrent. The converse is false. Here  $x$  is called an *almost periodic point* of  $f$  if for any neighborhood  $U$  of  $x$  there exists  $N \in \mathbb{N}$  such that  $\{f^{n+i}(x) : i = 0, 1, \dots, N\} \cap U \neq \emptyset$  for all  $n \in \mathbb{N}$ . Also,  $x$  is a *non-wandering point* of  $f$  provided that for any open set  $U$  containing  $x$  there exist  $y \in U$  and  $n \in \mathbb{N}$  such that  $f^n(y) \in U$ . Let  $P(f)$ ,  $AP(f)$ ,  $R(f)$  and  $\Omega(f)$  denote the set of periodic points, almost periodic points, recurrent points and non-wandering points respectively. Notice that  $\Omega(f)$  is closed and for the general topological system  $(X, f)$  there are no further relations except for  $P(f) \subset AP(f) \subset R(f) \subset \Omega(f)$ . But for one-dimensional systems one can say more. For a dendrite map Illanes [8] proved that  $\overline{P(f)} = \overline{R(f)}$  if and only if the dendrite does not contain any copy of the Gehman dendrite. For a graph map Mai and Shao [9] showed that  $\overline{R(f)} = \overline{P(f)} \cup R(f)$ . In Lemma 3.1, for a graph map, we will show that  $\overline{AP(f)} = \overline{R(f)}$ .

---

Received by the editors April 10, 2010 and, in revised form, June 2, 2010.

2010 *Mathematics Subject Classification*. Primary 37B20, 37E25.

*Key words and phrases*. Periodic point, recurrent point, almost periodic point, minimal set, relatively pointwise recurrent map, graph map.

©2010 American Mathematical Society  
 Reverts to public domain 28 years from publication

It is interesting to study maps such that  $P(f)$  or  $R(f)$  satisfies some additional properties. Montgomery [12], Weaver [15], Epstein [4], and others studied homeomorphisms such that  $P(f)$  is the whole space  $X$  ( $X$  is a connected manifold or a continuum embedded in a 2-manifold). For an interval map, Nitecki [13] showed that if  $P(f)$  is closed, then  $P(f) = \Omega(f)$ . For a graph map Mai [11] proved that  $R(f)$  is the whole space  $X$  if and only if one of the following statements holds:

- (1)  $X$  is a circle, and  $f$  is a homeomorphism topologically conjugate to an irrational rotation of the unit circle;
- (2)  $f$  is a periodic homeomorphism.

In this paper we will study a relatively pointwise recurrent graph map. Our main result is the following theorem:

**Main Theorem.** *Let  $G$  be a graph and let  $f : G \rightarrow G$  be a continuous map. Then  $f$  is relatively pointwise recurrent if and only if one of the following two statements holds:*

- (1)  $G$  is a circle, and  $f$  is a homeomorphism topologically conjugate to an irrational rotation of the unit circle  $\mathbb{S}^1$ ;
- (2)  $\overline{P(f)} = G$ .

A set  $W \subset X$  is called a *minimal set* of  $f$  if it is nonempty, closed, invariant ( $f(W) \subset W$ ) and no proper subset of  $W$  satisfies these three properties, which is equivalent to the fact that the orbit of every element of  $W$  is dense.

In a topological dynamical system there are two well-known theorems which exhibit the close relationship between almost periodic points and minimal sets; see Birkhoff [2]. In fact, if  $X$  is a locally compact metric space, then the set of almost periodic points  $AP(f)$  is the union of all minimal sets of  $f$ .

If  $X$  has no isolated point, then  $f$  is a *transitive map* if it has a dense orbit. If every orbit of  $f$  is dense in  $X$ , the map  $f$  is called *minimal*. For a transitive graph map Blokh [3] proved that  $P(f)$  is dense. In this paper we show that for a graph map if  $R(f)$  is dense, then  $P(f)$  is also dense, which extends this result of Blokh [3] (see Corollary 3.4).

## 2. RELATIVELY POINTWISE RECURRENT MAP

A map  $f$  of a compact metric space  $(X, d)$  is *recurrent* if it admits iterates arbitrarily close to the identity, i.e. if there exists a sequence  $n_k \rightarrow +\infty$  such that

$$d(f^{n_k}, Id) \rightarrow 0 \text{ as } k \rightarrow +\infty.$$

We say that  $f$  is *pointwise recurrent* if  $R(f) = X$ .

We continue, motivated by a desire to understand the mechanics of recurrent maps, by a desire to extend some known result and by the following:

In [6] Gottschalk proved that if  $X$  is a compact connected metric space,  $f$  is a homeomorphism, and if  $\overline{R(f)} = X$ , then every recurrent cut point of  $X$  is periodic.

We start by the following definition.

**Definition.**  $f$  is called a *relatively pointwise recurrent map* if  $\overline{R(f)} = X$ .

For example a transitive map is a relatively pointwise recurrent map.

Let  $X$  be a closed domain of finite volume of the  $n$ -Euclidian space  $\mathbb{R}^n$  or the  $n$ -torus  $\mathbb{T}^n$  and let  $f$  be an invertible volume-preserving self-map of  $X$ . Then  $f$  is a relatively pointwise recurrent map [7, Theorem 6.1.9].

We have the following implications:

$$\text{recurrent} \Rightarrow^1 \text{pointwise recurrent} \Rightarrow^2 \text{relatively pointwise recurrent}.$$

The following examples show that none of these implications can be reversed.

**Examples 2.1.** (1) An irrational rotation of  $\mathbb{R}^2$  is a pointwise recurrent non-recurrent map.

(2) In  $\mathbb{R}^2$  we consider the points  $A_n(0, \frac{1}{n})$  and  $B(n, \frac{1}{n})$  for every integer  $n > 0$ . Put  $X = (\bigcup_{n>0} [A_n, B_n]) \cup [0, +\infty[ \times \{0\}$ . Define the map  $f$  on  $X$  by:

- $f(x, \frac{1}{n}) = (\varphi(x), \frac{1}{n})$ , where

$$\varphi(x) = \begin{cases} x + 1 & \text{if } x \leq n - 1, \\ x + 1 - n & \text{if } n - 1 < x \leq n. \end{cases}$$

- $f(x, 0) = (x + 1, 0)$ .

$f$  is a relatively pointwise recurrent non-pointwise recurrent map.

In this example one can choose  $X$  to be connected.

**Theorem 2.2.** *If a continuous map of a topological space  $X$  to itself is either (1) recurrent, (2) pointwise recurrent, or (3) relatively pointwise recurrent, then so is  $f^k$ , for each integer  $k$ .*

*Proof.* (1) and (2) follow from [14] and [6] respectively.

We have  $R(f) \subset R(f^k)$ , and so if  $\overline{R(f)} = X$ , then  $\overline{R(f^k)} = X$ , which implies (3).  $\square$

**Proposition 2.3.** *If  $f$  is relatively pointwise recurrent, then  $f$  is surjective.*

*Proof.* If  $y \in R(f)$ , then there exists a point  $x \in X$  such that  $f(x) = y$ .

Let  $y$  be an element of  $X - R(f)$ . Then there exists a sequence  $y_n$  of  $R(f)$  which converges to  $y$ . For all  $n$  there exists  $x_n$  such that  $f(x_n) = y_n$ . Since  $X$  is compact, the sequence  $x_n$  has a limit point  $x$  and so  $f(x) = y$ . Therefore  $f$  is surjective.  $\square$

**Proposition 2.4.** *If  $f$  is relatively pointwise recurrent, then  $\Omega(f) = X$ .*

*Proof.* Let  $x$  be an element of  $X$ . If  $x$  is a recurrent point, it is also non-wandering. If  $x$  is a non-recurrent point, then every neighborhood of  $x$  contains a recurrent point, and so it is in  $\Omega(f)$ .  $\square$

The referee noticed that the converse of Proposition 2.4 also holds; see, for example, [5, Theorem 1.27].

**Proposition 2.5.** *If  $f$  is a relatively pointwise recurrent non-recurrent map, then  $f$  is not equicontinuous.*

*Proof.* Since  $f$  is a relatively pointwise recurrent non-recurrent map, then  $R(f)$  is not closed and so is not equal to  $\Omega(f)$ . From [10, Proposition 2.1] it follows that  $f$  is not equicontinuous.  $\square$

The following proposition can be derived from [10, Proposition 2.1].

**Proposition 2.6.** *If  $f$  is an equicontinuous relatively pointwise recurrent map, then  $f$  is a pointwise recurrent map.*

## 3. PROOF OF MAIN THEOREM

Before going into the proof of the main theorem we recall the definition of a graph. A (finite) graph  $G$  is a compact connected Hausdorff space which contains a finite non-empty set  $V$  (the set of *vertices*), such that every connected component of  $G \setminus V$  is homeomorphic to an open interval of the real line. These connected components are called *edges*. Since any graph can be embedded in  $\mathbb{R}^3$ , in what follows we will consider each graph endowed with the topology induced by the topology of  $\mathbb{R}^3$ . A *graph map* is a continuous map from a graph  $G$  to itself.

To prove the main theorem we need the following lemmas:

**Lemma 3.1.** *Let  $f$  be a graph map. Then  $\overline{R(f)} = \overline{AP(f)}$ .*

*Proof.* We recall that  $\overline{AP(f)}$  is the closure of the union of all minimal sets of  $f$  and  $\overline{AP(f)} \subset \overline{R(f)}$ .

In [1, Theorem 1] the authors showed that a minimal set of a graph map is a finite set or a Cantor set or a union of (finitely many) pairwise disjoint circles. One can deduce that each component of  $G - (\overline{AP(f)} \cup V)$  is an open arc of  $G$ .

Let  $]a, b[$  be a component of  $G - (\overline{AP(f)} \cup V)$  such that  $a \in \overline{AP(f)}$ . We suppose that  $]a, b[ \cap R(f) \neq \emptyset$ . Let  $x$  be an element of  $]a, b[ \cap R(f)$ . Since  $x$  is recurrent, there exists an increasing sequence  $(n_q)$  such that  $(f^{n_q}(x))$  converges to  $x$ . There exist three integers  $i < j < k$  such that one of the following two statements holds:

- (1)  $a < x < f^{n_k}(x) < f^{n_j}(x) < f^{n_i}(x)$ .
- (2)  $a < f^{n_i}(x) < f^{n_j}(x) < f^{n_k}(x) < x$ .

By applying [9, Lemma 2.2] we obtain:

- (1)  $\Rightarrow f^{n_k}(a) \in ]a, b[$ .
- (2)  $\Rightarrow f^{n_j - n_i}(a) \in ]a, b[$ .

In both cases the interval  $]a, b[$  will intersect  $\overline{AP(f)}$ , which is impossible.  $\square$

**Lemma 3.2.** *Let  $f$  be a relatively pointwise recurrent map of a graph  $G$ . If  $W$  is a proper minimal set of  $f$ , then  $W$  is not a union of (finitely many) pairwise disjoint circles.*

*Proof.* In [1, Theorem 1] the authors showed that a minimal set of a graph map is a finite set or a Cantor set or a union of (finitely many) pairwise disjoint circles.

Suppose that  $W$  is a union of (finitely many) pairwise disjoint circles. Then  $G$  is not a circle (because  $W \neq G$ ). From the fact that  $G$  is connected it follows that  $W$  contains a branching point  $w$ . Let  $A$  be an arc of  $G$  such that  $A \cap W = \{w\}$ . Since  $\overline{R(f)} = G$ , there exists in  $A$  a sequence  $(w_n)$  of recurrent points which converges to  $w$ . From the fact that  $O(w, f)$  is dense in  $W$  it follows that there exists an integer  $p$  such that  $f^p(w)$  is not a branching point and so by continuity of  $f^p$ , there exists an integer  $N$  such that  $f^p(w_n) \in W$  for all  $n > N$ . The recurrence of  $w_n$  implies that there exists an integer  $q > p$  such that  $f^q(w_n) \in A - \{w\}$ , which implies that  $f^{q-p}(f^p(w_n)) \in A - \{w\}$ , which contradicts the fact that  $W$  is invariant.  $\square$

**Lemma 3.3.** *Let  $f$  be a relatively pointwise recurrent map of a graph  $G$ . If  $f$  is not a minimal map, then  $\overline{P(f)} = \overline{AP(f)}$ .*

*Proof.* We always have the inclusion  $\overline{P(f)} \subset \overline{AP(f)}$ .

By applying [1, Theorem 1] and Lemma 3.2 it follows that every minimal set of  $f$  is a periodic orbit or a Cantor set. Let  $W$  be a minimal set which is a

Cantor set. Let  $w$  be an element of  $W$  and let  $w'$  be an element of  $G - V(G)$  such that the open arc  $(w, w')$  does not meet  $W$ . If  $(w, w') \cap P(f) = \emptyset$ , then from the fact that  $(w, w') \cap R(f) \neq \emptyset$  and by applying [9, Lemma 2.2] it follows that  $(w, w') \cap O(w, f) \neq \emptyset$ , which is impossible. Thus  $w \in \overline{P(f)}$ . Since  $P(f)$  is invariant,  $O(w, f) \subset \overline{P(f)}$  and so  $W \subset \overline{P(f)}$ . Therefore  $\overline{AP(f)} = \overline{P(f)}$ .  $\square$

*Proof of the main theorem.* The two statements imply that  $f$  is a relatively pointwise recurrent map.

Conversely, (1) if  $f$  is a minimal map, then first by [11, Theorem 3.2] it follows that  $G$  is a circle, and second by [11, Lemma 3.1] it follows that  $f$  is a homeomorphism. From Proposition 2.4 it follows that  $f$  is a pointwise non-wandering circle map without periodic points. Thus it is topologically conjugate to an irrational rotation.

(2) If  $f$  is not a minimal map, then from Lemma 3.1 and Lemma 3.3 it follows that  $\overline{P(f)} = G$ .  $\square$

The fact that the set of periodic points of a transitive graph map is dense was proved by Blokh in [3]. The following corollary extends this result.

**Corollary 3.4.** *Let  $f$  be a relatively pointwise recurrent map of a graph  $G$ . If  $f$  is not a minimal map, then  $\overline{P(f)} = G$ .*

#### ACKNOWLEDGEMENT

The author gratefully acknowledges the helpful corrections, comments and suggestions from the referee.

#### REFERENCES

1. F. Balibrea, R. Hric and L. Snoha, *Minimal sets on graphs and dendrites*, Internat. J. Bifur. Chaos Appl. Sci. Engrg. **13** (2003), 1721-1725. MR2015622 (2004i:37083)
2. G.D. Birkhoff, *Dynamical Systems*, revised edition, Amer. Math. Soc. Colloq. Publ., vol. **IX**, Amer. Math. Soc., Providence, RI, 1966. MR0209095 (35:1)
3. A. Blokh, *On the connection between entropy and transitivity for one-dimensional mappings*, Russian Math. Surveys **42:5** (1987), 165-166. MR928783 (89g:58117)
4. D.B.A. Epstein, *Pointwise periodic homeomorphisms*, Proc. London Math. Soc. (3) **42** (1981), 415-460. MR614729 (83e:57011)
5. H. Furstenberg, *Recurrence in ergodic theory and combinatorial number theory*, M. B. Porter Lectures. Princeton University Press, Princeton, 1981. MR603625 (82j:28010)
6. W.H. Gottschalk, *Powers of homeomorphisms with almost periodic properties*, Bull. Amer. Math. Soc. **50** (1944), 222-227. MR0009850 (5:213c)
7. B. Hasselblatt and A. Katok, *A First Course in Dynamics: With a Panorama of Recent Developments*, Cambridge University Press, 2003. MR1995704 (2004f:37001)
8. A. Illanes, *A characterization of dendrites with periodic-recurrent property*, Topology Proc. **23** (1998), 221-235. MR1803250 (2001j:54037)
9. J.H. Mai and S. Shao,  $\overline{R} = \overline{P} \cup R$  for graph maps, J. Math. Anal. Appl. **350** (2009), 9-11. MR2476886 (2010a:37082)
10. J.H. Mai, *The structure of equicontinuous maps*, Trans. Amer. Math. Soc. **355** (2003), 4125-4136. MR1990578 (2004e:54021)
11. J.H. Mai, *Pointwise-recurrent graph maps*, Ergod. Th. Dynam. Sys. **25** (2005), 629-637. MR2129113 (2005m:37094)
12. D. Montgomery, *Pointwise periodic homeomorphisms*, Amer. J. Math. **59** (1937), 118-120. MR1507223

13. Z. Nitecki, *Maps of the interval with closed periodic set*, Proc. Amer. Math. Soc. **85** (1982), 451-456. MR656122 (83k:58067)
14. L.G. Oversteegen and E.D. Tymchatyn, *Recurrent homeomorphisms on  $\mathbf{R}^2$  are periodic*, Proc. Amer. Math. Soc. **110** (1990), 1083-1088. MR1037216 (91k:54070)
15. N. Weaver, *Pointwise periodic homeomorphisms of continua*, Ann. of Math. (2) **95** (1972), 83-85. MR0293600 (45:2677)

INSTITUT SUPÉRIEUR D'INFORMATIQUE ET DU MULTIMEDIA, ROUTE DE TUNIS, KM 10, B.P.  
242, SFAX 3021, TUNISIA

*E-mail address:* `hattab.hawete@yahoo.fr`