SKEW CATEGORIES, SMASH PRODUCT CATEGORIES AND QUASI-KOSZUL CATEGORIES

DEKE ZHAO

(Communicated by Martin Lorenz)

ABSTRACT. Let $\mathcal{A}$ be a small additive Krull-Schmidt locally radical finite category over a field K and let G be a finite group. We show that if $\mathcal{A}$ is a free G -category (resp. G -graded category), then $\mathcal{A}$ is quasi-Koszul if and only if the skew (resp. smash product) category $G * \mathcal{A}$ (resp. $\mathcal{A} \# G$) is.

1. INTRODUCTION

Koszul algebras were originally defined by Priddy in [9] and have arisen in many contexts: algebraic geometry, Lie theory, combinatorics, non-commutative geometry and topology. There exist numerous generalizing notations of Koszul algebras. For example, Berger [2] introduced the notation of N -Koszul algebras, and Cassidy and Shelton [3] introduced the notation of K_2 algebras. Very recently, Martínez-Villa and Solberg [7] introduced the notation of (weakly, quasi-)Koszul categories to obtain a naturally associated Koszul theory for any finite dimensional algebra.

Throughout this paper we will consider small additive categories $\mathcal{A}$ over a field K , free G -categories and G -graded categories where G is a finite group. For G -categories and G -graded categories, Cibils and Marcos [4] defined their skew categories and smash products categories, respectively, which provide a generalized category version of the Cohen-Montgomery Duality Theorem [5]. Inspired by the closeness of the (quasi-)Koszulity of algebras under the skew products and smash products [6, 10], we show that

Theorem. *Let $\mathcal{A}$ be a small additive Krull-Schmidt locally radical finite category over a field K and let G be a finite group.*

- (i) *If $\mathcal{A}$ is a free G -category, then $\mathcal{A}$ is quasi-Koszul if and only if the skew category $G * \mathcal{A}$ is.*
- (ii) *If $\mathcal{A}$ is G -graded, then $\mathcal{A}$ is quasi-Koszul if and only if $\mathcal{A}$ is.*

In Section 2 we recall the definitions of G -category, of skew category, and of G -graded category and give some basic facts. In Section 3 we recall the definitions of the syzygy of functor and of quasi-Koszul category and prove the Theorem.

Received by the editors November 22, 2009 and, in revised form, July 12, 2010.

2010 *Mathematics Subject Classification.* Primary 18A25, 18G10; Secondary 18B99.

Key words and phrases. Skew categories, smash product categories, quasi-Koszul categories, group-graded categories.

2. SKEW CATEGORIES AND SMASH PRODUCT CATEGORIES

2.1. Skew categories. A G -category is a K -category $\mathcal{A}$ with firstly a set action of G on the set of objects $\mathcal{A}_0$ and secondly K -module maps $g : \text{Hom}_{\mathcal{A}}(X, Y) \rightarrow \text{Hom}_{\mathcal{A}}(gX, gY)$ for each $g \in G$ and $(X, Y) \in \mathcal{A}_0 \times \mathcal{A}_0$ satisfying $g(\varphi\psi) = (g\varphi)(g\psi)$ in case the morphism $\varphi\psi$ is composed in $\mathcal{A}$. Moreover for $g, h \in G$ and morphism ψ we have $(gh)\psi = g(h\psi)$ and $e\psi = \psi$ where e is the neutral element of G . In other words there is a group homomorphism from G to the group of autofunctors of $\mathcal{A}$. If the action of G is free on $\mathcal{A}_0$, we say that $\mathcal{A}$ is a free G -category.

Definition 1 ([4, Definition 2.3]). Let $\mathcal{A}$ be a G -category. The *skew category* $G * \mathcal{A}$ is a K -category with $(G * \mathcal{A})_0 = \mathcal{A}_0$ and morphisms $\text{Hom}_{G * \mathcal{A}}(X, Y) = \bigoplus_{g \in G} \text{Hom}_{\mathcal{A}}(gX, Y)$. Composition of morphisms is provided by the composition of $\mathcal{A}$ after adjustment.

An additive K -category $\mathcal{A}$ is said to be *Krull-Schmidt* if any object in $\mathcal{A}$ is a finite direct sum of objects with a local endomorphism algebra. Recall that the radical of a category $\mathcal{A}$, $\text{rad}_{\mathcal{A}}$, is a sub-bifunctor of $\text{Hom}_{\mathcal{A}}(-, -)$ given by

$$\text{rad}_{\mathcal{A}}(X, Y) = \{\psi \in \text{Hom}_{\mathcal{A}}(X, Y) \mid \varphi\psi \in \text{rad}(\text{End}_{\mathcal{A}}(X)) \text{ for all } \varphi \in \text{Hom}_{\mathcal{A}}(Y, X)\}.$$

Observe that we also have (see [8, Lemmas 4.1 and 4.2])

$$\text{rad}_{\mathcal{A}}(X, Y) = \{\psi \in \text{Hom}_{\mathcal{A}}(X, Y) \mid \psi\varphi \in \text{rad}(\text{End}_{\mathcal{A}}(Y)) \text{ for all } \varphi \in \text{Hom}_{\mathcal{A}}(Y, X)\}.$$

We say that $\psi \in \text{Hom}_{\mathcal{A}}(X, Y)$ is in $\text{rad}_{\mathcal{A}}^2(X, Y)$ if ψ is a finite sum of the form $X \xrightarrow{\psi'_i} X'_i \xrightarrow{\psi''_i} Y$ with $\psi'_i \in \text{rad}_{\mathcal{A}}(X, X'_i)$, $\psi''_i \in \text{rad}_{\mathcal{A}}(X'_i, Y)$ for $i = 1, \dots, m$ and $\psi = \sum_{i=1}^m \psi''_i \psi'_i$. Inductively define $\text{rad}_{\mathcal{A}}^n := \text{rad}_{\mathcal{A}} \cdot \text{rad}_{\mathcal{A}}^{n-1}$. A K -category $\mathcal{A}$ is called *locally radical finite* if $\text{rad}_{\mathcal{A}}^i(X, Y) / \text{rad}_{\mathcal{A}}^{i+1}(X, Y)$ is finite dimensional over K for all $X, Y \in \mathcal{A}$ and $i \geq 0$.

Lemma. If $X, Y \in \mathcal{A}$, then $\text{rad}_{G * \mathcal{A}}(X, Y) \simeq KG \otimes_K \text{rad}_{\mathcal{A}}(X, Y)$ and $\text{rad}_{G * \mathcal{A}}^n \simeq \overbrace{KG \otimes_K \cdots \otimes_K KG}^n \otimes_K \text{rad}_{\mathcal{A}}^n, n \geq 1$.

Proof. Clearly $\text{rad}_{\mathcal{A}}(gX, Y) \simeq \text{rad}_{\mathcal{A}}(X, Y)$ for all $g \in G$. Assume that $\psi = \sum_{g \in G} \psi_g$ and $\varphi = \sum_{h \in G} \varphi_h$ where $\psi_g \in \text{Hom}_{\mathcal{A}}(gX, Y)$ and $\varphi_h \in \text{Hom}_{\mathcal{A}}(hY, X)$. Then $\varphi \cdot \psi \in \text{rad}(\text{End}_{G * \mathcal{A}}(X))$ if and only if $\varphi_h(h\psi_g) \in \text{rad}(\text{End}_{G * \mathcal{A}}(X))$ for all $g, h \in G$ if and only if $\psi_g \in \text{rad}_{\mathcal{A}}(gX, Y)$ for all $g \in G$. Thus we obtain that

$$\begin{aligned} \text{rad}_{G * \mathcal{A}}(X, Y) &= \{\psi \in \text{Hom}_{G * \mathcal{A}}(X, Y) \mid \varphi\psi \in \text{rad}(\text{End}_{G * \mathcal{A}}(X)) \\ &\quad \text{for all } \varphi \in \text{Hom}_{G * \mathcal{A}}(X, Y)\} \\ &= \{\psi = \sum_{g \in G} \psi_g, \psi_g \in \text{Hom}_{\mathcal{A}}(gX, Y) \mid \sum_{g, h \in G} \varphi_h \psi_g \in \text{rad}(\text{End}_{G * \mathcal{A}}(X)) \\ &\quad \text{for all } \varphi_h \in \text{Hom}_{\mathcal{A}}(hY, X), h \in G\} \\ &= \{\psi = \sum_{g \in G} \psi_g, \psi_g \in \text{Hom}_{\mathcal{A}}(gX, Y) \mid \varphi_h(h\psi_g) \in \text{rad}(\text{End}_{G * \mathcal{A}}(X)) \\ &\quad \text{for all } \varphi_h \in \text{Hom}_{\mathcal{A}}(hY, X), h \in G\} \\ &= \{\psi = \sum_{g \in G} \psi_g \mid \psi_g \in \text{rad}_{\mathcal{A}}(gX, Y)\} \\ &\simeq KG \otimes_K \text{rad}_{\mathcal{A}}(gX, Y) \\ &= KG \otimes_K \text{rad}_{\mathcal{A}}(X, Y). \end{aligned}$$

By induction, we have

$$\begin{aligned} \operatorname{rad}_{G*\mathcal{A}}^n(X, Y) &= \operatorname{rad}_{G*\mathcal{A}} \cdot \operatorname{rad}_{G*\mathcal{A}}^{n-1}(X, Y) \\ &\simeq \operatorname{rad}_{G*\mathcal{A}} \cdot \overbrace{(KG \otimes_K \cdots \otimes_K KG)}^{n-1} \otimes_K \operatorname{rad}_{\mathcal{A}}^{n-1}(X, Y) \\ &\simeq \overbrace{KG \otimes_K \cdots \otimes_K KG}^n \otimes_K \operatorname{rad}_{\mathcal{A}}^n(X, Y). \end{aligned}$$

□

2.2. Smash product categories. A G -graded category $\mathcal{B}$ is a K -category together with a decomposition of morphisms $\operatorname{Hom}_{\mathcal{B}}(X, Y) = \bigoplus_{g \in G} \operatorname{Hom}_{\mathcal{B}}(X, Y)_g$ such that $\operatorname{Hom}_{\mathcal{B}}(Y, Z)_g \operatorname{Hom}_{\mathcal{B}}(X, Y)_h \subset \operatorname{Hom}_{\mathcal{B}}(X, Z)_{gh}$. In particular, $\operatorname{Hom}_{\mathcal{B}}(X, X)$ is a G -graded algebra for each $X \in \mathcal{B}$.

Definition 2 ([4, Definition 3.1]). Let $\mathcal{B}$ be a G -graded category. The *smash product category* $\mathcal{B} \# G$ is a category with $(\mathcal{B} \# G)_0 = \mathcal{B}_0 \times G$ and morphisms $\operatorname{Hom}_{\mathcal{B} \# G}((X, g), (Y, h)) = \operatorname{Hom}_{\mathcal{B}}(X, Y)_{h^{-1}g}$. Composition of morphisms is provided by the graded composition of $\mathcal{B}$.

Define a G -action on $\mathcal{B} \# G$ by $g(X, h) = (X, gh)$ and $gf = f \in \operatorname{Hom}_{\mathcal{B} \# G}((X, gh), (Y, gl))$ if $f \in \operatorname{Hom}_{\mathcal{B} \# G}((X, h), (Y, l))$. Then the category $\mathcal{B} \# G$ is a free G -category and the categories $G * (\mathcal{B} \# G)$ and $\mathcal{B}$ are equivalent [4, Corollary 3.3]. By the Lemma, we have the following:

Corollary. If $X, Y \in \mathcal{B}$, then $KG \otimes_K \operatorname{rad}_{\mathcal{B} \# G}((X, g), (Y, h)) = \operatorname{rad}_{\mathcal{B}}(X, Y)$ for all $g, h \in G$.

3. QUASI-KOSZUL CATEGORIES

3.1. Syzygy of functor. Let $\mathcal{A}$ be an additive Krull-Schmidt locally radical finite K -category. We denote by $\operatorname{Mod} \mathcal{A}$ the category of contravariant additive functors from $\mathcal{A}$ to the category $\operatorname{Mod} K$ of K -vector spaces with natural transforms as morphisms.

A family of objects $\{\mathcal{F}(Y_i) | Y_i \in \mathcal{A}, i \in I\}$ is said to be a *family of generators* for $\mathcal{F} \in \operatorname{Mod} \mathcal{A}$ if for every $Y \in \mathcal{A}$,

$$\mathcal{F}(Y) = \sum_{i \in I} \mathcal{F}(\phi)(\mathcal{F}(Y_i))$$

where all but a finite number of $\phi_i : Y \rightarrow Y_i$ are zero, and $\mathcal{F} \in \operatorname{Mod} \mathcal{A}$ is *finitely generated* if it has a finite set of generator.

Denote by $\operatorname{mod} \mathcal{A}$ the full subcategory of $\operatorname{Mod} \mathcal{A}$ consisting of all finitely generated contravariant additive functors of $\operatorname{Mod} \mathcal{A}$. Then the category $\operatorname{mod} \mathcal{A}$ has, by [1, Corollary 4.13], minimal projective resolution. This gives rise to one-to-one correspondence between the indecomposable objects in $\mathcal{A}$ and the simple objects in $\operatorname{Mod} \mathcal{A}$, where an indecomposable object X in $\mathcal{A}$ gives rise to the simple object $S_X := \operatorname{Hom}_{\mathcal{A}}(-, X) / \operatorname{rad}_{\mathcal{A}}(-, X)$ in $\operatorname{Mod} \mathcal{A}$. Furthermore Yoneda's Lemma implies that an $\mathcal{A}$ -module is projective in $\operatorname{mod} \mathcal{A}$ if and only if it is isomorphic to $\operatorname{Hom}_{\mathcal{A}}(-, X)$ for some $X \in \mathcal{A}$.

Definition 3. The *syzygy functor* $\Omega(\mathcal{F})$ of $\mathcal{F} \in \operatorname{Mod} \mathcal{A}$ is defined by the exact sequence

$$0 \rightarrow \Omega(\mathcal{F}) \rightarrow \operatorname{Hom}_{\mathcal{A}}(-, X) \rightarrow \mathcal{F} \rightarrow 0$$

for some $X \in \mathcal{A}$. Inductively, we can define the n -th syzygy functor $\Omega^n(\mathcal{F})$ of $\mathcal{F}$.

Remark. By Freyd's Theorem (see e.g. [8, p. 16, Theorem 3.1]) $\text{Mod}\mathcal{A}$ is an abelian category with coproduct and a faithful set of small projectives. Thus $\Omega(\mathcal{F}) : \mathcal{A} \rightarrow \text{Mod}K, M \mapsto \Omega(\mathcal{F})(M)$ is a contravariant functor for any $\mathcal{F} \in \text{mod}\mathcal{A}$.

3.2. Quasi-Koszul categories. Recall that an ideal in the category $\mathcal{A}$ is a subbifunctor of the Hom-functor. For two ideals $\mathcal{I}$ and $\mathcal{J}$ we define in a natural way inclusion, intersection and product. In particular the product of two ideals $\mathcal{I}$ and $\mathcal{J}$ is given by

$$\begin{aligned} \mathcal{I}\mathcal{J}(X, Z) &= \left\{ \psi \in \text{Hom}_{\mathcal{A}}(X, Z) \mid \psi \right. \\ &= \sum_{i=1}^n \varphi_i \phi_i, \phi_i \in \text{Hom}_{\mathcal{A}}(X, Y_i), \varphi_i \in \text{Hom}_{\mathcal{A}}(Y_i, Z), Y_i \in \mathcal{A} \left. \right\} \text{ for } X, Z \in \mathcal{A}. \end{aligned}$$

For an n -fold product of an ideal $\mathcal{I}$ with itself we write $\mathcal{I}^n$. Let $\mathcal{I}$ be an ideal in the category. For a contravariant functor $\mathcal{F} : \mathcal{A} \rightarrow \text{Mod}K$ define $\mathcal{I}\mathcal{F} : \mathcal{A} \rightarrow \text{Mod}K$ as the subfunctor of $\mathcal{F}$ given by

$$\mathcal{I}\mathcal{F}(M) = \sum_{X \in \mathcal{A}} \mathcal{F}(\mathcal{I}(M, X))$$

for all objects $M \in \mathcal{A}$. Clearly $\mathcal{I}\mathcal{F} \subseteq \mathcal{F}$ for any ideal $\mathcal{I}$ and any contravariant functor $\mathcal{F}$ in $\text{Mod}\mathcal{A}$.

Definition 4 ([7, §5]). A contravariant functor $\mathcal{F}$ in $\text{Mod}\mathcal{A}$ is *quasi-Koszul* if it has a finitely generated projective resolution

$$\cdots \rightarrow \text{Hom}_{\mathcal{A}}(-, X_i) \rightarrow \cdots \rightarrow \text{Hom}_{\mathcal{A}}(-, X_0) \rightarrow \mathcal{F} \rightarrow 0$$

and $\text{rad}_{\mathcal{A}}\Omega^i(\mathcal{F}) = \text{rad}_{\mathcal{A}}^2(-, X_{i-1}) \cap \Omega^i(\mathcal{F})$ for all $i \geq 1$.

The category $\mathcal{A}$ is *quasi-Koszul* if every simple functor in $\text{Mod}\mathcal{A}$ is quasi-Koszul.

3.3. Proof of Theorem. Since the categories $G * (\mathcal{A} \# G)$ and $\mathcal{A}$ are equivalent, it is sufficient to prove (i). Assume that $\mathcal{A}$ is quasi-Koszul and $\mathcal{F}$ is a simple contravariant functor in $\text{Mod}\mathcal{A}$ having a finitely generated projective resolution

$$\cdots \rightarrow \text{Hom}_{\mathcal{A}}(-, X_i) \rightarrow \cdots \rightarrow \text{Hom}_{\mathcal{A}}(-, X_0) \rightarrow \mathcal{F} \rightarrow 0$$

and that $\text{rad}_{\mathcal{A}}\Omega^i(\mathcal{F}) = \text{rad}_{\mathcal{A}}^2(-, X_{i-1}) \cap \Omega^i(\mathcal{F})$ for all $i \geq 1$.

Suppose that $\mathcal{F} = \text{Hom}_{\mathcal{A}}(-, X)/\text{rad}_{\mathcal{A}}(-, X)$ for some indecomposable object $X \in \mathcal{A}$ and define

$$\mathcal{F}^* := \text{Hom}_{G*\mathcal{A}}(-, X)/\text{rad}_{G*\mathcal{A}}(-, X).$$

Then $\mathcal{F}^*$ is a simple contravariant functor in $\text{Mod } G * \mathcal{A}$, and every simple functor in $\text{Mod } G * \mathcal{A}$ is of this form, which has a finitely generated projective resolution

$$\cdots \rightarrow \text{Hom}_{G*\mathcal{A}}(-, X_i) \rightarrow \cdots \rightarrow \text{Hom}_{G*\mathcal{A}}(-, X_0) \rightarrow \mathcal{F}^* \rightarrow 0.$$

Thus it is enough to show that

$$\begin{aligned} (*) \quad & \text{for all } i \geq 1, \text{rad}_{\mathcal{A}}\Omega^i(\mathcal{F}) = \text{rad}_{\mathcal{A}}^2(-, X_{i-1}) \cap \Omega^i(\mathcal{F}) \\ & \text{if and only if } \text{rad}_{G*\mathcal{A}}\Omega^i(\mathcal{F}^*) = \text{rad}_{G*\mathcal{A}}^2(-, X_{i-1}) \cap \Omega^i(\mathcal{F}^*). \end{aligned}$$

For any object $M \in \mathcal{A}$ and $i \geq 1$, we have

$$\begin{aligned} \text{rad}_{G*\mathcal{A}}\Omega^i(\mathcal{F}^*)(M) &= \sum_{Y \in \mathcal{A}} \Omega^i(\mathcal{F}^*)(\text{rad}_{G*\mathcal{A}}(M, Y)) \\ &= \sum_{Y \in \mathcal{A}} \Omega^i(\mathcal{F}^*)(KG \otimes_K \text{rad}_{\mathcal{A}}(M, Y)) \\ &= \sum_{Y \in \mathcal{A}} KG \otimes_K KG \otimes_K \Omega^i(\mathcal{F})(\text{rad}_{\mathcal{A}}(M, Y)) \\ &= KG \otimes_K KG \otimes_K \text{rad}_{\mathcal{A}}\Omega^i(\mathcal{F})(M); \end{aligned}$$

that is, for all $i \geq 1$,

$$(I) \quad \text{rad}_{G*\mathcal{A}}\Omega^i(\mathcal{F}^*) = KG \otimes_K KG \otimes_K \text{rad}_{\mathcal{A}}\Omega^i(\mathcal{F}).$$

By the Lemma, we have

$$\text{rad}_{G*\mathcal{A}}^2(-, X_{i-1}) \cap \Omega^i(\mathcal{F}^*) = KG \otimes_K KG \otimes_K \text{rad}_{\mathcal{A}}^2(-, X_{i-1}) \cap \Omega^i(\mathcal{F}^*) \quad \text{for all } i \geq 1.$$

Since $\Omega^i(\mathcal{F}^*) \subseteq \text{rad}_{G*\mathcal{A}}\text{Hom}_{G*\mathcal{A}}(-, X_{i-1})$ for all $i \geq 1$,

$$\begin{aligned} \Omega^i(\mathcal{F}^*)(M) &\subseteq \text{rad}_{G*\mathcal{A}}\text{Hom}_{G*\mathcal{A}}(-, X_{i-1}) \\ &= \sum_{Y \in \mathcal{A}} \text{Hom}_{G*\mathcal{A}}(-, X_{i-1})\text{rad}_{G*\mathcal{A}}(M, Y) \\ &= KG \otimes_K KG \otimes_K \text{Hom}_{\mathcal{A}}(-, X_{i-1})\text{rad}_{\mathcal{A}}(M, Y) \\ &= KG \otimes_K KG \otimes_K \text{rad}_{\mathcal{A}}\text{Hom}_{\mathcal{A}}(-, X_{i-1})(M). \end{aligned}$$

On the other hand, note that $\Omega^i(\mathcal{F}) \subseteq \text{rad}_{\mathcal{A}}\text{Hom}_{\mathcal{A}}(-, X_{i-1})$. This yields that $\Omega^i(\mathcal{F}^*) = KG \otimes_K KG \otimes_K \Omega^i(\mathcal{F})$. Hence

$$(II) \quad \text{rad}_{G*\mathcal{A}}^2(-, X_{i-1}) \cap \Omega^i(\mathcal{F}^*) = KG \otimes_K KG \otimes_K \text{rad}_{\mathcal{A}}^2(-, X_{i-1}) \cap \Omega^i(\mathcal{F}).$$

Now (*) follows by combining (I) and (II). This completes the proof. $\square$

REFERENCES

1. M. Auslander, *Representation theory of Artin algebras. I*, Comm. Algebra **1**(1974), 177–268. MR0349747 (50:2240)
2. R. Berger, *Koszulity for nonquadratic algebras*, J. Algebra **239**(2001), 705–734. MR1832913 (2002d:16034)
3. T. Cassidy and B. Shelton, *Generalizing the notion of Koszul algebra*, Math. Z. **260**(2008), 93–114. MR2413345 (2009e:16047)
4. C. Cibils and E.N. Marcos, *Skew category, Galois covering and smash product of a k-category*, Proc. Amer. Math. Soc. **134**(2006), 39–50. MR2170541 (2006e:18003)
5. M. Cohen and S. Montgomery, *Group-graded rings, smash products, and group actions*, Trans. Amer. Math. Soc. **282**(1984), 237–258. MR728711 (85i:16002)
6. R. Martínez-Villa, *Skew group algebras and their Yoneda algebras*, Math. J. Okayama Univ. **43**(2001), 1–16. MR1913868 (2003g:18013)
7. R. Martínez-Villa and Ø. Solberg, *Graded and Koszul categories*, Appl. Categor. Struct. **18**(2010), 615–652.
8. B. Mitchel, *Rings with several objects*, Adv. Math. **8**(1972), 1–161. MR0294454 (45:3524)

9. S.B. Priddy, *Koszul resolutions*, Trans. Amer. Math. Soc. **152**(1970), 39–60. MR0265437 (42:346)
10. D.K. Zhao and Y. Han, *Koszul algebras and finite Galois coverings*, Sci. China Ser. A **52**(2009), 2145–2153. MR2550272

ACADEMY OF MATHEMATICS AND SYSTEMS SCIENCE, CHINESE ACADEMY OF SCIENCES, BEIJING 100190, PEOPLE'S REPUBLIC OF CHINA

E-mail address: `deke@amss.ac.cn`

Current address: School of Applied Mathematics, Beijing Normal University at Zhuhai, Zhuhai, 519087, People's Republic of China