ON GEODESICS OF FINSLER METRICS VIA NAVIGATION PROBLEM

LIBING HUANG AND XIAOHUAN MO

(Communicated by Jianguo Cao)

ABSTRACT. This paper is devoted to a study of geodesics of Finsler metrics via Zermelo navigation. We give a geometric description of the geodesics of the Finsler metric produced from *any* Finsler metric and *any* homothetic field in terms of navigation representation, generalizing a result previously only known in the case of Randers metrics with constant S -curvature. As its application, we present explicitly the geodesics of the Funk metric on a strongly convex domain.

1. INTRODUCTION

A smooth curve in a Finsler manifold is called a *geodesic* if it is locally the shortest path connecting any two nearby points on this curve (cf. [21]). Recently, R. Bryant [5] showed that a Finsler metric on S^2 of constant flag curvature $K = 1$ with reversible geodesics is actually a Riemannian one. M. Crampin's result tells us that if a Randers metric $\alpha + \beta$ has reversible geodesics, then β is closed [8]. C. Robles investigated geodesics of Randers metrics of constant S -curvature [17]. Randers metrics of constant flag curvature (or quadratic Riemann curvature) have constant S -curvature [3, 10]. In terms of the navigation representation, they are produced from Riemannian metrics and homothetic vector fields [13, 16, 22, 23]. In fact, C. Robles classified geodesics in Randers manifolds of constant flag curvature [17].

The aim of this paper is to give a geometric description of the geodesics of the Finsler metric $\tilde{F}$ obtained from an *arbitrary* Finsler metric F and an *arbitrary* homothetic field V of F in terms of the navigation representation. Precisely we show the following:

Theorem 1.1. *Let $F = F(x, y)$ be a Finsler metric on a manifold M and let V be a vector field on M with $F(x, V_x) < 1$. Suppose that V is homothetic with dilation c . Let $\tilde{F} = \tilde{F}(x, y)$ denote the Finsler metric on M defined in (3.20). Then the geodesics of $\tilde{F}$ are given by $\psi_t(\gamma(a(t)))$ where ψ_t is the flow of $-V$, $\gamma(t)$*

Received by the editors June 30, 2010 and, in revised form, August 3, 2010.

2010 *Mathematics Subject Classification.* Primary 58B20.

This work is supported by the National Natural Science Foundation of China 11071005.

The second author is the corresponding author.

©2011 American Mathematical Society
 Reverts to public domain 28 years from publication

is a geodesic of F and $a(t)$ is defined by

$$a(t) := \begin{cases} \frac{e^{2ct}-1}{2c}, & \text{if } c \neq 0; \\ t, & \text{if } c = 0. \end{cases}$$

Our result generalizes a theorem previously only known in the case of Randers metrics with constant S -curvature [17]. As its application, we represent explicitly the geodesics of the Funk metric on a strongly convex domain (see Proposition 4.1).

Recall that a vector field V on a Finsler manifold (M, F) is said to be a *homothetic field of F with dilation c* if the corresponding flow ϕ_t is homothetic with dilation c . In particular V is called a *Killing field* if $c = 0$.

It is worth mentioning our recent result that for a non-Killing homothetic field V , the navigation representation has the flag curvature decreasing property [15].

For interesting results of geodesics on Finsler spheres, we refer the reader to [1, 9, 11].

2. PRELIMINARIES

Let us recall firstly the definition of the Finsler metrics.

Definition 2.1 ([2]). Let M be a finite-dimensional manifold. A function $F : TM \rightarrow [0, +\infty)$ is a *Finsler metric* if it satisfies

- (a) F is C^∞ on $TM \setminus \{0\}$;
- (b) $F(x, \lambda y) = \lambda F(x, y)$ for all $y \in T_x M$, $x \in M$ and $\lambda > 0$;
- (c) for every $y \in T_x M \setminus \{0\}$, the quadratic form

$$g_{x,y}(u, v) \equiv \frac{1}{2} \frac{\partial^2}{\partial s \partial t} F^2(x, y + su + tv)|_{t=s=0}, \quad \forall u, v \in T_x M$$

is positive definite.

In this case, (M, F) is called a *Finsler manifold*. Let SM be the projective sphere bundle of M , obtained from TM by identifying nonzero vectors which differ from each other by a positive multiplicative factor. Each geometrical quantity on TM , homogeneous of degree zero, is considered to sit on SM . Define

$$\omega := F_{y^i} dx^i.$$

Then ω is a differential form on SM . It is easy to verify that

$$\omega \wedge (d\omega)^{n-1} \neq 0, \quad n = \dim M$$

(cf. [6]), i.e., that ω defines a contact structure on SM . This form ω is known in the calculus of variations as the *Hilbert form*.

Since ω is a contact form, there exists a unique vector field X on SM that satisfies

$$\omega(X) = 1, \quad X \lrcorner (d\omega) = 0.$$

This vector field X is known as the *Reeb vector field* [4]. It is easy to see that a C^∞ -curve is a (constant Finslerian speed) geodesic if its canonical lift in SM is an integral curve of the Reeb vector field X [4].

Every vector $y \in T_x M \setminus \{0\}$ uniquely determines a covector $p \in T_x^* M \setminus \{0\}$ by

$$p(u) := \frac{1}{2} \frac{d}{dt} (F^2(x, y + tu))|_{t=0}, \quad u \in T_x M.$$

The resulting map $L_x^F : y \in T_x M \rightarrow p \in T_x^* M$ is called the *Legendre transformation* at x . The family $L^F := \{L_x^F \mid x \in M\}$ is called the *Legendre transformation*.

Define a non-negative scalar function $H = H(x, p)$ by

$$(2.1) \quad H(x, p) := \max_{y \in T_x M \setminus \{0\}} \frac{p(y)}{F(x, y)}.$$

The function H is C^∞ on $T^*M \setminus \{0\}$ and $H_x := H|_{T_x^*M}$ is a Minkowski norm on T_x^*M for $x \in M$. Such a function is called a *Cartan metric* [12, 15] (co-Finsler metric in an alternative terminology [18, 19]). The pair (M, H) is called a *Cartan manifold*.

Every covector $p \in T_x^*M \setminus \{0\}$ uniquely determines a vector $y \in T_x M \setminus \{0\}$ by

$$q(y) := \frac{1}{2} \frac{d}{dt} (H^2(x, p + tq))|_{t=0}, \quad q \in T_x^*M.$$

The resulting map $L_x^{F*} : p \in T_x^*M \rightarrow y \in T_x M$ is called the *inverse Legendre transformation* at x . Indeed L_x^F and L_x^{F*} are inverses of each other. Moreover, they preserve the Minkowski norms

$$H(x, p) = F(x, (L_x^F)^{-1}p).$$

Recently, one of the important approaches in discussing the Finsler metric is the (Zermelo) navigation problem. For instance, Bao-Robles-Shen have classified Randers metrics of constant flag curvature via the navigation problem in a Riemannian manifold [3].

The main technique of the navigation problem is described as follows. Given a Finsler metric F and a vector field V with $F(x, V_x) < 1$, define a new Finsler metric $\tilde{F}$ by

$$F(x, \frac{y}{\tilde{F}(x, y)} + V_x) = 1, \quad \forall x \in M, y \in T_x M.$$

A (local) flow (a *local one-parameter group* in an alternative terminology) on a manifold M is a map $\phi : (-\epsilon, \epsilon) \times M \rightarrow M$, also denoted by $\phi_t := \phi(t, \cdot)$, satisfying

- $\phi_0 = \text{id} : M \rightarrow M$;
- $\phi_s \cdot \phi_t = \phi_{s+t}$ for any $s, t \in (-\epsilon, \epsilon)$ with $s + t \in (-\epsilon, \epsilon)$.

Hence, the lift of a flow ϕ_t on M is a flow $\hat{\phi}_t$ on T^*M ,

$$(2.2) \quad \hat{\phi}_t(x, p) := (\phi_t(x), (\phi_t^*)^{-1}(p)).$$

By the relationship between vector fields and flows, (2.2) induces a natural way to lift a vector field V on M to a vector field X_V^* on T^*M .

A vector field V on a Finsler manifold (M, F) is called *homothetic with dilation c* if its flow ϕ_t satisfies

$$(2.3) \quad F(\phi_t(x), \phi_{t*}(y)) = e^{2ct} F(x, y), \quad \forall x \in M, y \in T_x M.$$

Similarly, a vector field V on a Cartan manifold (M, H) is called *homothetic with dilation c* if its flow ϕ_t satisfies

$$(2.4) \quad H(\phi_t(x), (\phi_t^*)^{-1}(p)) = e^{-2ct} H(x, p), \quad \forall x \in M, p \in T_x^*M.$$

Lemma 2.2. *Let V be a homothetic field on a Finsler manifold (M, F) with dilation c and H its Cartan metric defined by (2.1). Then V is a homothetic field of H with dilation c .*

Proof. By using (2.1) and (2.3) we have

$$\begin{aligned}
H(\phi_t(x), (\phi_t^*)^{-1}(p)) &= \max_{\tilde{y} \in T_{\phi_t(x)} M \setminus \{0\}} \frac{[(\phi_t^*)^{-1}(p)](\tilde{y})}{F(\phi_t(x), \tilde{y})} \\
&= \max_{\tilde{y} \in T_{\phi_t(x)} M \setminus \{0\}} \frac{p((\phi_{t*})^{-1}(\tilde{y}))}{F(\phi_t(x), \tilde{y})} \\
&= \max_{y \in T_x M \setminus \{0\}} \frac{p(y)}{F(\phi_t(x), \phi_{t*}(y))} \\
&= \max_{y \in T_x M \setminus \{0\}} \frac{p(y)}{e^{2ct} F(x, y)} \\
&= e^{-2ct} \max_{y \in T_x M \setminus \{0\}} \frac{p(y)}{F(x, y)} = e^{-2ct} H(x, p),
\end{aligned}$$

where $y := (\phi_{t*})^{-1}(\tilde{y})$. It follows that V is a homothetic field of H with dilation c . $\square$

3. GEODESICS OF A FINSLER METRIC VIA NAVIGATION PROBLEM

In this section, we give a geometric description of geodesics of Finsler metrics via the homothetic navigation problem in a Finsler manifold. First, we show the following:

Lemma 3.1. *Let N be a manifold, and let V and W be vector fields on N that satisfy*

$$(3.1) \quad [V, W] = -cV$$

for some constant c . Let ϕ_t and ψ_t be local 1-parameter groups of V and W respectively. Then $\psi_t \circ \phi_{a(t)}$ is a local 1-parameter group of the vector field $V + W$, where $a(t)$ is defined by

$$a(t) := \begin{cases} \frac{e^{ct}-1}{c}, & \text{if } c \neq 0; \\ t, & \text{if } c = 0. \end{cases}$$

Proof. Direct calculations yield

$$(3.2) \quad \frac{da}{dt} = e^{ct}.$$

Since V is the induced vector field from ϕ ,

$$(3.3) \quad \frac{d}{dt} \phi_t(x)|_{t=t_0} = V_{\phi_{t_0}(x)}.$$

Let $\eta_t := \phi_{a(t)}$, $u := a(t)$. By using (3.2) and (3.3) we obtain

$$(3.4) \quad \frac{d}{dt} [\eta_t(x)]_{t=s} = \frac{d\phi_u}{du}|_{u=a(s)} \frac{da}{dt}|_{t=s} = e^{cs} V|_{\phi_{a(s)}(x)} = e^{cs} V|_{\eta_s(x)}.$$

From (3.1), one obtains $[W, V] = cV$. It follows that

$$\begin{aligned}
(3.5) \quad c\psi_{t*}V &= \psi_{t*}(cV) \\
&= \psi_{t*}[W, V] \\
&= [\psi_{t*}W, \psi_{t*}V] = [W, \psi_{t*}V].
\end{aligned}$$

This gives

$$\begin{aligned}
 c\psi_{t*}V_p(f) &= [W, \psi_{t*}V]_p f \\
 &= [(\mathcal{L}_W(\psi_{t*}V))]_p f \\
 &= \lim_{\Delta t \rightarrow 0} \frac{1}{\Delta t} [\psi_{t*}V - (\psi_{t+\Delta t})_* V]_p f = -\frac{d}{dt} [\psi_{t*}V]_p f
 \end{aligned}
 \tag{3.6}$$

for a point $p \in N$ and a function $f \in \mathcal{C}^\infty(N)$. We set

$$y(t) := \psi_{t*}V_p(f).$$

Substituting (3.7) into (3.6) yields

$$\frac{dy}{dt} = -cy.$$

Solving (3.8), we get

$$y = C_1 e^{-ct}.$$

Plugging (3.7) into (3.9) yields

$$\psi_{t*}V_p(f) = C_1(p, f)e^{-ct}.$$

Taking $t = 0$, then $C_1(p, f) = V_p f$. Plugging this into (3.10) yields $(\psi_t)_* V_p(f) = e^{-ct} V_p f$ for any point $p \in N$ and any function $f \in \mathcal{C}^\infty(N)$. It follows that

$$(\psi_t)_* V = e^{-ct} V.$$

By using (3.4), for any function $f \in \mathcal{C}^\infty(N)$ we have

$$\begin{aligned}
 \frac{d}{dt} [\psi_t \circ \eta_t(x)]_{t=s} f &= \lim_{\Delta s \rightarrow 0} \frac{1}{\Delta s} [f \circ \psi_{s+\Delta s} \circ \eta_{s+\Delta s}(x) - f \circ \psi_s \circ \eta_s(x)] \\
 &= \lim_{\Delta s \rightarrow 0} \frac{1}{\Delta s} [f \circ \psi_{s+\Delta s} \circ \eta_{s+\Delta s}(x) - f \circ \psi_s \circ \eta_{s+\Delta s}(x)] \\
 &\quad + \lim_{\Delta s \rightarrow 0} \frac{1}{\Delta s} [f \circ \psi_s \circ \eta_{s+\Delta s}(x) - f \circ \psi_s \circ \eta_s(x)] \\
 &= \lim_{\Delta s \rightarrow 0} \frac{f \circ \psi_{s+\Delta s} - f \circ \psi_s}{\Delta s} (\eta_{s+\Delta s}(x)) \\
 &\quad + (f \circ \psi_s)_* \left(\lim_{\Delta s \rightarrow 0} \frac{\eta_{s+\Delta s}(x) - \eta_s(x)}{\Delta s} \right) \\
 &= W_{\psi_s \circ \eta_s(x)} f + \psi_{s*}(e^{cs} V|_{\eta_s(x)}) f.
 \end{aligned}$$

It follows that

$$\frac{d}{dt} [\psi_t \circ \eta_t(x)]_{t=s} = W_{\psi_s \circ \eta_s(x)} + \psi_{s*}(e^{cs} V|_{\eta_s(x)}).$$

By using (3.11) we have

$$\psi_{s*}(e^{cs} V|_{\eta_s(x)}) = e^{cs} \psi_{s*}(V|_{\eta_s(x)}) = e^{cs} e^{-cs} V_{\psi_s \circ \eta_s(x)} = V_{\psi_s \circ \eta_s(x)}.$$

Plugging this into (3.12) yields

$$\frac{d}{dt} [\psi_t \circ \eta_t(x)]_{t=s} = W_{\psi_s \circ \eta_s(x)} + V_{\psi_s \circ \eta_s(x)} = (W + V)_{\psi_s \circ \eta_s(x)}.$$

It follows that $\psi_t \circ \eta_t$ is a local 1-parameter group of the vector field $V + W$. $\square$

Lemma 3.2. *For a homothetic field V on a Cartan manifold (M, H) with dilation c , we have*

$$(3.13) \quad [X^\flat, X_V^*] = 2cX^\flat,$$

where $X^\flat = (L^F)_*X$ and X_V^* is the lift of V to T^*M .

Proof. In natural coordinates, we have

$$(3.14) \quad X_V^* = v^i \frac{\partial}{\partial x^i} - p_j \frac{\partial v^j}{\partial x^i} \frac{\partial}{\partial p_i},$$

where $V = v^i \frac{\partial}{\partial x^i}$ [15, (5.3)]. Note that V is homothetic with respect to H with dilation c . Differentiating (2.4) with respect to t at $t = 0$ yields

$$(3.15) \quad X_V^*(H) = -2cH.$$

By using (3.14), we have

$$X_V^*(H) = v^i \frac{\partial H}{\partial x^i} - p_j \frac{\partial v^j}{\partial x^i} \frac{\partial H}{\partial p_i}.$$

It follows that

$$(3.16) \quad v^i \frac{\partial H}{\partial x^i} - p_j \frac{\partial v^j}{\partial x^i} \frac{\partial H}{\partial p_i} = -2cH.$$

Differentiating (3.16) with respect to p_k gives

$$(3.17) \quad v^i \frac{\partial^2 H}{\partial x^i \partial p_k} - \sum_i \frac{\partial v^k}{\partial x^i} \frac{\partial H}{\partial p_i} - \sum_i p_j \frac{\partial v^j}{\partial x^i} \frac{\partial^2 H}{\partial p_i \partial p_k} = -2c \frac{\partial H}{\partial p_k}.$$

Differentiating (3.16) with respect to x^k yields

$$(3.18) \quad \frac{\partial v^i}{\partial x^k} \frac{\partial H}{\partial x^i} + v^i \frac{\partial^2 H}{\partial x^i \partial x^k} - p_j \frac{\partial^2 v^j}{\partial x^i \partial x^k} \frac{\partial H}{\partial p_i} - p_j \frac{\partial v^j}{\partial x^i} \frac{\partial^2 H}{\partial p_i \partial x^k} = -2c \frac{\partial H}{\partial x^k}.$$

By Lemma 4.5 in [15], $X^\flat$ is the Hamiltonian vector field for H . Hence it has the local expression

$$(3.19) \quad X^\flat = \frac{\partial H}{\partial p_i} \frac{\partial}{\partial x^i} - \frac{\partial H}{\partial x^i} \frac{\partial}{\partial p_i}.$$

It follows from (3.14) and (3.19) that

$$\begin{aligned} [X^\flat, X_V^*] &= \left[\frac{\partial H}{\partial p_i} \frac{\partial}{\partial x^i} - \frac{\partial H}{\partial x^i} \frac{\partial}{\partial p_i}, v^j \frac{\partial}{\partial x^j} - p_k \frac{\partial v^k}{\partial x^j} \frac{\partial}{\partial p_j} \right] \\ &= \left[\frac{\partial H}{\partial p_i} \frac{\partial}{\partial x^i}, v^j \frac{\partial}{\partial x^j} \right] - \left[\frac{\partial H}{\partial x^i} \frac{\partial}{\partial p_i}, v^j \frac{\partial}{\partial x^j} \right] \\ &\quad - \left[\frac{\partial H}{\partial p_i} \frac{\partial}{\partial x^i}, p_k \frac{\partial v^k}{\partial x^j} \frac{\partial}{\partial p_j} \right] + \left[\frac{\partial H}{\partial x^i} \frac{\partial}{\partial p_i}, p_k \frac{\partial v^k}{\partial x^j} \frac{\partial}{\partial p_j} \right]. \end{aligned}$$

Recall that

$$[fX, gY] = f(Xg)Y - g(Yf)X + fg[X, Y]$$

for any vector fields X, Y and functions f, g . It follows that

$$\begin{aligned}
 [X^\flat, X_V^*] &= \frac{\partial H}{\partial p_i} \frac{\partial v^j}{\partial x^i} \frac{\partial}{\partial x^j} - v^j \frac{\partial^2 H}{\partial p_i \partial x_j} \frac{\partial}{\partial x^i} + v^j \frac{\partial^2 H}{\partial x^i \partial x^j} \frac{\partial}{\partial p_i} \\
 &\quad - p_k \frac{\partial H}{\partial p_i} \frac{\partial^2 v^k}{\partial x^i \partial x^j} \frac{\partial}{\partial p_j} + p_k \frac{\partial v^k}{\partial x^j} \frac{\partial^2 H}{\partial p_i \partial p_j} \frac{\partial}{\partial x^i} \\
 &\quad + \frac{\partial H}{\partial x_i} \frac{\partial v^i}{\partial x^j} \frac{\partial}{\partial p_j} - p_k \frac{\partial v^k}{\partial x^j} \frac{\partial^2 H}{\partial x^i \partial p_j} \frac{\partial}{\partial p_i} \\
 &= - \left(v^j \frac{\partial^2 H}{\partial x^j \partial p_k} - \sum_i \frac{\partial v^k}{\partial x^i} \frac{\partial H}{\partial p_i} - \sum_j p_i \frac{\partial v^i}{\partial x^j} \frac{\partial^2 H}{\partial p_j \partial p_k} \right) \frac{\partial}{\partial x^k} \\
 &\quad + \left(\frac{\partial v^i}{\partial x^k} \frac{\partial H}{\partial x^i} + v^j \frac{\partial^2 H}{\partial x^j \partial x^k} - p_j \frac{\partial^2 v^j}{\partial x^i \partial x^k} \frac{\partial H}{\partial p_i} - p_i \frac{\partial v^i}{\partial x^j} \frac{\partial^2 H}{\partial p_j \partial x^k} \right) \frac{\partial}{\partial p_k} \\
 &= 2c \frac{\partial H}{\partial p_k} \frac{\partial}{\partial x^k} - 2c \frac{\partial H}{\partial x^k} \frac{\partial}{\partial p_k} = 2c X^\flat.
 \end{aligned}$$

Here we have used (3.17), (3.18) and (3.19). $\square$

Proof of Theorem 1.1. Let V be a homothetic field on a Finsler manifold (M, F) with dilation c and let X be the Reeb vector field. Applying the Legendre transformation, we obtain a Cartan metric $H(x, p)$. From Lemma 2.2, V is a homothetic field of (M, H) with dilation c . By Lemma 3.2 and (3.14), we have $[X^\flat, X_{-V}^*] = -2c X^\flat$. Let ϕ_t and $\hat{\psi}_t$ be local 1-parameter groups of $X^\flat$ and X_{-V}^* respectively. By using (3.14) we get $X^\flat + X_{-V}^* = X^\flat - X_V^*$. Taking this together with Lemma 3.1, we obtain that $\hat{\psi}_t \circ \phi_{a(t)}$ is a local 1-parameter group of $X^\flat - X_V^*$, where

$$a(t) = \begin{cases} \frac{e^{2ct}-1}{2c}, & \text{if } c \neq 0; \\ t, & \text{if } c = 0. \end{cases}$$

Recall that a navigation problem makes use of a Finsler metric F and a vector field V with $F(x, V_x) < 1$ and produces a new Finsler metric $\tilde{F}$ by solving the equation

$$(3.20) \quad F(x, y + \tilde{F}(x, y)V) = \tilde{F}(x, y).$$

Let $\tilde{X}$ be the Reeb vector field of $\tilde{F}$ and $\tilde{X}^\flat = (L^{\tilde{F}})_* \tilde{X}$. By Lemma 6.2 in [15], we have

$$\tilde{X}^\flat = X^\flat - X_V^*.$$

It follows that $\hat{\psi}_t \circ \phi_{a(t)}$ is a local 1-parameter group of $\tilde{X}^\flat$.

Since $L^F(x, y) = (x, L_x^F(y))$, we see that any geodesic of F is precisely the projection of an integral curve of $X^\flat$. Similarly, a geodesic of $\tilde{F}$ is precisely the projection of an integral curve of $\tilde{X}^\flat$. Note that

$$\hat{\psi}_t(x, p) = (\psi_t(x), (\psi_t^*)^{-1}(p)),$$

where ψ_t is the flow produced by $-V$.

Let $\pi : T^*M \setminus \{0\} \rightarrow M$ be the natural projection. It follows that

$$\pi \circ \hat{\psi}_t(x, p) = \pi(\psi_t(x), (\psi_t^*)^{-1}(p)) = \psi_t(x) = \psi_t \circ \pi(x, p)$$

for any $x \in M$ and $p \in T_x M \setminus \{0\}$. Hence we have

$$(3.21) \quad \pi \circ \hat{\psi}_t = \psi_t \circ \pi.$$

It follows that

$$\pi \circ \hat{\psi}_t \circ \phi_{a(t)} = \psi_t \circ \pi \circ \phi_{a(t)} = \psi_t(\gamma(a(t))),$$

where $\gamma(t) := \pi(\phi_t(x))$. Thus we have proved Theorem 1.1.

Remark 3.1. The reader should note that the navigation problem adopted here differs from those of C. Robles and Z. Shen [17, 20], where the navigation problem is defined by

$$F(x, \frac{y}{\tilde{F}(x, y)} - V) = 1;$$

i.e., the $\tilde{F}$ we define with (F, V) is precisely the $\tilde{F}$ that Shen defines with $(F, -V)$.

4. GEODESICS OF FUNK METRICS ON CONVEX DOMAINS

In this section we are going to represent explicitly the geodesics of the Funk metric on a strongly convex domain.

Given a Minkowski norm $\varphi : \mathbb{E} \rightarrow \mathbb{R}$ on a vector space $\mathbb{E}$, one can construct $\Omega := \{v \in \mathbb{E} \mid \varphi(v) < 1\}$, $T_x \Omega \simeq \mathbb{E}$. A domain Ω in $\mathbb{E}$ defined by a Minkowski norm φ is called a *strongly convex domain* [19]. Thus $(\Omega, F(x, y))$ is a Minkowski manifold, where $F(x, y) := \varphi(y)$. For each $x \in \Omega$, identify $T_x \Omega$ with $\mathbb{E}$. Thus $V_x := x$ is a radical vector field on Ω satisfying $F(x, V_x) = \varphi(x) < 1$. Moreover V is a homothetic field of F with dilation $c = \frac{1}{2}$ [15, Example 1]. Define a 1-parameter transformation ψ_t on Ω by

$$\psi_t(x) = e^{-t}x.$$

Note that $T_{\psi_t(x)}\Omega \simeq \mathbb{E}$ for any t . It is easy to see that

$$\psi_{t*}(y) = e^{-t}y, \quad \text{for } \forall y \in T_x \Omega.$$

Thus we have

$$\begin{aligned} F(\psi_t(x), \psi_{t*}(y)) &= \varphi(\psi_{t*}(y)) \\ &= \varphi(e^{-t}y) \\ &= e^{-t}\varphi(y) = e^{-t}F(x, y) \end{aligned}$$

for any $(x, y) \in T\Omega$. It follows that φ_t is homothetic. A direct calculation yields

$$\frac{d\psi_t(x)}{dt} \Big|_{t=0} = -x = -V_x.$$

Thus ψ_t is the flow of the vector field $-V$. By using the Minkowski metric F and the homothetic field V , we produce a new Finsler metric $\tilde{F}$ in terms of the navigation problem. $\tilde{F}$ is called the *Funk metric* on a strongly convex domain Ω (cf. [7, Example 3.4.3]).

Since F is Minkowskian, it is locally projectively flat. This implies that the unit speed geodesic through x with tangent vector $y(\neq 0)$ is $\gamma(t) := x + \frac{t}{\varphi(y)}y$. Note that V is a homothetic field with dilation $c = \frac{1}{2}$. We have

$$a(t) = e^t - 1$$

(see Theorem 1.1). It follows that

$$\psi_t(\gamma(a(t))) = e^{-t} \left[x + \frac{e^t - 1}{\varphi(y)} y \right].$$

By Theorem 1.1, we have the following.

Proposition 4.1. *Let $\varphi : \mathbb{E} \rightarrow \mathbb{R}$ be a Minkowski norm and Ω its strongly convex domain. Assume that $\tilde{F}$ is the Funk metric on Ω defined by*

$$\varphi\left(\frac{y}{\tilde{F}(x, y)} + x\right) = 1.$$

Then the geodesics of $\tilde{F}$ are given by

$$e^{-t} \left[x + \frac{e^t - 1}{\varphi(y)} y \right].$$

REFERENCES

1. V. Bangert and Y. Long, *The existence of two closed geodesics on every Finsler 2-sphere*, Math. Ann. 346 (2010), 335-366. MR2563691
2. D. Bao, S.-S. Chern and Z. Shen, *An introduction to Riemann-Finsler geometry*. Graduate Texts in Mathematics, 200. Springer-Verlag, New York, 2000. MR1747675 (2001g:53130)
3. D. Bao, C. Robles and Z. Shen, *Zermelo navigation on Riemannian manifolds*. J. Differential Geom. 66 (2004), 377-435. MR2106471 (2005k:58023)
4. R. Bryant, *Some remarks on Finsler manifolds with constant flag curvature*, special issue for S. S. Chern. Houston J. Math. 28 (2002), 221-262. MR1898190 (2003h:53102)
5. R. Bryant, *Geodesically reversible Finsler 2-spheres of constant curvature*, inspired by S. S. Chern, 95-111, Nankai Tracts Math., 11, World Sci. Publ., Hackensack, NJ, 2006. MR2313331 (2008a:53075)
6. S.-S. Chern, *Riemannian geometry as a special case of Finsler geometry*, Contemp. Math., 196. Amer. Math. Soc., 1996, 51-58. MR1403576 (98e:53026)
7. S.-S. Chern and Z. Shen, *Riemann-Finsler geometry*. Nankai Tracts in Mathematics, 6. World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2005. MR2169595 (2006d:53094)
8. M. Crampin, *Randers spaces with reversible geodesics*, Publ. Math. Debrecen 67 (2005), 401-409. MR2162131 (2006e:53130)
9. H. Duan and Y. Long, *Multiplicity and stability of closed geodesics on bumpy Finsler 3-spheres*, Calc. Var. Partial Differential Equations 31 (2008), 483-496. MR2372902 (2008m:58025)
10. B. Li and Z. Shen, *On Randers metrics of quadratic Riemann curvature*, Internat. J. Math. 20 (2009), 369-376. MR2500075 (2010a:53161)
11. Y. Long and H. Duan, *Multiple closed geodesics on 3-spheres*, Adv. Math. 221 (2009), 1757-1803. MR2522828
12. R. Miron, D. Hrimiuc, H. Shimada and S. Sabau, *The geometry of Hamilton and Lagrange spaces*. Fundamental Theories of Physics, 118. Kluwer Academic Publishers Group, Dordrecht, 2001. MR1839409 (2002e:53111)
13. X. Mo, *A global classification result for Randers metrics of scalar curvature on closed manifolds*, Nonlinear Anal. 69 (2008), 2996-3004. MR2452109 (2009k:53195)
14. X. Mo, *On some Finsler metrics of constant (or scalar) flag curvature*, accepted for publication in Houston Journal of Mathematics.
15. X. Mo and L. Huang, *On curvature decreasing property of a class of navigation problems*, Publ. Math. Debrecen 71 (2007), 141-163. MR2340039 (2008g:53092)
16. X. Mo and C. Yu, *On the Ricci curvature of a Randers metric of isotropic S-curvature*, Acta Math. Sin. (Engl. Ser.) 24 (2008), 911-916. MR2428109 (2009b:53128)
17. C. Robles, *Geodesics in Randers spaces of constant curvature*, Trans. Amer. Math. Soc. 359 (2007), 1633-1651. MR2272144 (2007h:53119)
18. Z. Shen, *Two-dimensional Finsler metrics with constant flag curvature*, Manuscripta Math. 109 (2002), 349-366. MR1948019 (2003k:53091)

19. Z. Shen, *Landsberg curvature, S-curvature and Riemann curvature. A sampler of Riemann-Finsler geometry*, 303-355, Math. Sci. Res. Inst. Publ. 50. Cambridge Univ. Press, Cambridge, 2004. MR2132662 (2005k:53132)
20. Z. Shen, *Finsler metrics with $\mathbf{K} = 0$ and $\mathbf{S} = 0$* , Canad. J. Math. 55 (2003), 112-132. MR1952328 (2004e:53112)
21. Z. Shen, *Lectures on Finsler geometry*, World Scientific Publishing Co., Singapore, 2001. MR1845637 (2002f:53032)
22. Z. Shen, and H. Xing, *On Randers metrics with isotropic S-curvature*, Acta Math. Sin. (Engl. Ser.) 24 (2008), 789-796. MR2403114 (2010c:53109)
23. E. Zermelo, *Über das Navigationsproblem bei ruhender oder veränderlicher Windverteilung*, Z. Angew. Math. Mech. 11 (1931), 114-124.

SCHOOL OF MATHEMATICAL SCIENCES, NANKAI UNIVERSITY, TIANJIN 300071, PEOPLE'S REPUBLIC OF CHINA

E-mail address: `huanglb@nankai.edu.cn`

KEY LABORATORY OF PURE AND APPLIED MATHEMATICS, SCHOOL OF MATHEMATICAL SCIENCES, PEKING UNIVERSITY, BEIJING 100871, PEOPLE'S REPUBLIC OF CHINA

E-mail address: `moxh@pku.edu.cn`