MONOMIALS AS SUMS OF POWERS: THE REAL BINARY CASE

MATS BOIJ, ENRICO CARLINI, AND ANTHONY V. GERAMITA

(Communicated by Irena Peeva)

ABSTRACT. We generalize an example, due to Sylvester, and prove that any monomial of degree d in $\mathbb{R}[x_0, x_1]$, which is not a power of a variable, cannot be written as a linear combination of fewer than d powers of linear forms.

1. INTRODUCTION

It is well-known, and easy to prove, that if k is a field of characteristic zero and $R = k[x_0, \dots, x_n] = \bigoplus_{i=0}^{\infty} R_i$ is the standard graded polynomial algebra, then the k -vector space R_d (for any d) has a basis consisting of polynomials $\{L_1^d, \dots, L_s^d\}$, where $s = \binom{d+n}{n} = \dim_k R_d$ and the L_i are pairwise linearly independent forms in R_1 . It follows that every form in R_d is a k -linear combination of at most s d^{th} powers of linear forms and, if k is algebraically closed, simply a sum of at most s d^{th} powers of linear forms. We will call such a way of writing $F \in R_d$ a *Waring expansion* of F because of the echo of Waring's problem from number theory. We will further refer to such an expression as a *minimal Waring expansion* for F if the number of summands in such an expression for F is minimal among all such representations.

If $n > 0$ and $d = 2$ it is a classical fact that although $s = \binom{n+2}{2}$, every quadratic form has a Waring expansion involving $\leq n + 1 < s$ summands. In general, i.e. for $[F]$ belonging to a non-empty Zariski open subset of $\mathbb{P}(R_2)$, a minimal Waring expansion for the quadric F has exactly $n + 1$ summands.

These observations have led to a series of problems, usually called **Waring Problems**, which ask for information on minimal Waring expansions for forms of degree d in R .

The long outstanding problem of finding the number of summands in a minimal Waring expansion of the generic form of degree d was solved, after being open for almost 100 years, by J. Alexander and A. Hirschowitz (see [AH95]), when k is an algebraically closed field of characteristic zero. (For positive characteristic, see [IK99, Theorem 1.61].)

Of course, solving this problem for the generic form of degree d does not always give information about any specific form of degree d , and the problem of finding the length of the minimal Waring expansion for specific forms has also been a continuing

Received by the editors May 17, 2010.

2010 *Mathematics Subject Classification.* Primary 14P99; Secondary 13A02, 15A21.

source of interesting speculations and lovely results. For example, it was Sylvester ([Har92]) who first observed that although for $R = \mathbb{C}[x_0, x_1]$, the generic form of degree d has a Waring expansion with $s = \lceil \frac{d+1}{2} \rceil$ summands, the monomial $x_0 x_1^{d-1}$ has d summands in its minimal Waring expansion (the maximum possible).

The Waring problem for specific forms has been considered in depth by B. Reznick in his monograph (see [Rez92]) and by G. Comas and M. Seiguer, who, to our knowledge, were the first to resolve the problem completely and algorithmically in $\mathbb{C}[x_0, x_1]$ (cf. [CS01]).

It is interesting to note that although the Waring problem is a very interesting and stimulating problem in purely algebraic terms, it has a surprising number of intimate connections with problems in areas as seemingly disparate as algebraic geometry and communication theory (see for example [RS00], [CC03] and [CM96]).

Indeed, if $k = \mathbb{R}$, the field of real numbers, the connection with real world problems is very direct. This has prompted a reexamination of the Waring problem for $R = \mathbb{R}[x_0, x_1]$, and a recent, very suggestive paper of P. Comon and G. Ottaviani (see [CO09]) considered this very problem for degrees $d \leq 5$.

Our main result in this paper follows the line of Sylvester's examples and concerns the minimal Waring expansion for monomials in $\mathbb{R}[x_0, x_1]$. We first see how the Apolarity Lemma produces a simple proof of the fact that the minimal Waring expansion of the monomial $x_0^a x_1^b$ in $\mathbb{C}[x_0, x_1]$ with $0 < a \leq b$ has $b + 1$ summands. In sharp contrast to this we show that in $\mathbb{R}[x_0, x_1]$ every monomial of degree d (except x_0^d and x_1^d) has d summands in its minimal Waring expansion.

2. BASIC RESULTS

Let $S = k[x_0, x_1]$ and $T = k[y_0, y_1]$, where k is an algebraically closed field of characteristic zero. We make S into a T -module using differentiation; i.e., we think of $y_0 = \partial/\partial x_0$ and $y_1 = \partial/\partial x_1$. We refer to a polynomial in T as ∂ instead of using capital letters. In particular, for any form F in S_d we define the ideal $F^\perp \subseteq T$ as follows:

$$F^\perp = \{\partial \in T : \partial F = 0\}.$$

The following *Apolarity Lemma* is a classical result whose proof can be found in [IK99, Lemma 1.31].

Lemma 2.1. *A homogeneous form $F \in S$ can be written as*

$$F(x_0, x_1) = \sum_{i=1}^r \alpha_i (L_i)^d, \quad L_i \text{ pairwise linearly independent, } \alpha_i \in k,$$

i.e. has a Waring expansion with r summands, if and only if the ideal $F^\perp$ contains the product of r distinct linear forms.

3. BINARY MONOMIALS: THE COMPLEX CASE

The complex case is straightforward for monomials.

Proposition 3.1. *Let $M = x_0^a x_1^b$ be a monomial in $\mathbb{C}[x_0, x_1]$. If $0 < a \leq b$, then M has a minimal Waring expansion with $b + 1$ summands, i.e. is a sum of $b + 1$ powers of linear forms and no fewer.*

Proof. Let $I = M^\perp = (y_0^{a+1}, y_1^{b+1})$ and notice that the linear system defined by I_{b+1} is base point free on $\mathbb{P}^1 = \mathbb{P}S_1$. Applying Bertini's Theorem, we get that the generic element of I_{b+1} defines a set of $b+1$ distinct points and hence it is the product of $b+1$ distinct linear forms. Thus the Apolarity Lemma yields that M is the sum of $b+1$ powers of linear forms. If $r < b+1$, then r powers do not suffice as no element in $I_r = (y_0^{a+1})_r$ is a product of r distinct linear forms. $\square$

4. BINARY MONOMIALS: THE REAL CASE

We can also ask for a real Waring expansion of a monomial M . More precisely, we want to write

$$M(x_0, x_1) = \sum_{i=1}^r \alpha_i (L_i)^d, \quad \alpha_i \in \{1, -1\},$$

where the linear forms L_i are in $\mathbb{R}[x_0, x_1]$. In order to do this, we have to increase the number of summands in Proposition 3.1.

The following elementary facts will be extremely useful.

Lemma 4.1. *Consider the degree d polynomial*

$$F(x) = c_d x^d + \cdots + c_1 x + c_0 \in \mathbb{R}[x].$$

If $c_i = c_{i-1} = 0$ for some $1 \leq i \leq d$, then $F(x)$ does not have d real distinct roots.

Proof. The proof is obvious if $i = 1$ or $i = d$, so we may as well assume that $1 < i < d$.

Consider all the pairs (c_r, c_s) of non-zero coefficients such that $r > s$ and $c_j = 0$ if $r > j > s$. Let α be the number of pairs such that $r - s$ is odd and β the number of pairs such that $r - s$ is even. Notice that, by hypothesis, $\alpha + 2\beta < d - 1$.

Now we apply Descartes' rule of signs. For a pair (c_r, c_s) such that $r - s$ is odd we get a real root of $F(x)$. To see this notice that if $c_r c_s < 0$, then we have a non-negative real root of $F(x)$. If $c_r c_s > 0$, then we have a non-negative real root of $F(-x)$, hence a negative real root of $F(x)$. Similarly, for a pair (c_r, c_s) such that $r - s$ is even we get either two real roots of $F(x)$ or none.

In conclusion, the number of real roots of $F(x)$ is at most $\alpha + 2\beta$, and we are done. $\square$

Lemma 4.2. *For each $i < d$ there exists a degree d polynomial $F(x) = c_d x^d + \cdots + c_1 x + c_0 \in \mathbb{R}[x]$ having d real distinct roots and such that $c_i = 0$.*

Proof. Choose $a_1, \dots, a_d \in \mathbb{R}$ and consider the polynomial $F(x) = (x - a_1) \cdots (x - a_d)$. This polynomial can also be written as

$$F(x) = \sum_{i=0}^d E_i(a_1, \dots, a_d) x^i,$$

where E_i is the degree i elementary symmetric function in its arguments. The vanishing of the i -th coefficient of $F(x)$ can be written as

$$E_i(a_1, \dots, a_{d-1}) + a_d E_{i-1}(a_1, \dots, a_{d-1}) = 0.$$

Hence, if we choose the $a_1, \dots, a_{d-1} > 0$ and distinct there exists a unique, negative value of a_d such that the coefficient of x^i in $F(x)$ is zero. As the roots of $F(x)$ are $a_1, \dots, a_d$ the polynomial has d real, distinct roots. $\square$

Using the previous results we immediately get a lower bound on the number of summands in the minimal Waring expansion of a monomial in $\mathbb{R}[x_0, x_1]$.

Lemma 4.3. *Let $M = x_0^a x_1^b$ be a monomial in $\mathbb{R}[x_0, x_1]$. If $0 < a \leq b$, then M does not have a Waring expansion with $r \leq a + b - 1$ real summands.*

Proof. Let $I = M^\perp = (y_0^{a+1}, y_1^{b+1})$. The general degree r element in I has the form

$$\begin{aligned} F(y_0, y_1) \\ = c_r y_0^r + c_{r-1} y_0^{r-1} y_1 + \cdots + c_{a+1} y_0^{a+1} y_1^{r-a-1} + c_{r-b-1} y_0^{r-b-1} y_1^{b+1} + \cdots + c_0 y_1^r. \end{aligned}$$

If $a + 1 \geq r - b + 2$, then by Lemma 4.1, $F(y_0, y_1)$ is not the product of r real linear forms. The conclusion follows by the Apolarity Lemma. $\square$

Proposition 4.4. *Let $M = x_0^a x_1^b$ be a monomial in $\mathbb{R}[x_0, x_1]$. If $0 < a \leq b$, then M has a minimal Waring expansion with $a + b$ summands which are powers of real linear forms.*

Proof. We have that $M^\perp = I = (y_0^{a+1}, y_1^{b+1})$. Notice that I_{a+b} is the subspace of T_{a+b} of polynomials which are missing all the monomials having factor y_0^a or y_1^b . Thus, Lemma 4.2 and the Apolarity Lemma yield the result. $\square$

Remark 4.5. A few months after this paper was submitted, Bruce Reznick posted an article on the arXiv [Rez10] in which he considers the Waring expansion for binary forms over several fields k . Reznick's paper has many connections with this work, and, for example, his Corollary 4.11 contains our Proposition 4.4.

ACKNOWLEDGEMENT

This project started with a research visit supported by a grant for visiting researchers from the Göran Gustafsson Foundation.

REFERENCES

- [AH95] J. Alexander and A. Hirschowitz. Polynomial interpolation in several variables. *J. Algebraic Geom.*, 4(2):201–222, 1995. MR1311347 (96f:14065)
- [CC03] E. Carlini and J. Chipalkatti. On Waring's problem for several algebraic forms. *Comment. Math. Helv.*, 78(3):494–517, 2003. MR1998391 (2005b:14097)
- [CS01] G. Comas and M. Seiguer. On the rank of a binary form. *Found. Comput. Math.*, 11(1):65–78, 2010.
- [CM96] P. Comon and B. Mourrain. Decomposition of quantics in sums of power of linear forms. *Signal Processing*, 53(2):93–107, 1996. Special issue on High-Order Statistics.
- [CO09] P. Comon and G. Ottaviani. On the typical rank of real binary forms. arXiv:0909.4865v1, 2009.
- [Har92] J. Harris. *Algebraic geometry, A first course*. Graduate Texts in Math., Springer-Verlag, New York, 1992. MR1182558 (93j:14001)
- [IK99] A. Iarrobino and V. Kanev. Power sums, Gorenstein algebras, and determinantal loci. *Lecture Notes in Mathematics*, 1721, Springer-Verlag, 1999. MR1735271 (2001d:14056)
- [RS00] K. Ranestad and F.-O. Schreyer. Varieties of sums of powers. *J. Reine Angew. Math.*, 525:147–181, 2000. MR1780430 (2001m:14009)
- [Rez92] B. Reznick. Sums of even powers of real linear forms. *Mem. Amer. Math. Soc.*, 96(463):viii+155, 1992. MR1096187 (93h:11043)
- [Rez10] B. Reznick. On the length of a binary form. Preprint arXiv:1007.5485.

DEPARTMENT OF MATHEMATICS, KTH, SE-100 44 STOCKHOLM, SWEDEN

E-mail address: `boij@kth.se`

DIPARTIMENTO DI MATEMATICA, POLITECNICO DI TORINO, 10129 TURIN, ITALY

E-mail address: `enrico.carlini@polito.it`

DEPARTMENT OF MATHEMATICS AND STATISTICS, QUEEN'S UNIVERSITY, KINGSTON, ONTARIO,
CANADA, K7L 3N6 – AND – DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI GENOVA, 16126
GENOA, ITALY

E-mail address: `Anthony.Geramita@gmail.com`

E-mail address: `geramita@dima.unige.it`