THE BRAUER GROUP OF MODULI SPACES OF VECTOR BUNDLES OVER A REAL CURVE

INDRANIL BISWAS, NORBERT HOFFMANN, AMIT HOGADI,
 AND ALEXANDER H. W. SCHMITT

(Communicated by Lev Borisov)

ABSTRACT. Let X be a geometrically connected smooth projective curve of genus $g_X \geq 2$ over $\mathbb{R}$. Let $M(r, \xi)$ be the coarse moduli space of geometrically stable vector bundles E over X of rank r and determinant ξ , where ξ is a real point of the Picard variety $\text{Pic}^d(X)$. If $g_X = r = 2$, then let d be odd. We compute the Brauer group of $M(r, \xi)$.

1. INTRODUCTION

Let $X_{\mathbb{C}}$ be a connected smooth projective curve of genus $g_X \geq 2$ over $\mathbb{C}$. Fix integers $r \geq 2$ and d . Given a line bundle $\xi_{\mathbb{C}}$ of degree d over $X_{\mathbb{C}}$, we denote by $M(r, \xi_{\mathbb{C}})$ the coarse moduli space of stable vector bundles over $X_{\mathbb{C}}$ of rank r and determinant $\xi_{\mathbb{C}}$.

The Picard group of such moduli spaces has been studied intensively; see for example [DN, KN, LS, So, BLS, Fa, Te, BHo1]. We view the Brauer group as a natural higher order analogue of the Picard group. It is related to the classical rationality problem [CS].

We assume that d is odd if $g_X = r = 2$; otherwise d is arbitrary. The Brauer group of $M(r, \xi_{\mathbb{C}})$ has been computed in [BBGN]; the result is a canonical isomorphism

$$\text{Br}(M(r, \xi_{\mathbb{C}})) \cong \mathbb{Z}/\gcd(r, d).$$

The corresponding generator $\beta_{\mathbb{C}} \in \text{Br}(M(r, \xi_{\mathbb{C}}))$ can be viewed as the obstruction against the existence of a Poincaré bundle, or universal vector bundle, over $M(r, \xi_{\mathbb{C}}) \times X_{\mathbb{C}}$.

Now suppose $X_{\mathbb{C}} = X \otimes_{\mathbb{R}} \mathbb{C}$ for a smooth projective curve X over $\mathbb{R}$. Then some of the above moduli spaces carry interesting real algebraic structures, and there has been a growing interest in understanding these structures [BhB, BHH, BHu, Sch]. In this paper we compute the Brauer group of such real algebraic moduli spaces.

More precisely, assume that the line bundle $\xi_{\mathbb{C}}$ comes from a real point ξ of the Picard variety $\text{Pic}^d(X)$. Let $M(r, \xi)$ be the coarse moduli space of geometrically stable vector bundles E over X of rank r and determinant ξ . It is a smooth quasiprojective variety over $\mathbb{R}$, with $M(r, \xi) \otimes_{\mathbb{R}} \mathbb{C} \cong M(r, \xi_{\mathbb{C}})$; see Section 2. Our main result, Theorem 3.3, describes the Brauer group of $M(r, \xi)$ as follows.

Received by the editors August 19, 2010 and, in revised form, October 13, 2010.

2010 *Mathematics Subject Classification*. Primary 14F22, 14D20, 14P99.

Key words and phrases. Brauer group, moduli space, real algebraic curve.

©2011 American Mathematical Society
 Reverts to public domain 28 years from publication

Theorem 1.1. *With $\chi := r(1 - g_X) + d$, there is a canonical isomorphism*

$$\mathrm{Br}(M(r, \xi)) \cong \begin{cases} \mathbb{Z}/\mathrm{gcd}(r, \chi) \oplus \mathbb{Z}/2 & \text{if } \xi \text{ comes from a line bundle defined over } \mathbb{R}, \\ \mathbb{Z}/\mathrm{gcd}(2r, \chi) & \text{otherwise.} \end{cases}$$

Note that $\mathrm{gcd}(r, \chi) = \mathrm{gcd}(r, d)$. The groups $\mathbb{Z}/\mathrm{gcd}(r, \chi)$ and $\mathbb{Z}/\mathrm{gcd}(2r, \chi)$ are generated by a canonical class $\beta \in \mathrm{Br}(M(r, \xi))$, the obstruction against a Poincaré bundle over $M(r, \xi) \times X$. The order of this obstruction class β is computed in Proposition 3.2. The remaining direct summand $\mathbb{Z}/2$ comes from the Brauer group of $\mathbb{R}$.

2. MODULI OF VECTOR BUNDLES OVER A REAL CURVE

Let X be a geometrically connected smooth projective algebraic curve of genus $g_X \geq 2$ defined over $\mathbb{R}$. We will denote the base change from $\mathbb{R}$ to $\mathbb{C}$ by a subscript $\mathbb{C}$. In particular, $X_{\mathbb{C}} := X \otimes_{\mathbb{R}} \mathbb{C}$ is the associated algebraic curve over $\mathbb{C}$.

Let $\sigma : \mathbb{C} \rightarrow \mathbb{C}$ denote complex conjugation. The involutive morphism of schemes

$$\sigma_X := \mathrm{id}_X \otimes \sigma : X_{\mathbb{C}} \rightarrow X_{\mathbb{C}}$$

lies over $\sigma : \mathbb{C} \rightarrow \mathbb{C}$. The closed points of $X_{\mathbb{C}}$ fixed by σ_X are the real points of X .

Let ξ be a real point of the Picard variety $\mathrm{Pic}(X)$. Viewing the associated complex point $\xi_{\mathbb{C}}$ of $\mathrm{Pic}(X_{\mathbb{C}})$ as a line bundle over $X_{\mathbb{C}}$, we have $\xi_{\mathbb{C}} \cong \sigma_X^*(\xi_{\mathbb{C}})$.

A *real* (respectively, *quaternionic*) structure on $\xi_{\mathbb{C}}$ is by definition an isomorphism

$$\eta : \xi_{\mathbb{C}} \rightarrow \sigma_X^*(\xi_{\mathbb{C}})$$

of line bundles over $X_{\mathbb{C}}$ with $\sigma_X^*\eta \circ \eta = \mathrm{id}_{\xi_{\mathbb{C}}}$ (respectively, $\sigma_X^*\eta \circ \eta = -\mathrm{id}_{\xi_{\mathbb{C}}}$). The line bundle $\xi_{\mathbb{C}}$ admits either a real structure η or a quaternionic structure η , and in both cases the resulting pair $(\xi_{\mathbb{C}}, \eta)$ is uniquely determined up to an isomorphism; cf. for example [Ve, Proposition 2.5] or [BHH, Proposition 3.1].

The real point ξ of $\mathrm{Pic}(X)$ is called *quaternionic* if $\xi_{\mathbb{C}}$ admits a quaternionic structure. Otherwise, $\xi_{\mathbb{C}}$ admits a real structure, so we can view ξ as a real line bundle over X .

A vector bundle E over X is called *geometrically stable* if the vector bundle $E_{\mathbb{C}}$ over $X_{\mathbb{C}}$ is stable. Not every stable vector bundle E over X is geometrically stable, but it is always geometrically polystable. Fix integers $r \geq 2$ and d . We denote by

$$(2.1) \quad \mathcal{M}(r, d) \supset \mathcal{M}(r, d)^s \rightarrow M(r, d)$$

the moduli stack of vector bundles E over X of rank r and degree d , the open substack of geometrically stable E , and their coarse moduli scheme, respectively. Since geometrically stable E have only scalar automorphisms, $\mathcal{M}(r, d)^s$ is a gerbe with band $\mathbb{G}_m$ over $M(r, d)$.

Let $\mathcal{L}(\det)$ denote the determinant of the cohomology line bundle over $\mathcal{M}(r, d)$. Its fiber over the moduli point of a vector bundle E is by definition $\det H^0(E) \otimes \det^{-1} H^1(E)$.

All three moduli spaces or stacks in (2.1) come with a determinant map to the Picard variety $\mathrm{Pic}^d(X)$. Given a real point ξ of $\mathrm{Pic}^d(X)$, we denote by

$$\mathcal{M}(r, \xi) \supset \mathcal{M}(r, \xi)^s \rightarrow M(r, \xi)$$

the corresponding fibers over ξ . So $M(r, \xi)$ is a smooth quasiprojective variety over $\mathbb{R}$, whose base change $M(r, \xi)_{\mathbb{C}}$ is the moduli space of stable vector bundles over $X_{\mathbb{C}}$ of rank r and determinant $\xi_{\mathbb{C}}$. By restriction, $\mathcal{M}(r, \xi)^s$ is a gerbe with band $\mathbb{G}_m$ over $M(r, \xi)$.

Suppose for the moment that ξ is a real line bundle. Then we can define a line bundle $\mathcal{L}(\xi)$ over $\mathcal{M}(r, \xi)$ whose fiber over the moduli point of a vector bundle E is $\text{Hom}(\xi, \det E)$. To state this more precisely, let S be a scheme over $\mathbb{R}$. Then the pullback of $\mathcal{L}(\xi)$ along the classifying morphism $S \rightarrow \mathcal{M}(r, \xi)$ of a vector bundle $\mathcal{E}$ over $X \times S$ is by definition the line bundle $\text{pr}_{2,*}(\text{pr}_1^* \xi^{-1} \otimes \det \mathcal{E})$ over S . This defines a line bundle $\mathcal{L}(\xi)$ over $\mathcal{M}(r, \xi)$.

Now suppose that ξ is quaternionic. Then the same recipe defines a line bundle over $\mathcal{M}(r, \xi)_{\mathbb{C}}$ endowed with a quaternionic structure. We denote this pair again by $\mathcal{L}(\xi)$.

In both cases, $\mathcal{L}(\xi)$ gives us a line bundle $\mathcal{L}(\xi)_{\mathbb{C}}$ over $\mathcal{M}(r, \xi)_{\mathbb{C}}$. If we trivialize the fiber of $\xi_{\mathbb{C}}$ over one closed point $x_0 \in X_{\mathbb{C}}$, we can identify $\mathcal{L}(\xi)_{\mathbb{C}}$ with the line bundle whose fiber at the moduli point of a vector bundle $E_{\mathbb{C}}$ over $X_{\mathbb{C}}$ is the fiber of $\det E_{\mathbb{C}}$ over x_0 .

Proposition 2.1. *The Picard group $\text{Pic}(\mathcal{M}(r, \xi))$ is generated by*

- i) $\mathcal{L}(\det)$ and $\mathcal{L}(\xi)$ if ξ is a real line bundle,
- ii) $\mathcal{L}(\det)$ and $\mathcal{L}(\xi)^{\otimes 2}$ if ξ is quaternionic.

The restrictions of these line bundles also generate $\text{Pic}(\mathcal{M}(r, \xi)^s)$.

Proof. Let $\widetilde{\mathcal{M}}(r, \xi_{\mathbb{C}})$ denote the moduli stack of vector bundles E of rank r over $X_{\mathbb{C}}$ together with an isomorphism $\xi_{\mathbb{C}} \cong \det E$. The forgetful map

$$\pi : \widetilde{\mathcal{M}}(r, \xi_{\mathbb{C}}) \rightarrow \mathcal{M}(r, \xi)_{\mathbb{C}}$$

is the $\mathbb{G}_m$ -torsor given by the line bundle $\mathcal{L}(\xi)_{\mathbb{C}}$. It is easy to check that the kernel of

$$\pi^* : \text{Pic}(\mathcal{M}(r, \xi)_{\mathbb{C}}) \rightarrow \text{Pic}(\widetilde{\mathcal{M}}(r, \xi_{\mathbb{C}}))$$

is generated by $\mathcal{L}(\xi)_{\mathbb{C}}$; cf. the proof of [BL, Lemma 7.8]. The Picard group of $\widetilde{\mathcal{M}}(r, \xi_{\mathbb{C}})$ is generated by $\pi^*(\mathcal{L}(\det)_{\mathbb{C}})$, according to [BL, Remark 7.11 and Proposition 9.2].

This shows that $\text{Pic}(\mathcal{M}(r, \xi)_{\mathbb{C}})$ is generated by $\mathcal{L}(\det)_{\mathbb{C}}$ and $\mathcal{L}(\xi)_{\mathbb{C}}$. We have just seen that all these line bundles admit a real or quaternionic structure. This real or quaternionic structure is unique, since $\Gamma(\mathcal{M}(r, \xi)_{\mathbb{C}}, \mathcal{O}^*) = \mathbb{C}^*$. It follows that $\text{Pic}(\mathcal{M}(r, \xi))$ is the subgroup of line bundles in $\text{Pic}(\mathcal{M}(r, \xi)_{\mathbb{C}})$ which are real, not quaternionic. Hence $\text{Pic}(\mathcal{M}(r, \xi))$ is generated by the line bundles as claimed.

As $\mathcal{M}(r, \xi)$ is smooth, the restriction map $\text{Pic}(\mathcal{M}(r, \xi)) \rightarrow \text{Pic}(\mathcal{M}(r, \xi)^s)$ is surjective; cf. for example [BHo2, Lemma 7.3]. So these line bundles also generate $\text{Pic}(\mathcal{M}(r, \xi)^s)$. $\square$

Now let $\mathcal{M} \rightarrow M$ be a gerbe with band $\mathbb{G}_m$ over an irreducible Noetherian scheme M . As a basic example, we have the gerbe $\mathcal{M}(r, d)^s \rightarrow M(r, d)$ in mind.

Definition 2.2. Let $\mathcal{L}$ be a line bundle over $\mathcal{M}$. Then the automorphism groups $\mathbb{G}_m$ in $\mathcal{M}$ act on the fibers of $\mathcal{L}$. These $\mathbb{G}_m$ act by the same power $w \in \mathbb{Z}$ on every fiber of $\mathcal{L}$, since $\mathcal{M}$ is connected. The integer w is called the *weight* of $\mathcal{L}$.

The weight of a quaternionic line bundle $\mathcal{L}$ is by definition the weight of the associated complex line bundle $\mathcal{L}_{\mathbb{C}}$. For example, the real or quaternionic line bundle $\mathcal{L}(\xi)$ over $\mathcal{M}(r, \xi)^s$ has weight r . The real line bundle $\mathcal{L}(\det)$ over $\mathcal{M}(r, d)^s$ has weight

$$\chi := r(1 - g_X) + d$$

according to Riemann-Roch. Consider the integers

$$\chi' := \chi / \gcd(r, \chi) \quad \text{and} \quad r' := r / \gcd(r, \chi).$$

The real or quaternionic line bundle

$$\mathcal{L}(\Theta) := \mathcal{L}(\det)^{\otimes -r'} \otimes \mathcal{L}(\xi)^{\otimes \chi'}$$

over $\mathcal{M}(r, \xi)^s$ has weight 0. Hence it descends to a real or quaternionic line bundle over $M(r, \xi)$, which we again denote by $\mathcal{L}(\Theta)$. The line bundle $\mathcal{L}(\Theta)_{\mathbb{C}}$ is ample on $M(r, \xi)_{\mathbb{C}}$, and it generates the Picard group $\text{Pic}(M(r, \xi)_{\mathbb{C}})$ according to [DN, Théorèmes A & B].

Proposition 2.3. *The Picard group $\text{Pic}(M(r, \xi))$ is generated by*

- i) $\mathcal{L}(\Theta)$ if ξ is a real line bundle or χ' is even,
- ii) $\mathcal{L}(\Theta)^{\otimes 2}$ if ξ is quaternionic and χ' is odd.

Proof. The line bundles over $M(r, \xi)$ are the line bundles of weight 0 over $\mathcal{M}(r, \xi)^s$. According to Proposition 2.1, these are of the form $\mathcal{L}(\det)^{\otimes a} \otimes \mathcal{L}(\xi)^{\otimes b}$ with $a\chi + br = 0$, where moreover b has to be even if ξ is quaternionic. $\square$

3. THE BRAUER GROUP

The Brauer group $\text{Br}(S)$ of a Noetherian scheme S is by definition the abelian group of Azumaya algebras over S up to Morita equivalence. It is a torsion group, and it embeds canonically into the étale cohomology group $H_{\text{ét}}^2(S, \mathbb{G}_m)$.

If S is smooth and quasiprojective over a field, then $H_{\text{ét}}^2(S, \mathbb{G}_m)$ is also a torsion group [Gr, Proposition 1.4], and the embedding of $\text{Br}(S)$ into $H_{\text{ét}}^2(S, \mathbb{G}_m)$ is an isomorphism [dJ].

Our aim is to compute the Brauer group of the real moduli space $M(r, \xi)$. Let

$$(3.1) \quad \beta \in H_{\text{ét}}^2(M(r, \xi), \mathbb{G}_m) = \text{Br}(M(r, \xi))$$

denote the class given by the gerbe $\mathcal{M}(r, \xi)^s \rightarrow M(r, \xi)$ with band $\mathbb{G}_m$. Since a section of this gerbe would yield a Poincaré bundle over $M(r, \xi) \times X$, we can view the class β as the obstruction against the existence of such a Poincaré bundle.

Remark 3.1. Choose an effective divisor $D \subset X$ defined over $\mathbb{R}$, for example a closed point in X . The Brauer class β over $M(r, \xi)$ can also be described by the Azumaya algebra with fibers $\text{End } H^0(D, E|_D)$, or by the projective bundle with fibers $\mathbb{P}H^0(D, E|_D)$.

We first compute the exponent of β , i.e., the order of β as an element in the torsion group $\text{Br}(M(r, \xi))$. This will in particular re-prove results of [BH_u, Section 5].

Proposition 3.2. *Let ξ be a real point of the Picard variety $\underline{\text{Pic}}^d(X)$.*

- i) *If ξ is a real line bundle, then $\beta \in \text{Br}(M(r, \xi))$ has exponent $\gcd(r, \chi)$.*
- ii) *If ξ is quaternionic, then $\beta \in \text{Br}(M(r, \xi))$ has exponent $\gcd(2r, \chi)$.*

Proof. An integer $n \in \mathbb{Z}$ annihilates the class $\beta \in H_{\text{ét}}^2(M(r, \xi), \mathbb{G}_m)$ of the gerbe $\mathcal{M}(r, \xi)^s$ if and only if there is a line bundle $\mathcal{L}$ over $\mathcal{M}(r, \xi)^s$ which has weight n ; see for example [Ho, Lemma 4.9]. Hence the claim follows from Proposition 2.1. $\square$

We denote by $\mathbb{Z} \cdot \beta \subseteq \text{Br}(M(r, \xi))$ the subgroup generated by the class β in (3.1). Let

$$(3.2) \quad f : M(r, \xi) \longrightarrow \text{Spec}(\mathbb{R})$$

be the structure morphism. Recall that $\text{Br}(\mathbb{R}) \cong \mathbb{Z}/2$, the nontrivial element being the class $[\mathbb{H}] \in \text{Br}(\mathbb{R})$ of the quaternion algebra $\mathbb{H} = \mathbb{R} \oplus \mathbb{R} \cdot i \oplus \mathbb{R} \cdot j \oplus \mathbb{R} \cdot k$.

Theorem 3.3. *Let ξ be a real point of $\underline{\text{Pic}}^d(X)$, with d odd if $g_X = r = 2$. We have*

$$\text{Br}(M(r, \xi)) = \begin{cases} \mathbb{Z} \cdot \beta \oplus f^*(\text{Br}(\mathbb{R})) \cong \mathbb{Z}/\gcd(r, \chi) \oplus \mathbb{Z}/2 & \text{if } \xi \text{ is a real line bundle,} \\ \mathbb{Z} \cdot \beta \cong \mathbb{Z}/\gcd(2r, \chi) & \text{if } \xi \text{ is quaternionic.} \end{cases}$$

Proof. The structure morphism f in (3.2) yields a Leray spectral sequence

$$(3.3) \quad E_2^{p,q} = H_{\text{ét}}^p(\mathbb{R}, R^q f_* \mathbb{G}_m) \Rightarrow H_{\text{ét}}^{p+q}(M(r, \xi), \mathbb{G}_m).$$

We have $R^1 f_* \mathbb{G}_m = \text{Pic}(M(r, \xi)_{\mathbb{C}}) \cong \mathbb{Z}$. The action of $\text{Gal}(\mathbb{C}/\mathbb{R}) = \mathbb{Z}/2$ on it is trivial, for example because it preserves ampleness. From this we deduce

$$E_2^{1,1} = H_{\text{ét}}^1(\mathbb{R}, \mathbb{Z}) = \text{Hom}(\mathbb{Z}/2, \mathbb{Z}) = 0.$$

Hence the spectral sequence (3.3) provides in particular an exact sequence

$$H_{\text{ét}}^1(M(r, \xi), \mathbb{G}_m) \longrightarrow E_2^{0,1} \longrightarrow E_2^{2,0} \longrightarrow H_{\text{ét}}^2(M(r, \xi), \mathbb{G}_m) \longrightarrow E_2^{0,2}.$$

Using $f_* \mathbb{G}_m = \mathbb{G}_m$ and $R^2 f_* \mathbb{G}_m = \text{Br}(M(r, \xi)_{\mathbb{C}})$, we thus obtain an exact sequence

$$\text{Pic}(M(r, \xi)) \xrightarrow{g^1} \text{Pic}(M(r, \xi)_{\mathbb{C}}) \longrightarrow \text{Br}(\mathbb{R}) \xrightarrow{f^*} \text{Br}(M(r, \xi)) \xrightarrow{g^2} \text{Br}(M(r, \xi)_{\mathbb{C}}),$$

where g^1 and g^2 are pullback maps along the projection $g : M(r, \xi)_{\mathbb{C}} \longrightarrow M(r, \xi)$. Note that g^2 is surjective, since $g^2(\beta) = \beta_{\mathbb{C}}$ generates $\text{Br}(M(r, \xi)_{\mathbb{C}})$ by [BBGN].

Suppose that ξ is a real line bundle. Then g^1 is surjective due to Proposition 2.3, so f^* is injective. Since β has the same exponent as its image $\beta_{\mathbb{C}}$ by Proposition 3.2, it follows that $\text{Br}(M(r, \xi))$ is the direct sum of its subgroups $\mathbb{Z} \cdot \beta$ and $f^*(\text{Br}(\mathbb{R}))$, as required.

Now suppose that ξ is quaternionic and that $\chi' = \chi/\gcd(r, \chi)$ is even. Then f^* is injective as before, but the exponent $\gcd(2r, \chi)$ of β is twice the exponent $\gcd(r, \chi)$ of its image $\beta_{\mathbb{C}}$. Hence $\gcd(r, \chi) \cdot \beta = f^*([\mathbb{H}])$, and the class β generates $\text{Br}(M(r, \xi))$.

Finally, suppose that ξ is quaternionic and that χ' is odd. Then the cokernel of g^1 has two elements according to Proposition 2.3, so f^* is the zero map, and g^2 is an isomorphism. In particular, the class β again generates $\text{Br}(M(r, \xi))$. $\square$

ACKNOWLEDGEMENTS

Part of this work was done while the first author was visiting Freie Universität Berlin, with support from the SFB 647: Space - Time - Matter. The second and the fourth authors were also supported by SFB 647.

REFERENCES

- [BBGN] V. Balaji, I. Biswas, O. Gabber and D. S. Nagaraj, Brauer obstruction for a universal vector bundle, *Comp. Rend. Math. Acad. Sci. Paris* **345** (2007), 265–268. MR2353678 (2008h:14031)
- [BL] A. Beauville and Y. Laszlo, Conformal blocks and generalized theta functions. *Commun. Math. Phys.* **164** (1994), 385–419. MR1289330 (95k:14011)
- [BLS] A. Beauville, Y. Laszlo and C. Sorger, The Picard group of the moduli of G -bundles on a curve, *Compositio Math.* **112** (1998), 183–216. MR1626025 (99i:14011)
- [BhB] U. N. Bhosle and I. Biswas, Stable real algebraic vector bundles over a Klein bottle. *Trans. Amer. Math. Soc.* **360** (2008), 4569–4595. MR2403697 (2009f:14063)
- [BHo1] I. Biswas and N. Hoffmann, The line bundles on moduli stacks of principal bundles on a curve, *Documenta Math.* **15** (2010), 35–72. MR2628848
- [BHo2] I. Biswas and N. Hoffmann, Poincaré families of G -bundles on a curve, preprint arXiv:1001.2123, available at www.arXiv.org, to appear in *Math. Ann.*
- [BHH] I. Biswas, J. Huisman and J. Hurtubise, The moduli space of stable vector bundles over a real algebraic curve, *Math. Ann.* **347** (2010), 201–233. MR2593289
- [BHu] I. Biswas and J. Hurtubise, Universal vector bundle over the reals, preprint arXiv:0909.0041, available at www.arXiv.org, to appear in *Trans. Amer. Math. Soc.*
- [CS] J.-L. Colliot-Thélène and J.-J. Sansuc, The rationality problem for fields of invariants under linear algebraic groups (with special regard to the Brauer group), in: V. Mehta, editor, *Algebraic groups and homogeneous spaces* (Mumbai, 2004), pages 113–186, 2007. MR2348904 (2008m:13010)
- [dJ] A. de Jong: A result of Gabber, preprint, <http://www.math.columbia.edu/~dejong/>
- [DN] J.-M. Drezet and M. S. Narasimhan, Groupe de Picard des variétés de modules de fibrés semi-stables sur les courbes algébriques, *Invent. Math.* **97** (1989), 53–94. MR999313 (90d:14008)
- [Fa] G. Faltings, Algebraic loop groups and moduli spaces of bundles, *J. Eur. Math. Soc. (JEMS)* **5** (2003), 41–68. MR1961134 (2003k:14011)
- [Gr] A. Grothendieck, Le groupe de Brauer II: Théorie cohomologique, *Dix Exposés sur la Cohomologie des Schémas*, 67–87. North-Holland, Amsterdam, 1968. MR0244270 (39:5586b)
- [Ho] N. Hoffmann, Rationality and Poincaré families for vector bundles with extra structure on a curve, *Int. Math. Res. Not.*, Article ID rnm010 (2007), 30 pp. MR2337034 (2009b:14066)
- [KN] S. Kumar and M. S. Narasimhan, Picard group of the moduli spaces of G -bundles, *Math. Ann.* **308** (1997), 155–173. MR1446205 (98d:14014)
- [LS] Y. Laszlo and C. Sorger, The line bundles on the moduli of parabolic G -bundles over curves and their sections, *Ann. Sci. École Norm. Sup. (4)* **30** (1997), 499–525. MR1456243 (98f:14007)
- [Sch] F. Schaffhauser, Moduli varieties of real and quaternionic vector bundles over a curve, preprint arXiv:1003.5285, available at www.arXiv.org
- [So] C. Sorger, On moduli of G -bundles of a curve for exceptional G , *Ann. Sci. École Norm. Sup. (4)* **32** (1999), 127–133. MR1670528 (99m:14031)
- [Te] C. Teleman, Borel-Weil-Bott theory on the moduli stack of G -bundles over a curve, *Invent. Math.* **134** (1998), 1–57. MR1646586 (2000b:14014)
- [Ve] J.-L. Verdier, Instantons. *Les équations de Yang-Mills*, Séminaire École Norm. Sup. 1977-1978, 105–134. Astérisque 71–72, Société Mathématique de France, Paris, 1980. MR589891 (84j:14026)

SCHOOL OF MATHEMATICS, TATA INSTITUTE OF FUNDAMENTAL RESEARCH, HOMI BHABHA
ROAD, BOMBAY 400005, INDIA

E-mail address: `indranil@math.tifr.res.in`

INSTITUT FÜR MATHEMATIK, FREIE UNIVERSITÄT, ARNIMALLEE 3, 14195 BERLIN, GERMANY

E-mail address: `norbert.hoffmann@fu-berlin.de`

SCHOOL OF MATHEMATICS, TATA INSTITUTE OF FUNDAMENTAL RESEARCH, HOMI BHABHA
ROAD, BOMBAY 400005, INDIA

E-mail address: `amit@math.tifr.res.in`

INSTITUT FÜR MATHEMATIK, FREIE UNIVERSITÄT, ARNIMALLEE 3, 14195 BERLIN, GERMANY

E-mail address: `alexander.schmitt@fu-berlin.de`