ON THE NORMS OF DISCRETE ANALOGUES OF CONVOLUTION OPERATORS

OLEG KOVRIZHKIN

(Communicated by Michael T. Lacey)

ABSTRACT. We consider a discrete analogue of convolution operator $T(f) = K * f$ from $L^p(\mathbb{R}^d)$ to $L^p(\mathbb{R}^d)$: $T_{dis}(g) = K_{dis} * g$ from $\ell^p(\mathbb{Z}^d)$ to $\ell^p(\mathbb{Z}^d)$ where $K_{dis} = K|_{\mathbb{Z}^d}$ and $\hat{K}$ is supported in the fundamental cube. We show that the estimate $\|T_{dis}\|_p \leq C^d \|T\|_p$ with $C > 1$ cannot be improved for a certain range of p .

We consider a convolution operator $T(f) = K * f$ from $L^p(\mathbb{R}^d)$ to $L^p(\mathbb{R}^d)$ with a distribution kernel K whose Fourier transform $\hat{K}(\xi)$ is supported in the fundamental cube $Q = \{\xi = (\xi_1, \xi_2, \dots, \xi_d) : -\frac{1}{2} < \xi_i \leq \frac{1}{2}, i = 1, 2, \dots, d\} = (-\frac{1}{2}, \frac{1}{2}]^d$. The Fourier transform $(T(f))^\wedge = \hat{K}(\xi)\hat{f}(\xi)$. It is well known that if $T(f) = K * f$ acts from L^p to L^p , then it also acts from $L^{p'}$ to $L^{p'}$ with the same norm. Thus we restrict ourselves to the case $1 \leq p \leq 2$. In particular, we can say that $T(f) = K * f$ is an operator from L^r to L^r where $p \leq r \leq 2$. We will be interested in the norms of T from L^p to L^p . If $p = 1$, then $\|T\|_1 = \|K\|_1$, and if $p = 2$, then $\|T\|_2 = \|\hat{K}\|_\infty$. Interpolating, we get $\|T\|_p \leq \|K\|_1^{\frac{2}{p}-1} \|\hat{K}\|_\infty^{2-\frac{2}{p}}$ for $1 < p < 2$.

As we mentioned before, T is also a bounded operator from L^2 to L^2 and therefore $\hat{K}(\xi)$ is bounded, which makes $K(x) = \int_{\mathbb{R}^d} \hat{K}(\xi) e^{i2\pi\xi x} d\xi$ an infinitely differentiable function extendable to an entire function of exponential type. We define $K_{dis} = K|_{\mathbb{Z}^d}$ and $T_{dis}(g) = K_{dis} * g$ as an operator on $\mathbb{Z}^d$:

$$T_{dis}(g)(n) = \sum_{m \in \mathbb{Z}^d} K(n-m)g(m).$$

Since $\hat{K}$ is supported in the fundamental cube Q , we can view $K(n)$ as its Fourier coefficients and thus $\hat{K}(\xi) = \sum_{n \in \mathbb{Z}^d} K(n) e^{i2\pi n \xi} \chi_Q(\xi)$; i.e., K_{dis} and K are determined by each other. We will be interested in norms of T_{dis} from $\ell^p(\mathbb{Z}^d)$ to $\ell^p(\mathbb{Z}^d)$. What was said above about $\|T\|_p$ is also true for $\|T_{dis}\|_p$. If $p = 1$, then $\|T_{dis}\|_1 = \|K_{dis}\|_1$, and if $p = 2$, then $\|T_{dis}\|_2 = \|\hat{K}\|_\infty$. Interpolating, we get $\|T_{dis}\|_p \leq \|K_{dis}\|_1^{\frac{2}{p}-1} \|\hat{K}\|_\infty^{2-\frac{2}{p}}$ for $1 < p < 2$. We will find the following estimate useful.

Received by the editors December 16, 2010 and, in revised form, January 4, 2011.

2010 *Mathematics Subject Classification.* Primary 42A99, 42B99.

Key words and phrases. Discrete analogues, convolutions.

This research was partially supported by NSF grant DMS 0201099.

Proposition 1.

$$(1) \quad \|T_{dis}\|_p \geq \|K_{dis}\|_p.$$

To show this, just take $g(n) = \delta_0(n)$.

The main question we are interested in is how $\|T_{dis}\|_p$ is controlled by $\|T\|_p$. As we said earlier, we restrict ourselves to the case $1 \leq p \leq 2$ since $\|T_{dis}\|_{p'} = \|T_{dis}\|_p$ and $\|T\|_{p'} = \|T\|_p$. The following result is due to Magyar, Stein and Wainger [1, Proposition 2.1]: *If T is a bounded operator from $L^p(\mathbb{R}^d)$ to $L^p(\mathbb{R}^d)$, then T_{dis} is a bounded operator from $\ell^p(\mathbb{Z}^d)$ to $\ell^p(\mathbb{Z}^d)$ and*

$$(2) \quad \|T_{dis}\|_p \leq C(d)\|T\|_p$$

where C depends only on the dimension d but not on p .

The actual statement is more general. Magyar, Stein and Wainger raised the following question in their paper (see Remark 1 after Proposition 2.1 in [1]): *Is it possible to take C independently of the dimension d and, in particular, if $C = 1$?*

We will answer this question negatively.

Theorem 1. *There is p_0 with $1 < p_0 < 2$ such that for all p with $1 \leq p < p_0$ the optimal $C(d)$ in (2) has a lower bound $C(d) \geq B^d$ where $B > 1$.*

Actually the proof of Proposition 2.1 in [1] shows that $C(d) \leq A^d$, and thus this estimate of $C(d)$ cannot be improved. If $p = 2$, then $\|T\|_2 = \|T_{dis}\|_2 = \|\hat{K}\|_\infty$. Therefore, $C(d) = 1$ for $p = 2$. If $K \geq 0$, then $\|T\|_p = \|T_{dis}\|_p = \|\hat{K}\|_\infty = \|K\|_1$. Thus $C(d) \geq 1$.

Proof of Theorem 1. First we consider the case when the dimension $d = 1$ and $p = 1$. We will construct a kernel K such that $\|T_{dis}\|_1 = \sum_{n \in \mathbb{Z}} |K(n)| > \int_{\mathbb{R}} |K(x)| dx = \|T\|_1$. Let $\phi(x)$ be a nonnegative Schwartz function whose Fourier transform $\hat{\phi}(\xi)$ is also nonnegative and supported in $[-\frac{1}{2}, \frac{1}{2}]$, for example, $\phi(x) = \psi(x) \cdot \bar{\psi}(x)$ where $\hat{\psi}(\xi)$ is nonnegative and supported in $[-\frac{1}{4}, \frac{1}{4}]$. Define $\hat{K}(\xi) = \hat{\phi}(2(\xi - \frac{1}{4})) + \hat{\phi}(2(\xi + \frac{1}{4}))$. Then $\hat{K}(\xi)$ is supported in $[-\frac{1}{2}, \frac{1}{2}]$ and

$$K(x) = \cos\left(\frac{\pi x}{2}\right) \phi\left(\frac{x}{2}\right).$$

Therefore

$$|K(n)| = \begin{cases} \phi(\frac{n}{2}) & \text{if } n \text{ is even,} \\ 0 & \text{if } n \text{ is odd.} \end{cases}$$

Hence,

$$\sum_{n \in \mathbb{Z}} |K(n)| = \sum_{m \in \mathbb{Z}} \phi(m) = \hat{\phi}(0).$$

Now we will estimate $\|K\|_1$:

$$\begin{aligned} \int |K(x)|dx &= \int \phi\left(\frac{x}{2}\right) \left|\cos \frac{\pi x}{2}\right| dx \\ &= 2 \int \phi(x) |\cos \pi x| dx \\ &> 2 \int \phi(x) \cos^2 \pi x dx \\ &= \int \phi(x) (1 + \cos 2\pi x) dx \\ &= \hat{\phi}(0) + \frac{1}{2}\hat{\phi}(1) + \frac{1}{2}\hat{\phi}(-1) = \hat{\phi}(0). \end{aligned}$$

Thus we obtain $\int |K(x)|dx > \sum_{n \in \mathbb{Z}} |K(n)|$. This is the opposite of what we need to get. Now consider a family of kernels $K_t(x) = K(x+t)$ where $0 \leq t \leq 1$. We have $\|K_t\|_1 = \|K\|_1$. On the other hand $\int_0^1 \sum_{n \in \mathbb{Z}} |K(n+t)| dt = \int |K(x)| dx$. Since for $t=0$ we have $\int |K(x)| dx > \sum_{n \in \mathbb{Z}} |K(n)|$, there exists t such that $\int |K(x)| dx < \sum_{n \in \mathbb{Z}} |K_t(n)|$. For this kernel K_t we have $\|T_{dis}\|_1 > \|T\|_1$. Basically, the idea is to show that $h(t) = \sum_{n \in \mathbb{Z}} |K(n+t)|$ is not a constant. This can also be done by calculating Fourier coefficients $\hat{h}(m) = |K|^\wedge(m)$ and showing that they do not vanish; take for example $K(x) = \left(\frac{\sin \frac{\pi x}{3}}{\pi x}\right)^3$.

Now we consider the case $p > 1$. Suppose T is a bounded operator from L^1 to L^1 . Then T_{dis} is also a bounded operator from ℓ^1 to ℓ^1 . As we said earlier, $\|T\|_\infty = \|T\|_1$ and $\|T_{dis}\|_\infty = \|T_{dis}\|_1$. Interpolating between $p=1$ and $p=\infty$, we have that T and T_{dis} are bounded operators from L^p to L^p and $\log \|T\|_p$ and $\log \|T_{dis}\|_p$ are convex functions. Thus $\|T\|_p$ and $\|T_{dis}\|_p$ are continuous functions for $p \geq 1$. Let K be a kernel such that $\|T_{dis}\|_1 > \|T\|_1$. Then $\|T_{dis}\|_p > \|T\|_p$ for some range of p : $1 \leq p < p_0$. We can prove this in a different way which also uses the continuous dependence of $\|K_{dis}\|_p$ on p . Recall from (1) that $\|T_{dis}\|_p \geq \|K_{dis}\|_p$ and $\|T\|_p \leq \|K\|_1$. If K is a kernel such that $\sum_{n \in \mathbb{Z}} |K(n)| > \int |K(x)| dx$, then

$$\begin{aligned} \|T_{dis}\|_p &\geq \left(\sum_{n \in \mathbb{Z}} |K(n)|^p \right)^{\frac{1}{p}} \\ &> \int |K(x)| dx \geq \|T\|_p \end{aligned}$$

for some range of p : $1 \leq p < p_0$.

Now we will do the case when the dimension $d > 1$. Let T be a convolution operator from $L^p(\mathbb{R})$ to $L^p(\mathbb{R})$ with kernel K . Define a convolution operator $\tilde{T}$ from $L^p(\mathbb{R}^d)$ to $L^p(\mathbb{R}^d)$ with kernel $\tilde{K}(x) = K(x_1)K(x_2) \cdots K(x_d)$. It is easy to show that $\|\tilde{T}\|_p = \|T\|_p^d$. The same is true for discrete analogues. Let K be such a

kernel that $\frac{\|T_{dis}\|_p}{\|T\|_p} = B > 1$. Note that $\tilde{K}_{dis} = (\tilde{K})_{dis}$. Then

$$\frac{\|(\tilde{T})_{dis}\|_p}{\|\tilde{T}\|_p} = \frac{\|T_{dis}\|_p^d}{\|T\|_p^d} = B^d.$$

It would be interesting to know if there are kernels K with $\hat{K}$ supported in $[-\frac{1}{2}, \frac{1}{2}]$ such that $\|T_{dis}\|_p > \|T\|_p$ for $1 \leq p < 2$, not just for $1 \leq p < p_0$.

REFERENCES

- [1] A. Magyar, E. M. Stein, S. Wainger, *Discrete analogues in harmonic analysis: Spherical averages*, Annals of Mathematics (2), 155, 2002, 189-208. MR1888798 (2003f:42028)

E-mail address: olegk@alum.mit.edu