ON MULTIVALUED FIXED-POINT FREE MAPS ON $\mathbb{R}^n$

RAUSHAN Z. BUZYAKOVA

(Communicated by Alexander N. Dranishnikov)

ABSTRACT. To formulate our results let f be a continuous multivalued map from $\mathbb{R}^n$ to $2^{\mathbb{R}^n}$ and k a natural number such that $|f(x)| \leq k$ for all x . We prove that f is fixed-point free if and only if its continuous extension $\tilde{f} : \beta\mathbb{R}^n \rightarrow 2^{\beta\mathbb{R}^n}$ is fixed-point free. If one wishes to stay within metric terms, the result can be formulated as follows: f is fixed-point free if and only if there exists a continuous fixed-point free extension $\tilde{f} : b\mathbb{R}^n \rightarrow 2^{b\mathbb{R}^n}$ for some metric compactification $b\mathbb{R}^n$ of $\mathbb{R}^n$. Using the classical notion of colorability, we prove that such an f is always colorable. Moreover, a number of colors sufficient to paint the graph can be expressed as a function of n and k only. The mentioned results also hold if the domain is replaced by any closed subspace of $\mathbb{R}^n$ without any changes in the range.

1. INTRODUCTION

A series of topological results about fixed-point free maps is motivated by these two classical set-theoretical statements (see, in particular, [3]):

S1. *If $f : X \rightarrow X$ is a fixed-point free map, then there exists a finite cover $\mathcal{F}$ of X such that $f(F)$ misses F for each $F \in \mathcal{F}$.*

S2. *Let $\mathcal{P}(X)$ be the set of all non-empty subsets of X and let $f : X \rightarrow \mathcal{P}(X)$ be a map with the property that $x \notin f(x)$. If there exists a natural number k such that $|f(x)| \leq k$ for all $x \in X$, then there exists a finite cover $\mathcal{F}$ of X such that F misses $\bigcup\{f(x) : x \in F\}$ for each $F \in \mathcal{F}$.*

One of the first results of a topological nature related to these statements was obtained by Katětov in [6] by translating the first statement into this form: *For a discrete space X , a map $f : X \rightarrow X$ is fixed-point free if and only if its continuous extension $\tilde{f} : \beta X \rightarrow \beta X$ is fixed-point free.* This topological version suggests that if one wants to have a similar criterion for a non-discrete space X one has to at least demand that elements of covers $\mathcal{F}$ in the statements under discussion be closed subsets of X . Thus, it is commonly accepted that when working with a topological space X and a continuous map f from a closed subspace A of X to X , any closed subset F of A that misses its image under f is called a *color* of f . If there exists a finite cover (“coloring”) of X by colors, f is said to be *colorable*. Katětov’s paper [6] and van Douwen’s work [4] have made a significant impact on topologists’ interest in the topic. A number of interesting results of a topological nature in the direction of the first statement have been published since the mentioned papers (see, in particular, [8, Section 3.2] for references). In this paper we consider one of the natural topological versions of the second statement for Euclidean space $\mathbb{R}^n$ and

Received by the editors November 22, 2010 and, in revised form, March 7, 2011.

2010 *Mathematics Subject Classification.* Primary 54H25, 58C30, 54B20.

Key words and phrases. Fixed point, hyperspace, multivalued function.

its hyperspace $2^{\mathbb{R}^n}$. In [2] it is proved that a continuous fixed-point free map from a closed subspace of $\mathbb{R}^n$ to $\mathbb{R}^n$ is colorable. In this paper we consider fixed-point free multivalued maps on $\mathbb{R}^n$ and its closed subspaces. To formulate our main results we first introduce the necessary terminology related to multivalued maps.

For a topological space X , we use the symbol 2^X to denote the space of all non-empty closed subsets of X endowed with the Vietoris topology and the symbol $\exp_k(X)$ to denote the subspace of 2^X that consists of only those $A \in 2^X$ for which $|A| \leq k$. A multivalued map $f : X \subset Z \rightarrow 2^Z$ is *fixed-point free* if $x \notin f(x)$ for every $x \in X$. A closed set $F \subset X$ is a *color* of a continuous map f from $X \subset Z$ to 2^Z if F misses $\bigcup\{f(x) : x \in F\}$. If X can be covered by finitely many colors of f , then f is said to be *colorable*. The main results of the paper are Theorems 2.7 and 2.8, which state that any continuous fixed-point free map from a closed subspace of $\mathbb{R}^n$ to $\exp_k(\mathbb{R}^n)$ is colorable and there exists a formula that computes a number of colors sufficient for painting in terms of n and k only. Using the main result we also show that a criterion similar to Katětov's holds for multivalued maps as well. Namely, we show that a continuous map f from a closed subspace X of $\mathbb{R}^n$ to $\exp_k(\mathbb{R}^n)$ is fixed-point free if and only if its continuous extension $\tilde{f} : \beta X \rightarrow \exp_k(\beta \mathbb{R}^n)$ is fixed-point free. To justify the requirement on sizes of $f(x)$ in our main results let us consider one simple example. Define f from $\omega \setminus \{0\}$ to the space of finite subsets of ω as follows: $f(n) = \{n+1, n+2, \dots, 2n\}$. The map f is continuous because ω and the space of finite subsets of ω are discrete. Since for every $n \in \omega \setminus \{0\}$, all elements of $\{n+1, n+2, \dots, 2n\}$ must be of a different color we conclude that f is not colorable. This example justifies our requirement in the main results that the set $\{|f(x)| : x \in X\}$ is bounded by a positive integer. Before we dive into the technical part of the paper we would like to outline a short transparent argument of the main results of the paper for a fixed-point free map $f : \mathbb{R} \rightarrow \exp_2(\mathbb{R})$. For this let $f_1(x) = \min f(x)$ and $f_2(x) = \max f(x)$. Since f is fixed-point free, the maps f_1 and f_2 are fixed-point free real-valued maps. By [1, Theorem 2.5], both functions are colorable. If one lets $\mathcal{F}_1$ and $\mathcal{F}_2$ be colorings of these maps, then it is easy to verify that the family $\{A \cap B : A \subset \mathcal{F}_1, B \subset \mathcal{F}_2\}$ is a coloring of f . If one wishes to extend the argument for the case $\exp_3(\mathbb{R})$ using a straightforward inductive approach, then one may find oneself dealing with open colors or with a single-valued map with the domain being a proper subset of the range. The existence of open colors in a single-valued case can be easily deduced from the definition of colorability when one deals with self-maps. However, if one works with maps from a subspace X of Y into Y , work needs to be done. Nevertheless, the presented argument can be extended for $\exp_k(\mathbb{R})$ for any k with some more work and suitable references. Although our argument for any n and k that we present in the paper may seem different from the one just described, a closer look will reveal that it carries the same idea hidden behind technical details naturally arising when dealing with higher dimensions. Throughout the paper we will follow standard notation and terminology as in [5].

2. RESULTS

For simplicity, but without loss of generality, most of our arguments related to $\mathbb{R}^n$ will be carried out for $\mathbb{R}^5$. This will free the letter “ n ” for other purposes. Since throughout our discussion we will juggle several spaces at a time we agree that by

$\bar{S}$ we denote the closure of S in $\mathbb{R}^k$ (where the value of k is always understood from the context) while $cl_X(P)$ will denote the closure of P in X .

A standard neighborhood in 2^X will be denoted as

$$\langle U_1, \dots, U_m \rangle = \{A \in 2^X : A \subset U_1 \cup \dots \cup U_m \text{ and } U_i \cap A \neq \emptyset \text{ for all } i = 1, \dots, m\},$$

where $U_1, \dots, U_m$ are open sets of X .

By $\exp_n(X)$ we denote the subspace $\{F \in 2^X : |F| \leq n\}$.

Definition 2.1. Let f be a continuous map from $X \subset Z$ to 2^Z . A closed set $F \subset X$ is a bright color of f if F misses $cl_Z[\bigcup\{f(x) : x \in F\}]$.

Proposition 2.2. Suppose Z is normal, X is closed in Z , $f : X \rightarrow 2^Z$ is continuous, and F is a bright color of f . Then there exists an open neighborhood U of F such that $cl_X(U)$ is a bright color of f .

Proof. Since F is a bright color of f and X is closed in Z , the sets F and $cl_Z(\bigcup\{f(x) : x \in F\})$ are disjoint closed sets in Z . Since Z is normal, there exist V an open neighborhood of $cl_Z[\bigcup\{f(x) : x \in F\}]$ in Z and W an open neighborhood of F in Z such that $cl_Z(W)$ misses $cl_Z(V)$. Consider the open set $\langle V \rangle$ in 2^Z . Since f is continuous and $f(F) \subset \langle V \rangle$ there exists an open neighborhood U of F in X such that $U \subset W$ and $f(cl_Z(U)) \subset \langle V \rangle$. Thus, $cl_Z(\bigcup\{f(x) : x \in cl_Z(U)\})$ is in $cl_Z(V)$. The latter misses $cl_Z(W)$ and therefore $cl_Z(U)$. Therefore, U is as desired. $\square$

Proposition 2.2 implies that if X is closed in $\mathbb{R}^k$ and $\mathcal{F}$ is an m -sized bright coloring of a continuous map $f : X \rightarrow \exp_n(\mathbb{R}^k)$, then there exists an m -sized open cover $\mathcal{U}$ of X the closures of whose elements are bright colors of f . We will use this observation throughout the paper without formally referencing it.

In the following discussion that leads to the main result we will restrict ourselves to closed subspaces of $\mathbb{R}^5$. This is done to avoid accumulation of too many variables. All arguments are valid if one replaces “5” with any natural number.

Definition 2.3. $S(5, n)$ denotes the following statement: there exists a smallest integer $K(5, n)$ such that every continuous fixed-point free map f from a closed subset $X \subset \mathbb{R}^5$ to $\exp_n(\mathbb{R}^5)$ is colorable by at most $K(5, n)$ many bright colors.

Lemma 2.4. Suppose $f, g, h : X \subset Z \rightarrow 2^Z$ are maps and g and h are colorable by at most N_g and N_h (bright) colors, respectively. If $f(x) \subset g(x) \cup h(x)$ for every $x \in X$, then f is colorable by at most $N_g \cdot N_h$ (bright) colors.

Proof. Let $\mathcal{G}$ and $\mathcal{H}$ be bright colorings of g and h of sizes N_g and N_h , respectively. Put $\mathcal{F} = \{G \cap H : G \cap H \neq \emptyset, H \in \mathcal{H}, G \in \mathcal{G}\}$. Clearly, $|\mathcal{F}| \leq N_g \cdot N_h$. Since $\mathcal{G}$ and $\mathcal{H}$ are closed covers of X , so is $\mathcal{F}$. Fix $F = H \cap G$ in $\mathcal{F}$. Since, by hypothesis, $f(x) \subset g(x) \cup h(x)$, we conclude that

$$cl_Z \left[\bigcup \{f(x) : x \in H \cap G\} \right] \subset cl_Z \left[\bigcup \{g(x) : x \in H \cap G\} \right] \cup cl_Z \left[\bigcup \{h(x) : x \in H \cap G\} \right].$$

Since $\mathcal{G}$ and $\mathcal{H}$ are bright colorings, $cl_Z[\bigcup\{g(x) : x \in H \cap G\}]$ and $cl_Z[\bigcup\{h(x) : x \in H \cap G\}]$ miss $H \cap G$. Therefore, the left side of the above set inclusion formula misses $H \cap G$ as well, meaning that $H \cap G$ is a bright color for f . $\square$

In what follows, by π_i we denote the projection of $\mathbb{R}^5$ onto its i -th coordinate axis. The next two statements (Lemmas 2.5 and 2.6) hold if we replace π_1 by π_i for any $i \in \{1, \dots, n\}$. However, we will carry out our arguments for π_1 for the already-mentioned reason of avoiding unnecessary variables.

Lemma 2.5. *Suppose $n > M \geq 1$, A is closed in $\mathbb{R}^5$, $f : A \rightarrow \exp_n(\mathbb{R}^5) \setminus \exp_{n-1}(\mathbb{R}^5)$ is continuous and fixed-point free, $|\{y \in f(x) : \pi_1(y) = \max \pi_1(f(x))\}| = M$ for all $x \in A$, and $S(5, n-1)$ is true. Then f is colorable by at most $[K(5, n-1)]^2$ bright colors.*

Proof. Define g and h from A to $\exp_{n-1}(\mathbb{R}^5)$ as follows:

$$g(x) = \{z \in f(x) : \pi_1(z) = \max \pi_1(f(x))\} \text{ and } h(x) = f(x) \setminus g(x).$$

Since $f(x)$ is finite, $\max \pi_1(f(x))$ exists. Hence $g(x)$ is defined. Since $|\{y \in f(x) : \pi_1(y) = \max \pi_1(f(x))\}| = M$ and $1 \leq M < n$, we conclude that $0 < |f(x) \setminus g(x)| < n$ and $0 < |g(x)| < n$. Therefore, g and h are well-defined functions from A to $\exp_{n-1}(\mathbb{R}^5)$. Observe that $f(x) = g(x) \cup h(x)$ for each x . Therefore, by Lemma 2.4, to reach the conclusion of our lemma we need to show that g and h are colorable by at most $K(5, n-1)$ bright colors each. Since we assume that $S(5, n-1)$ holds it suffices to show that g and h are continuous and fixed-point free. The latter property follows from the facts that f is fixed-point free and $f(x) = g(x) \cup h(x)$.

To prove continuity of g and h , fix $x \in A$ and open neighborhoods $\mathcal{U}_{g(x)}$ and $\mathcal{V}_{h(x)}$ of $g(x)$ and $h(x)$ in $\exp_{n-1}(\mathbb{R}^5)$. We need to find an open neighborhood of x in A whose images under $g(x)$ and $h(x)$ are in $\mathcal{U}_{g(x)}$ and $\mathcal{V}_{h(x)}$, respectively. Without loss of generality we may assume that the selected neighborhoods are standard, that is, in the form

$$\mathcal{U}_{g(x)} = \langle U_y : y \in g(x) \rangle \text{ and } \mathcal{V}_{h(x)} = \langle U_y : y \in f(x) \setminus g(x) \rangle,$$

where U_y is a fixed bounded open neighborhood of y in $\mathbb{R}^5$ for each $y \in f(x)$. We may also assume that the following properties hold:

P1: $\min \pi_1(\overline{U}_y) > \max \pi_1(\overline{U}_z)$ whenever $y \in g(x)$ and $z \in f(x) \setminus g(x)$.

P2: $\overline{U}_y \cap \overline{U}_z = \emptyset$ for any distinct $y, z \in f(x)$.

The property P1 can be achieved since by the definition of g , the set $\pi_1(g(x))$ is a singleton and its element is strictly greater than any element of $\pi_1(f(x) \setminus g(x))$. Put $\mathcal{W}_{f(x)} = \langle U_y : y \in f(x) \rangle$. Clearly, $\mathcal{W}_{f(x)}$ is an open neighborhood of $f(x)$. By continuity of f , there exists an open O of x in A such that $f(O) \subset \mathcal{W}_{f(x)}$. To finish the proof of continuity of g and h it suffices to show that $g(O) \subset \mathcal{U}_{g(x)}$ and $h(O) \subset \mathcal{V}_{h(x)}$. For this, fix an arbitrary $x' \in O$. By the choice of O , we have $f(x') \in \mathcal{W}_{f(x)}$. Let $f(x') = \{z_1, \dots, z_n\} \in \exp_n(\mathbb{R}^5) \setminus \exp_{n-1}(\mathbb{R}^5)$ and $\pi_1(z_i) = \max \pi_1(f(x'))$ for any $i = 1, \dots, M$. By the lemma's condition on M , we have $\pi_1(z_j) < \max \pi_1(f(x))$ for any $j = M+1, \dots, n$. By P1, we have

P3: $z_i \in \bigcup \{U_y : y \in g(x)\}$ for any $i = 1, \dots, M$.

By P2 and P3, we have

P4: $z_j \in \bigcup \{U_y : y \in f(x) \setminus g(x)\}$ for any $j = M+1, \dots, n$.

By P2 and the fact that $f(x') \in \mathcal{W}_{f(x)} = \langle U_y : y \in f(x) \rangle$, we conclude that each U_y , participating in the definition of $\mathcal{W}_{f(x)}$, contains exactly one $z_i \in f(x')$. By P3 and P4, we have $\{z_1, \dots, z_M\} \in \mathcal{U}_{g(x)}$ and $\{z_{M+1}, \dots, z_n\} \in \mathcal{V}_{h(x)}$. Since $g(x') = \{z_1, \dots, z_M\}$ and $h(x') = \{z_{M+1}, \dots, z_n\}$, we are done. $\square$

Lemma 2.6. *Suppose $n > 1$, A is closed in $\mathbb{R}^5$, $f : A \rightarrow \exp_n(\mathbb{R}^5) \setminus \exp_{n-1}(\mathbb{R}^5)$ is continuous and fixed-point free, $|\{y \in f(x) : \pi_1(y) = \max \pi_1(f(x))\}| > 1$ for all $x \in A$, and $S(5, n-1)$ is true. Then f is colorable by at most $n \cdot [K(5, n-1)]^2$ bright colors.*

Proof. For $m = 1, \dots, n-1$, define $O_m \subset A$ as follows: $x \in O_m$ if and only if $M_x = |\{y \in f(x) : \pi_1(y) = \max \pi_1(f(x))\}| < n - m$.

Claim. O_m is open. To prove the claim, fix $x \in O_m$ and let $\mathcal{V}_{f(x)} = \langle V_y : y \in f(x) \rangle$ be an open neighborhood of $f(x)$ such that the following hold:

- (1) V_y is a bounded neighborhood of y for every $y \in f(x)$;
- (2) $V_y \cap V_z = \emptyset$ for any distinct $y, z \in f(x)$;
- (3) $\min \pi_1(\overline{V}_y) > \max \pi_1(\overline{V}_z)$ whenever $\pi_1(y) = \max \pi_1(f(x))$ and $\pi_1(z) < \max \pi_1(f(x))$.

It suffices to show now that $f^{-1}(\mathcal{V}_{f(x)}) \subset O_m$. For this pick $x' \in f^{-1}(\mathcal{V}_{f(x)})$. We have $f(x') \in \mathcal{V}_{f(x)}$. By (2) and (3), $M_{x'} \leq M_x$. Hence, $M_{x'} < n - m$. By the definition of O_m , $x' \in O_m$. The claim is proved.

We will construct our coloring inductively. For short put $K = [K(5, n-1)]^2$.

Step 1. Put $A_1 = A \setminus O_1$. Thus, an element x of A is in A_1 if and only if $M_x \geq n - 1$. Since $|\pi_1(f(x))| > 1$ for every $x \in A$ we conclude that $M_x = n - 1$ for every $x \in A_1$. Therefore, by Lemma 2.5, there exists a K -sized open cover $\mathcal{U}_1$ of A_1 the closures of whose elements are bright colors for f .

Assumption. Assume for $m - 1$ that an open family $\mathcal{U}_{m-1}$ of size at most $(m - 1) \cdot K$ is defined and that the following conditions are met:

- P1: $\overline{U}$ is a bright color of f for every $U \in \mathcal{U}_{m-1}$;
- P2: If $M_x \geq n - (m - 1)$, then $x \in \bigcup \mathcal{U}_{m-1}$.

Step $m < n$. Put $A_m = A \setminus [O_m \cup (\bigcup \mathcal{U}_{m-1})]$. By the claim, the set A_m is closed. Pick any $x \in A_m$. Then $x \notin O_m$, meaning that $M_x \geq n - m$. Also $x \notin \bigcup \mathcal{U}_{m-1}$, which, by P2, implies that $M_x < n - (m - 1)$. Thus, $n - m \leq M_x < n - m + 1$. Therefore, $M_x = n - m$. Therefore, by Lemma 2.5, there exists a K -sized open cover $\mathcal{V}_m$ of A_m , the closures of whose elements are bright colors for f . Put $\mathcal{U}_m = \mathcal{U}_{m-1} \cup \mathcal{V}_m$. Clearly, the size of this family is at most $m \cdot K$. Let us verify P1 and P2 for m . Property P1 holds since $\mathcal{U}_m$ is the union of two families satisfying P1. For P2, observe that $M_x \geq n - m$ if and only if $x \notin O_m$. But $\mathcal{U}_m$ covers the complement of O_m .

The construction is complete. It suffices to show now that $A = \bigcup \mathcal{U}_{n-1}$. For this, pick $x \in A$. By the lemma's hypothesis, $M_x > 1 = n - (n - 1)$. By P2, $x \in \bigcup \mathcal{U}_{n-1}$. $\square$

The base step in the proof of our main theorem uses the following statement ([2, Proposition 2.9]): *If f is a continuous fixed-point free map from a closed subset X of $\mathbb{R}^m$ to $\mathbb{R}^m$, then f is colorable by at most $m + 3$ bright colors.*

Theorem 2.7. *There exists an integer $K(m, n)$ such that every continuous fixed-point free map from a closed subspace X of $\mathbb{R}^m$ into $\exp_n(\mathbb{R}^m)$ is colorable by at most $K(m, n)$ many bright colors.*

Proof. As in previous statements, to avoid accumulation of extra variables, let us deal with $m = 5$. Thus, we need to show that $K(5, n)$ exists for every natural number n . By [2, Proposition 2.9], $K(5, 1)$ exists. Assume that $K(5, n - 1)$ exists. To prove the conclusion of the theorem for n , fix any fixed-point free continuous

map f from a closed subspace X of $\mathbb{R}^5$ into $\exp_n(\mathbb{R}^5)$. Next define $L \subset X$ as follows: $x \in L$ if and only if $|f(x)| < n$. Then L is closed and the range of $f|_L$ is a subset of $\exp_{n-1}(\mathbb{R}^5)$ (notice that L can be empty). Therefore, by our inductive assumption, there exists a $K(5, n-1)$ -sized open cover $\mathcal{U}_L$ of L the closures of whose elements are bright colors. Put $E = X \setminus \bigcup \mathcal{U}_L$. Then E is closed and $|f(x)| = n$ for every $x \in E$.

For $1 \leq i \leq n$, define S_i as follows: $x \in S_i$ if and only if $x \in E$ and $|\pi_i(f(x))| = 1$. Notice that S_i can be empty. Clearly, S_i is closed. Inductively, we will first cover $\bigcup_{i \leq n} S_i = \{x \in E : |\pi_i(f(x))| = 1 \text{ for some } i\}$ by bright colors and then we will cover the rest of E .

Step 1. Put $E_i = \bigcap_{j \neq i} S_j$. Notice that E_i can be empty. Since $|f(x)| = n$ for every $x \in E$ we conclude that $|\pi_i(f(x))| = n$ for all $x \in E_i$. Since $n > 1$, by Lemma 2.5 (with π_1 replaced by π_i and $M = 1$), there exists a finite open cover $\mathcal{U}_1$ of $\bigcup_{i \leq n} E_i$ the closures of whose elements are bright colors for f .

Assumption. Suppose an open finite family $\mathcal{U}_{k-1}$, where $1 \leq k-1 < n$, is defined and the following hold:

P1: For every at most $(k-1)$ -sized subset I of $\{1, \dots, n\}$ the inclusion $\bigcap_{j \notin I} S_j \subset \bigcup \mathcal{U}_{k-1}$ holds;

P2: $\overline{U}$ is a bright color for every $U \in \mathcal{U}_{k-1}$.

Step $k < n$. For every k -sized $I \subset \{1, \dots, n\}$ put $E_I = [\bigcap_{j \notin I} S_j] \setminus [\bigcup \mathcal{U}_{k-1}]$. Pick $i^* \in I$. Then $|I \setminus \{i^*\}| = k-1$. By P1, the set $\bigcup \mathcal{U}_{k-1}$ contains $\bigcap \{S_j : j \in I \setminus \{i^*\}\}$. Since E_I misses $\bigcup \mathcal{U}_{k-1}$, we conclude that $|\pi_{i^*}(f(x))| > 1$ for every x in E_I . By Lemma 2.6, there exists a finite open cover $\mathcal{U}_I$ of E_I the closures of whose elements are bright colors. Put $\mathcal{U}_k = [\bigcup \{\mathcal{U}_I : I \subset \{1, \dots, n\}, |I| = k\}] \cup \mathcal{U}_{k-1}$. Properties P1 and P2 clearly hold for k . The construction is complete.

Let us show that $\{x \in E : |\pi_i(f(x))| = 1 \text{ for some } i\}$ is covered by $\mathcal{U}_{n-1}$. For this pick any z in this set. Put $I_z = \{i \leq n : |\pi_i(f(x))| > 1\}$. Clearly, $|I_z| < n$. Since $x \in E$ we conclude that $|f(x)| = n$. Therefore, $I_z \neq \emptyset$, and therefore $z \in \bigcap_{j \notin I_z} S_j$. By P1, $z \in \bigcup \mathcal{U}_{n-1}$.

To finish the proof we need to cover $E \setminus [\bigcup \mathcal{U}_{n-1}]$ by bright colors. For this observe that $E \setminus [\bigcup \mathcal{U}_{n-1}]$ contains only those x for which $|\pi_i(f(x))| = n > 1$ for all i . Therefore, by Lemma 2.5 (with $M = 1$), the set in question is covered by bright colors. Since we always used Lemmas 2.5 and 2.6 to construct our coloring, the size of the coloring depends on n and number 5 only. Thus, $K(5, n)$ exists. $\square$

A non-technical version of the above result is the following theorem, where n and k denote any natural numbers.

Theorem 2.8. *Any continuous fixed-point free map from a closed subspace X of $\mathbb{R}^k$ into $\exp_n(\mathbb{R}^k)$ is brightly colorable.*

Observe that if a continuous map $f : X \rightarrow 2^Z$ has the property that $f(x)$ is compact in Z for all x , then there exists a continuous extension $\tilde{f} : \beta X \rightarrow 2^{\beta Z}$. For our next discussion put $\mathcal{K}(X) = \{A \in 2^X : A \text{ is compact}\}$. In [2] it is proved that a continuous map f from a closed subspace X of $\mathbb{R}^k$ to $\mathbb{R}^k$ is fixed-point free if and only if its continuous extension $\tilde{f} : \beta X \rightarrow \beta \mathbb{R}^k$ is fixed-point free. It is natural to ask if the corresponding statement holds for a multivalued map $f : X \rightarrow \exp_n(\mathbb{R}^k)$ and its continuous extension $\tilde{f} : \beta X \rightarrow \exp_n(\beta \mathbb{R}^k)$. Observe first

that the continuous extension exists since $\exp_n(\beta\mathbb{R}^k)$ is compact. The affirmative answer to this question follows from the next statement.

Proposition 2.9. *Let X be a closed subspace of a normal space Z . If $\mathcal{F}$ is a bright coloring of a continuous map $f : X \rightarrow \mathcal{K}(Z)$, then $\{\beta F : F \in \mathcal{F}\}$ is a bright coloring of $\tilde{f} : \beta X \rightarrow \mathcal{K}(\beta Z)$.*

Proof. Since X is normal and $\mathcal{F}$ is a finite closed cover of X , the family $\{\beta F : F \in \mathcal{F}\}$ is a closed cover of βX . Therefore, we only need to show that βF is a bright color for $\tilde{f}$. Since F is a bright color for f the set F misses $cl_Z(\bigcup\{f(x) : x \in F\})$. Since F and $cl_Z(\bigcup\{f(x) : x \in F\})$ are disjoint closed subsets of the normal space Z we conclude that βF misses $cl_{\beta Z}(\bigcup\{f(x) : x \in F\})$. Since $\tilde{f}$ is a continuous extension of f we conclude that $cl_{\beta Z}(\bigcup\{\tilde{f}(x) : x \in \beta F\}) = cl_{\beta Z}(\bigcup\{f(x) : x \in F\})$. Thus βF misses $cl_{\beta Z}(\bigcup\{\tilde{f}(x) : x \in \beta F\})$, whence βF is a bright color of $\tilde{f}$. $\square$

Theorem 2.10. *Let f be a continuous map from a closed subspace X of $\mathbb{R}^k$ to $\exp_n(\mathbb{R}^k)$. Then f is fixed-point free if and only if $\tilde{f} : \beta X \rightarrow \exp_n(\beta\mathbb{R}^k)$ is fixed-point free.*

Proof. Sufficiency is obvious. To prove necessity, let $\mathcal{F}$ be a bright coloring of f . By Proposition 2.9, $\{\beta F : F \in \mathcal{F}\}$ is a coloring for $\tilde{f}$. Since $\{\beta F : F \in \mathcal{F}\}$ covers βX and $f(\beta F)$ misses βF for each $F \in \mathcal{F}$, we conclude that $\tilde{f}$ does not fix any point. $\square$

Using spectral techniques it is observed in [8, Corollary 3.5.7] that if f is a continuous colorable self-map on a separable metric space X , then one can find a continuous fixed-point free extension $\tilde{f} : bX \rightarrow bX$, where bX is a metric compactification of X of the same dimensionality as that of X . Using the same technique we will next outline a proof for the following metric version of Theorem 2.10 for natural numbers n and k .

Theorem 2.11. *Let $f : \mathbb{R}^k \rightarrow \exp_n(\mathbb{R}^k)$ be continuous. Then f is fixed-point free if and only if there exists a continuous fixed-point free extension $\tilde{f} : b\mathbb{R}^k \rightarrow \exp_n(b\mathbb{R}^k)$, where $b\mathbb{R}^k$ is some metric compactification of $\mathbb{R}^k$ of dimension k .*

Proof. Sufficiency is obvious. Let us outline a proof of necessity. By Theorem 2.10, $\tilde{f} : \beta\mathbb{R}^k \rightarrow \exp_n(\beta\mathbb{R}^k) \subset 2^{\beta\mathbb{R}^k}$ is fixed-point free. By the Ščepin spectral theorem [7], we can find spectra $\{b_\alpha(\mathbb{R}^k), \pi_\alpha^\gamma, A\}$ and $\{2^{b_\alpha(\mathbb{R}^k)}, p_\alpha^\gamma, A\}$ with inverse limits $\beta\mathbb{R}^k$ and $2^{\beta\mathbb{R}^k}$, respectively, and a family of maps $\{f_\alpha : \alpha \in A\}$ such that the following hold:

- (1) $b_\alpha(\mathbb{R}^k)$ is a metric compactification of $\mathbb{R}^k$ of dimension k for all α .
- (2) π_α^γ and p_α^γ are identity maps on $\mathbb{R}^k$ and $2^{\mathbb{R}^k}$, respectively.
- (3) $f_\alpha \circ \pi_\alpha = p_\alpha \circ f$.

Since $\tilde{f}$ is fixed-point free and $b_\alpha(\mathbb{R}^k)$ is compact for every α we may assume that f_α is fixed-point free for every α . By (2) and (3), each f_α coincides with f on $\mathbb{R}^k$. Therefore, each $\{f_\alpha, b_\alpha(\mathbb{R}^k)\}$ serves our purpose. $\square$

We would like to finish the paper by commenting on a number of colors sufficient to paint a given graph. If one follows the argument of Theorem 2.7 or the argument for the reals outlined in the introduction, one will quickly see that the size of the coloring constructed for the case $\exp_n(\mathbb{R}^k)$ is at least k^n . Therefore, it is natural

to wonder if an estimate for the required number of colors can be represented as a polynomial of both n and k .

REFERENCES

- [1] R. Buzyakova, *Fixed Point Free Maps on the Reals and More*, Top. and its Appl. **156** (2008), 465-472. MR2475129 (2009i:54065)
- [2] R. Buzyakova and A. Chigogidze, *Fixed-point free maps of Euclidean spaces*, Fundamenta Mathematica **212** (2011), 1-16. MR2771585
- [3] N. G. de Bruijn and P. Erdős, *A color problem for infinite graphs and a problem in the theory of relations*, Indagationes Math. **13** (1951), 371-373. MR0046630 (13:763g)
- [4] E. van Douwen, *βX and fixed-point free maps*, Topology Appl. **51** (1993), 191-195. MR1229715 (94g:54030)
- [5] R. Engelking, *General Topology*, PWN, Warszawa, 1977. MR0500780 (58:18316b)
- [6] M. Katětov, *A theorem on mappings*, Comment Math. Univ. Carolinae **8** (1967), 431-433. MR0229228 (37:4802)
- [7] E. V. Ščepin, *Topology of limit spaces with uncountable inverse spectra* (Russian), Uspekhi Mat. Nauk **31** (1976), 191-226. MR0464137 (57:4072)
- [8] J. van Mill, *The infinite-dimensional topology of function spaces*, Elsevier, Amsterdam, 2001. MR1851014 (2002h:57031)

DEPARTMENT OF MATHEMATICS AND STATISTICS, THE UNIVERSITY OF NORTH CAROLINA AT GREENSBORO, GREENSBORO, NORTH CAROLINA 27402

E-mail address: `rzbouzia@uncg.edu`