ON A DIFFERENCE BETWEEN QUANTITATIVE WEAK SEQUENTIAL COMPLETENESS AND THE QUANTITATIVE SCHUR PROPERTY

O. F. K. KALENDA AND J. SPURNÝ

(Communicated by Thomas Schlumprecht)

ABSTRACT. We study quantitative versions of the Schur property and weak sequential completeness, proceeding with investigations started by G. Godefroy, N. Kalton and D. Li and continued by H. Pfitzner and the authors. We show that the Schur property of ℓ_1 holds quantitatively in the strongest possible way and construct an example of a Banach space which is quantitatively weakly sequentially complete, has the Schur property, but fails the quantitative form of the Schur property.

1. INTRODUCTION

We recall that a Banach space is *weakly sequentially complete* if every weakly Cauchy sequence is weakly convergent. Further, a Banach space has the *Schur property* if any weakly convergent sequence is norm convergent. This is easily seen to be equivalent to the fact that any weakly Cauchy sequence is norm Cauchy (and hence norm convergent), and thus any space with the Schur property is weakly sequentially complete. A classical example of a Banach space with the Schur property is the space ℓ_1 of all absolutely summable sequences. An example of a weakly sequentially complete Banach space without the Schur property is the Lebesgue space $L_1(0, 1)$.

In the present paper we study a quantitative version of the Schur property and its relation to quantitative weak sequential completeness. A quantitative version of weak sequential completeness was studied in [5, 3, 8]. The existence of a quantitative Schur property was pointed out to us by M. Fabian, who observed in 2005 that the space ℓ_1 has, in the terminology defined below, the 5-Schur property. The referee pointed out to us that in the paper [4] the authors use the notion of “1-strong Schur property” which is another quantification of the Schur property. We explain the relation to our notions in the last section. Before stating the known results and our contribution we shall define the quantitative properties.

Received by the editors March 15, 2011 and, in revised form, April 5, 2011.

2010 *Mathematics Subject Classification.* Primary 46B20, 46B25.

Key words and phrases. Weakly sequentially complete Banach space, Schur property, quantitative versions of weak sequential completeness, quantitative versions of the Schur property, L -embedded Banach space.

The authors were supported by the Research Project MSM 0021620839 from the Czech Ministry of Education.

The first author was additionally supported in part by grant GAAV IAA 100190901.

The second author was partly supported by grant GAČR 201/07/0388.

©2012 American Mathematical Society
 Reverts to public domain 28 years from publication

For a bounded sequence (x_k) in a Banach space X , we write $\text{clust}_{X^{**}}(x_k)$ for the set of all weak* cluster points of (x_k) in X^{**} , and by $\delta(x_k)$ we denote its diameter. This quantity measures in a way how far the sequence is from being weakly Cauchy. Similarly, the quantity

$$\text{ca}(x_k) = \inf_{n \in \mathbb{N}} \text{diam}\{x_k : k \geq n\}$$

measures how far the sequence is from being norm Cauchy.

Further, if A, B are nonempty subsets of a Banach space X , then

$$d(A, B) = \inf\{\|a - b\| : a \in A, b \in B\}$$

denotes the usual distance between A and B , and the Hausdorff nonsymmetrized distance from A to B is defined by

$$\hat{d}(A, B) = \sup\{d(a, B) : a \in A\}.$$

We will say that a Banach space X has the C -Schur property (where $C \geq 0$) if

$$\text{ca}(x_k) \leq C\delta(x_k)$$

for each bounded sequence (x_k) in X . Similarly, X is said to be C -weakly sequentially complete if

$$\hat{d}(\text{clust}_{X^{**}}(x_k), X) \leq C\delta(x_k)$$

for each bounded sequence (x_k) in X . It is clear that any space with the C -Schur property has the Schur property and that any C -weakly sequentially complete space is weakly sequentially complete. It follows from Proposition 1.1 below that the C -Schur property implies C -weak sequential completeness.

It is proved in [5, Lemma IV.7] that any L -embedded Banach space is 1-weakly sequentially complete. We recall that a Banach space X is called L -embedded if there exists a projection $P : X^{**} \rightarrow X$ such that

$$\|Px^{**}\| + \|(I - P)x^{**}\| = \|x^{**}\|, \quad x^{**} \in X^{**}.$$

This result was mentioned in [3, p. 829] together with the question of which weakly sequentially complete Banach spaces satisfy a quantitative version of this property. Recently some new results were obtained in [8]. It is proved there that any L -embedded space is $\frac{1}{2}$ -weakly sequentially complete, where the constant $\frac{1}{2}$ is optimal as witnessed by the space ℓ_1 . The quoted paper further contains an example of a Schur space which is not C -weakly sequentially complete for any $C \geq 0$. Conversely, C -weak sequential completeness does not imply the Schur property (consider for example reflexive spaces or the space $L^1(0, 1)$).

Inspired by the above-mentioned results and a remark of M. Fabian, we investigate in this paper a quantification of the Schur property. We will assume that our Banach spaces are real. However, although some methods work only in real spaces, almost all the results are valid for complex spaces as well. We will discuss this in the final section.

The first proposition contains two easy inequalities and their consequences on the relationship of the quantitative Schur property and quantitative weak sequential completeness.

Proposition 1.1. *Let (x_k) be a bounded sequence in a Banach space X . Then the following inequalities hold:*

- (1) $\widehat{d}(\text{clust}_{X^{**}}(x_k), X) \leq \text{ca}(x_k),$
- (2) $\delta(x_k) \leq \text{ca}(x_k).$

In particular, if X has the C -Schur property, then it is C -weakly sequentially complete.

Proof. We first observe that by the weak* lower semicontinuity of the norm we have

$$\text{diam}\{x_k : k \geq n\} = \overline{\text{diam}\{x_k : k \geq n\}}^{w*},$$

and thus

$$d(x^{**}, X) \leq \overline{\text{diam}\{x_k : k \geq n\}}^{w*}, \quad x^{**} \in \text{clust}_{X^{**}}(x_k), n \in \mathbb{N}.$$

From this we deduce that

$$\text{ca}(x_k) = \inf_{n \in \mathbb{N}} \text{diam}\{x_k : k \geq n\} = \inf_{n \in \mathbb{N}} \overline{\text{diam}\{x_k : k \geq n\}}^{w*} \geq \widehat{d}(\text{clust}_{X^{**}}(x_k), X)$$

and

$$\text{ca}(x_k) \geq \text{diam} \bigcap_{n=1}^{\infty} \overline{\{x_k : k \geq n\}}^{w*} = \delta(x_k).$$

This proves (1) and (2). The remaining statement follows immediately from (1). $\square$

Inequalities (2) and (1) are optimal. Indeed, let $X = c_0$ and (y_k) be the summing basis (i.e., $y_k = (1, \dots, 1, 0, \dots)$, where the last ‘1’ is on the k -th place). Denote by (x_k) the sequence

$$y_1, 0, y_2, 0, y_3, 0, \dots$$

Then clearly $\delta(x_k) = \text{ca}(x_k) = \widehat{d}(\text{clust}_{X^{**}}(x_k), X) = 1$.

Inequality (2) shows in particular that a nontrivial space cannot have the C -Schur property for any $C < 1$. Another trivial consequence is that in spaces with the C -Schur property the quantities δ and ca are equivalent.

Let us remark that the situation with weak sequential completeness is different: any L -embedded space is $\frac{1}{2}$ -weakly sequentially complete by [8] and reflexive spaces are trivially 0-weakly sequentially complete. The constant $\frac{1}{2}$ is optimal by the following proposition.

Proposition 1.2. *Let X be a Banach space which is C -weakly sequentially complete for some $C < \frac{1}{2}$. Then X is reflexive.*

Moreover, even a stronger version holds: Suppose that there is $C < \frac{1}{2}$ such that

$$d(\text{clust}_{X^{**}}(x_k), X) \leq C\delta(x_k)$$

for each bounded sequence (x_k) in X . Then X is reflexive.

Proof. It is clear that the first part follows from the stronger version, so let us show the stronger statement. Let X be a nonreflexive Banach space. Let $\varepsilon > 0$ be arbitrary. It follows from [6, Theorem 1] that there is $x^* \in X^*$ such that any $x^{**} \in B_{X^{**}}$ such that $x^{**}(x^*) = \|x^*\|$ satisfies $d(x^{**}, X) \geq 1 - \varepsilon$.

Indeed, suppose that for any $y^* \in X^*$ there is some $x^{**} = x_{y^*}^{**} \in B_{X^{**}}$ such that $x^{**}(y^*) = \|y^*\|$ and $d(x^{**}, X) < 1 - \varepsilon$. Let $B = \{x_{y^*}^{**} : y^* \in X^*\}$. Then B is a boundary for X^* and $\widehat{d}(B, X) < 1$, which contradicts [6, Theorem 1].

So, let x^* be as in the first paragraph. Let (x_k) be a sequence in B_X such that $x^*(x_k) \rightarrow \|x^*\|$. Then clearly $\delta(x_k) \leq 2$. Moreover, $d(\text{clust}_{X^{**}}(x_k), X) \geq 1 - \varepsilon$, as any $x^{**} \in \text{clust}_{X^{**}}(x_k)$ satisfies $x^{**}(x^*) = \|x^*\|$. This completes the proof. $\square$

Let us remark that the converse of the previous proposition is trivially valid as well, as any reflexive space is 0-weakly sequentially complete.

We continue by our first main result, which is an improvement of Schur's theorem for ℓ_1 .

Theorem 1.3. *The space ℓ_1 has the 1-Schur property; i.e.,*

$$\text{ca}(x_k) = \delta(x_k)$$

for any bounded sequence (x_k) in ℓ_1 .

It follows from [8] that there is a Banach space with the Schur property which fails its quantitative version. Indeed, a space is constructed there with the Schur property which is not C -weakly sequentially complete for any $C \geq 0$. It follows from Proposition 1.1 that the same space fails the C -Schur property for each $C \geq 0$. Our second result shows that a quantitative version of the Schur property may fail even for an L -embedded space (which is $\frac{1}{2}$ -weakly sequentially complete).

Example 1.4. There exists a separable L -embedded Banach space with the Schur property which fails the C -Schur property for every $C \geq 0$.

Following the suggestion of the referee, we include the following diagram of implications between the classes of Banach spaces we study in this paper ("wsc" is an abbreviation of "weakly sequentially complete"):

$$\begin{array}{ccccccc} & & & \text{Schur} & & & \\ & & \nearrow & & \searrow & & \\ C\text{-Schur} & \Rightarrow & C\text{-wsc and Schur} & \Rightarrow & C\text{-wsc} & \Rightarrow & \text{wsc} \end{array}$$

All these implications are easy to check. If a constant C is included both in the assumption and in the conclusion, the constant in the conclusion has the same value.

None of the displayed implications can be reversed, even if we allow the respective constants to change. For the first implication from the left, it follows from Example 1.4. A reasoning for the implication $\nearrow$ and the third one in the bottom line is a result of [8]. For the remaining two implications one can use the classical space $L^1(0, 1)$.

2. PROOF OF THEOREM 1.3

We start the proof with a simple fact based upon the standard "sliding hump" argument. If $x \in \ell_1$ and $N \subset \mathbb{N}$, we denote by $x|_N$ the element arising from x by setting its coordinates outside N to zero, i.e.,

$$x|_N(n) = \begin{cases} x(n), & n \in N, \\ 0, & n \in \mathbb{N} \setminus N. \end{cases}$$

Lemma 2.1. *Let (y_n) be a sequence in ℓ_1 converging pointwise to y with $\|y\| < \varepsilon$. Then there exist a subsequence (y_{n_k}) and indices $0 = N_0 < N_1 < N_2 < \dots$ such that*

$$\|y_{n_k}|_{(N_{k-1}, N_k]}\| > \|y_{n_k}\| - \varepsilon, \quad k \in \mathbb{N}.$$

Proof. First we set $N_0 = 0$, $n_1 = 1$ and choose $N_1 > 0$ such that $\|y_{n_1}|_{(0, N_1]}\| > \|y_{n_1}\| - \varepsilon$. Suppose now that $k \in \mathbb{N}$ is such that we have already constructed n_j for $1 \leq j \leq k$ and N_j for $0 \leq j \leq k$.

Since (y_n) converges pointwise to y , we can select $n_{k+1} > n_k$ so large that $\|y_n|_{(N_0, N_k]}\| < \varepsilon$ for every $n \geq n_{k+1}$. Thus, in particular,

$$\|y_{n_{k+1}}|_{(N_k, +\infty)}\| > \|y_{n_{k+1}}\| - \varepsilon,$$

so we can find $N_{k+1} > N_k$ such that

$$\|y_{n_{k+1}}|_{(N_k, N_{k+1}]}\| > \|y_{n_{k+1}}\| - \varepsilon.$$

This completes the construction. $\square$

Now we proceed with the proof of Theorem 1.3. Let (x_k) be a bounded sequence in $X = \ell_1$ and $\varepsilon > 0$. We consider an arbitrary $c < \text{ca}(x_k)$.

We extract subsequences (a_n) and (b_n) from (x_k) such that $c < \|a_n - b_n\|$ for $n \in \mathbb{N}$. Denoting $y_n = a_n - b_n$, we pass to a subsequence if necessary and assume thus that (y_n) pointwise converges to $y \in \ell_1$. Let $m \in \mathbb{N}$ be chosen such that $\|y|_{(m, +\infty)}\| < \varepsilon$. By omitting finitely many elements of (y_n) we achieve that

$$(3) \quad \|(y_n - y)|_{[1, m]}\| < \varepsilon, \quad n \in \mathbb{N}.$$

Hence

$$(4) \quad c < \|y_n|_{[1, m]}\| + \|y_n|_{(m, +\infty)}\| \leq \|y|_{[1, m]}\| + \varepsilon + \|y_n|_{(m, +\infty)}\|, \quad n \in \mathbb{N}.$$

Using Lemma 2.1 applied to $(y_n|_{(m, +\infty)})$ we obtain a subsequence (y_{n_k}) and indices $m = N_0 < N_1 < \dots$ such that

$$(5) \quad \|y_{n_k}|_{(N_{k-1}, N_k]}\| > \|y_{n_k}|_{(m, +\infty)}\| - \varepsilon, \quad k \in \mathbb{N}.$$

Let $x^* \in \ell_\infty = \ell_1^*$ be defined as

$$x^*(j) = \begin{cases} \text{sign } y(j), & j \in [1, m], \\ \text{sign } y_{n_k}(j), & j \in (N_{k-1}, N_k], \quad k \in \mathbb{N}. \end{cases}$$

Then $\|x^*\| \leq 1$, and, for each $k \in \mathbb{N}$, we have using (3), (5) and (4),

$$\begin{aligned} x^*(y_{n_k}) &= x^*(y_{n_k}|_{[1, m]}) + x^*(y_{n_k}|_{(N_{k-1}, N_k]}) + \sum_{j \in (m, +\infty) \setminus (N_{k-1}, N_k]} x^*(j)y_{n_k}(j) \\ &> x^*(y|_{[1, m]}) - \varepsilon + \|y_{n_k}|_{(N_{k-1}, N_k]}\| - \|y_{n_k}|_{(m, +\infty) \setminus (N_{k-1}, N_k]}\| \\ &> \|y|_{[1, m]}\| + \|y_{n_k}|_{(m, +\infty)}\| - 3\varepsilon \\ &> c - 4\varepsilon. \end{aligned}$$

Therefore we have

$$x^*(a_{n_k} - b_{n_k}) \geq c - 4\varepsilon.$$

Up to a passing to a subsequence we can suppose that the sequence $(x^*(a_{n_k}))$ converges. Up to passing to a further subsequence we can assume that the sequence $(x^*(b_{n_k}))$ converges as well. Let a^{**} and b^{**} be weak* cluster points of (a_{n_k}) and (b_{n_k}) , respectively. Then $a^{**}, b^{**} \in \text{clust}_{X^{**}}(x_k)$ and

$$\|a^{**} - b^{**}\| \geq (a^{**} - b^{**})(x^*) = \lim_{k \rightarrow \infty} x^*(a_{n_k} - b_{n_k}) \geq c - 4\varepsilon.$$

Since c and ε are arbitrary, $\text{ca}(x_k) \leq \delta(x_k)$.

Inequality (2) from Proposition 1.1 finishes the proof.

3. CONSTRUCTION OF EXAMPLE 1.4

The construction is based upon Example 4 of [8]. It uses a standard renorming technique (see, e.g., [7, Proposition III.2.11]). We recall that $\beta\mathbb{N}$ is the Čech–Stone compactification of $\mathbb{N}$ and $M(\beta\mathbb{N})$ is the space of all signed Radon measures on $\beta\mathbb{N}$ considered as the dual of ℓ_∞ .

Let us fix $\alpha > 0$ and consider the space

$$Y_\alpha = (\ell_1, \alpha \|\cdot\|_1) \oplus_1 (\ell_2, \|\cdot\|_2).$$

Here $\|\cdot\|_1$ and $\|\cdot\|_2$ denote the usual norms on ℓ_1 and ℓ_2 , respectively. Note that we have the following canonical identifications:

$$\begin{aligned} Y_\alpha^* &= (\ell_\infty, \tfrac{1}{\alpha} \|\cdot\|_\infty) \oplus_\infty (\ell_2, \|\cdot\|_2) \text{ and} \\ Y_\alpha^{**} &= (M(\beta\mathbb{N}), \alpha \|\cdot\|_{M(\beta\mathbb{N})}) \oplus_1 (\ell_2, \|\cdot\|_2). \end{aligned}$$

For $k \in \mathbb{N}$, let $x_k = (e_k, e_k) \in Y_\alpha$, where e_k denotes the k -th canonical basic vector. Let X_α be the closed linear span of the set $\{x_k : k \in \mathbb{N}\}$.

We claim that

$$(6) \quad X_\alpha = \{(x, y) \in Y_\alpha : x(n) = y(n), n \in \mathbb{N}\}.$$

Indeed, the set on the right-hand side is a closed linear subspace of Y_α containing x_k for each $k \in \mathbb{N}$ which yields the inclusion ‘ $\subset$ ’. To prove the converse inclusion, let us take any element $(z, y) \in Y_\alpha$ satisfying $z(n) = y(n)$ for all $n \in \mathbb{N}$. Since $z \in \ell_1$, we get

$$(z, y) = \sum_{k=1}^{\infty} z(k)x_k \in X_\alpha$$

as the series is absolutely convergent.

Let $T : \ell_1 \rightarrow \ell_2$ denote the identity mapping. Since T maps the unit ball of ℓ_1 into the unit ball of ℓ_2 , we get $\|T\| \leq 1$. Therefore, for an arbitrary element $(x, y) \in X_\alpha$, we have

$$\alpha \|x\|_1 \leq \|(x, y)\|_{X_\alpha} = \alpha \|x\|_1 + \|Tx\|_2 \leq (\alpha + 1) \|x\|_1.$$

Thus the projection on the first coordinate is an isomorphism of X_α onto ℓ_1 . In particular, X_α has the Schur property.

We further observe that X_α^{**} is canonically identified with the weak* closure of X_α in Y_α^{**} ; thus

$$(7) \quad X_\alpha^{**} = \{(\mu, y) \in M(\beta\mathbb{N}) \times \ell_2 : \mu(n) = y(n), n \in \mathbb{N}\}.$$

Indeed, the set on the right-hand side is a weak* closed linear subspace of Y_α^{**} containing X_α , which proves the inclusion ‘ $\subset$ ’. To prove the converse inclusion, we fix (μ, y) in the set on the right-hand side. Take a bounded net (u_τ) in ℓ_1 which weak* converges to μ . For each τ there is a unique $y_\tau \in \ell_2$ such that $(u_\tau, y_\tau) \in X_\alpha$. Then (y_τ) is clearly a bounded net in ℓ_2 . Moreover, we will show that (y_τ) weak* (i.e. weakly) converges to y . Since the weak topology on bounded sets in ℓ_2 coincides with the topology of pointwise convergence, it suffices to show that the y_τ pointwise converge to y . Indeed,

$$y_\tau(n) = \mu_\tau(n) \rightarrow \mu(n) = y(n), \quad n \in \mathbb{N}.$$

It follows that X_α is L -embedded in X_α^{**} because the projection $P : X_\alpha^{**} \rightarrow X_\alpha$ defined as

$$P(\mu, y) = (\mu|_{\mathbb{N}}, y), \quad (\mu, y) \in M(\beta\mathbb{N}) \times \ell_2,$$

satisfies

$$\|(I - P)(\mu, y)\|_{X_\alpha^{**}} + \|P(\mu, y)\|_{X_\alpha^{**}} = \|(\mu, y)\|_{X_\alpha^{**}}, \quad (\mu, y) \in X_\alpha^{**}.$$

Further, for the sequence (x_k) , its weak* cluster points in X_α^{**} are equal to

$$\{(\varepsilon_t, 0) : t \in \beta\mathbb{N} \setminus \mathbb{N}\},$$

where ε_t denotes the Dirac measure at a point $t \in \beta\mathbb{N}$.

We claim that, for our sequence (x_k) , we have

$$(8) \quad \text{ca}(x_k) = 2\alpha + \sqrt{2} \quad \text{and} \quad \delta(x_k) = 2\alpha.$$

To see the first inequality, we observe that for each distinct $k, k' \in \mathbb{N}$ we have

$$\|x_k - x_{k'}\|_{X_\alpha} = \alpha\|e_k - e_{k'}\|_1 + \|e_k - e_{k'}\|_2 = 2\alpha + \sqrt{2}.$$

The second inequality follows from the fact that, given $t, t' \in \beta\mathbb{N} \setminus \mathbb{N}$ distinct, then

$$\|(\varepsilon_t, 0) - (\varepsilon_{t'}, 0)\|_{X_\alpha^{**}} = \|(\varepsilon_t - \varepsilon_{t'}, 0)\|_{X_\alpha^{**}} = \alpha\|\varepsilon_t - \varepsilon_{t'}\|_{M(\beta\mathbb{N})} = 2\alpha.$$

This verifies (8).

Now we use the described procedure to construct the desired space X . For $n \in \mathbb{N}$, let $\alpha_n = \frac{1}{n}$ and let $X_{\frac{1}{n}}$ be the space constructed for α_n . Let

$$X = \left(\sum_{n=1}^{\infty} X_{\frac{1}{n}} \right)_{\ell_1}$$

be the ℓ_1 -sum of the spaces $X_{\frac{1}{n}}$. We claim that X is the required space.

First, since each $X_{\frac{1}{n}}$ has the Schur property, X , as their ℓ_1 -sum, possesses this property as well (this follows by a straightforward modification of the proof that ℓ_1 has the Schur property; see [2, Theorem 5.19]).

Similarly, as an ℓ_1 -sum of L -embedded spaces, X is L -embedded as well (see [7, Proposition 1.5]).

Finally, fix $n \in \mathbb{N}$. We consider a sequence $\widehat{x}_k = (0, \dots, 0, x_k^{n\text{-th}}, 0, \dots)$, where the elements $x_k \in X_{\frac{1}{n}}$, $k \in \mathbb{N}$, are defined above. Then, for any $k, k' \in \mathbb{N}$ distinct,

$$\|\widehat{x}_k - \widehat{x}_{k'}\|_X = \|x_k - x_{k'}\|_{X_{\frac{1}{n}}};$$

hence $\text{ca}(\widehat{x}_k) = \text{ca}(x_k) > \sqrt{2}$ by (8).

On the other hand,

$$\delta(\widehat{x}_k) = \delta(x_k) = \frac{2}{n},$$

again by (8). So,

$$\text{ca}(\widehat{x}_k) > \frac{n}{\sqrt{2}} \delta(\widehat{x}_k).$$

Since $n \in \mathbb{N}$ is arbitrary, the conclusion follows.

4. FINAL REMARKS

In [4, p. 57] a Banach space is said to have the *1-strong Schur property*, whenever for any $\delta \in (0, 2]$, any $\varepsilon > 0$ and any normalized δ -separated sequence in X there is a subsequence which is $(\frac{2}{\delta} + \varepsilon)$ -equivalent to the standard ℓ_1 basis.

Although this property has the adjective “strong”, it is in fact weaker than our 1-Schur property, as witnessed by the following proposition. We do not know whether these two properties are in fact equivalent but we conjecture that it is not the case. The reason for this opinion is the fact that the 1-Schur property is a property of all

bounded sequences while the 1-strong Schur property is a property of very special sequences.

Proposition 4.1. • *Any Banach space with the 1-Schur property has the 1-strong Schur property.*
 • *Any Banach space with the 1-strong Schur property has the 5-Schur property.*

Proof. Suppose that X is a Banach space with the 1-Schur property and fix $\delta \in (0, 2]$, $\varepsilon > 0$ and a normalized δ -separated sequence (x_k) in X . Since (x_k) is δ -separated, $\text{ca}(x_{k_n}) \geq \delta$ for each subsequence (x_{k_n}) of (x_k) . By the 1-Schur property we get $\delta(x_{k_n}) \geq \delta$ for each subsequence. By [1, Theorem 3.2] there is a subsequence (x_{k_n}) such that

$$\left\| \sum_{n=1}^N \alpha_n x_{k_n} \right\| \geq \frac{1}{\frac{2}{\delta} + \varepsilon} \sum_{n=1}^N |\alpha_n|$$

for each $N \in \mathbb{N}$ and each choice of real numbers $\alpha_1, \dots, \alpha_N$. Note that we used that

$$\frac{1}{\frac{2}{\delta} + \varepsilon} < \frac{\delta}{2}.$$

It follows that (x_{k_n}) is $(\frac{2}{\delta} + \varepsilon)$ -equivalent to the standard ℓ_1 basis.

Conversely, suppose that X is a Banach space which has the 1-strong Schur property. Let (x_k) be a bounded sequence in X and $c > 0$ be such that $\text{ca}(x_k) > 5c$. We will show that $\delta(x_k) \geq c$. We will distinguish two cases:

- (a) There is $n \in \mathbb{N}$ such that for each $k \in \mathbb{N}$ we have $d(x_k, \{x_1, \dots, x_n\}) \leq c$.
- (b) For each $n \in \mathbb{N}$ there is $k > n$ such that $d(x_k, \{x_1, \dots, x_n\}) > c$.

It is clear that exactly one of these two cases takes place.

First suppose that the case (a) occurs. As $\text{ca}(x_k) > 5c$, there are two subsequences (u_k) and (v_k) of (x_k) such that $\|u_k - v_k\| > 5c$ for each $k \in \mathbb{N}$. For each $k \in \mathbb{N}$ fix $p_k, q_k \in \{1, \dots, n\}$ such that $\|u_k - x_{p_k}\| \leq c$ and $\|v_k - x_{q_k}\| \leq c$. Up to taking a subsequence we can assume that the sequence (p_k) is constant. Denote the common value by p . Up to taking a further subsequence we can suppose that the sequence (q_k) is constant as well. Let q be the common value. Set $a = x_p$ and $b = x_q$. Let u^{**} be a weak* cluster point of (u_k) in X^{**} . Similarly, let v^{**} be a weak* cluster point of (v_k) . Then

$$\begin{aligned} \|v^{**} - u^{**}\| &\geq \|b - a\| - \|v^{**} - b\| - \|u^{**} - a\| \geq \|b - a\| - 2c \\ &\geq \|v_1 - u_1\| - \|b - v_1\| - \|a - u_1\| - 2c > 5c - 4c = c. \end{aligned}$$

Hence $\delta(x_k) > c$.

Next suppose that the case (b) occurs and fix an arbitrary $\varepsilon > 0$. It is easy to construct a c -separated subsequence (y_k) of (x_k) . As (y_k) is bounded, without loss of generality we can suppose that the sequence $(\|y_k\|)$ converges to some $\alpha \geq 0$. As the sequence is c -separated, necessarily $\alpha \geq \frac{c}{2}$. Up to omitting a finite number of (y_k) we can suppose that $|\|y_k\| - \alpha| < \varepsilon\alpha$ for each $k \in \mathbb{N}$. Then the sequence $(\frac{y_k}{\|y_k\|})$ is a normalized sequence which is $(\frac{c}{\alpha} - 2\varepsilon)$ -separated. Indeed, for any distinct

$k, l \in \mathbb{N}$ we have

$$\begin{aligned} \left\| \frac{y_k}{\|y_k\|} - \frac{y_l}{\|y_l\|} \right\| &\geq \left\| \frac{y_k}{\alpha} - \frac{y_l}{\alpha} \right\| - \left\| \frac{y_k}{\|y_k\|} - \frac{y_k}{\alpha} \right\| - \left\| \frac{y_l}{\|y_l\|} - \frac{y_l}{\alpha} \right\| \\ &> \frac{c}{\alpha} - \left| \frac{\alpha - \|y_k\|}{\alpha} \right| - \left| \frac{\alpha - \|y_l\|}{\alpha} \right| > \frac{c}{\alpha} - 2\varepsilon. \end{aligned}$$

By the 1-strong Schur property we can extract a subsequence of $\left(\frac{y_k}{\|y_k\|}\right)$ which is $\left(\frac{2}{\frac{c}{\alpha} - 2\varepsilon} + \varepsilon\right)$ -equivalent to the standard ℓ_1 -basis. Thus

$$\delta\left(\frac{y_k}{\|y_k\|}\right) \geq \frac{2}{\frac{2}{\frac{c}{\alpha} - 2\varepsilon} + \varepsilon}$$

by [8, Lemma 5]. Since $\varepsilon > 0$ is arbitrary, we get $\delta\left(\frac{y_k}{\|y_k\|}\right) \geq \frac{c}{\alpha}$. It follows that $\delta(y_k) \geq c$; thus $\delta(x_k) \geq c$. This completes the proof. $\square$

As we remarked in the beginning of the paper, we worked only with real spaces. However, all the results except possibly for Proposition 4.1 are true also for complex spaces. Let us explain this.

Let X be a complex Banach space. Denote by X_R the real version of X , i.e., the same space considered over the reals. Given a bounded sequence (x_k) in X , all the considered quantities ($d, \widehat{d}, \delta, ca$) are the same for X and for X_R . For the quantities $d, \widehat{d}$ and δ , this is explained in [8, Section 5]; for the quantity ca , it is trivial. Hence, in particular, X is C -weakly sequentially complete (or has the C -Schur property) if and only if X_R has this property. Thus Propositions 1.1 and 1.2 are valid for complex spaces as well.

The proof of Theorem 1.3 can be easily adapted to the complex case. It is enough to define x^* using complex signs (i.e., appropriate complex units) and to estimate from below the real part of $x^*(y_{n_k})$. The proof of Example 1.4 works in the complex case without any change.

As for Proposition 4.1, the second part works for complex spaces without any change. It is not clear whether the first part is valid, because the proof uses a result of [1] which works for real spaces and it is an open problem whether it is valid for complex spaces as well. It is also not clear whether the 1-strong Schur property of X is equivalent to that of X_R .

ACKNOWLEDGEMENT

We are grateful to the referee for helpful comments which led to an improvement of our paper.

REFERENCES

- [1] E. Behrends. New proofs of Rosenthal's l^1 -theorem and the Josefson-Nissenzweig theorem. *Bull. Polish Acad. Sci. Math.*, 43(4):283–295 (1996), 1995. MR1414785 (97k:46011)
- [2] M. Fabian, P. Habala, P. Hájek, V. Montesinos Santalucía, J. Pelant, and V. Zizler. *Functional analysis and infinite-dimensional geometry*. CMS Books in Mathematics/Ouvrages de Mathématiques de la SMC, 8. Springer-Verlag, New York, 2001. MR1831176 (2002f:46001)
- [3] G. Godefroy. Renormings of Banach spaces. In *Handbook of the geometry of Banach spaces*, Vol. I, pages 781–835. North-Holland, Amsterdam, 2001. MR1863706 (2003f:46023)
- [4] G. Godefroy, N. J. Kalton and D. Li. On subspaces of L^1 which embed into ℓ_1 . *J. Reine Angew. Math.*, 471:43–75, 1996. MR1374918 (97d:46017)

- [5] G. Godefroy, N. J. Kalton and D. Li. Operators between subspaces and quotients of L^1 . *Indiana Univ. Math. J.*, 49(1):245–286, 2000. MR1777031 (2001e:46025)
- [6] A.S. Granero, J.M. Hernández and H. Pfitzner. The distance $\text{dist}(\mathcal{B}, X)$ when $\mathcal{B}$ is a boundary of $B(X^{**})$. *Proc. Amer. Math. Soc.* 139(3):1095–1098, 2011. MR2745660
- [7] P. Harmand, D. Werner, and W. Werner. *M-ideals in Banach spaces and Banach algebras*, volume 1547 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1993. MR1238713 (94k:46022)
- [8] O. Kalenda, H. Pfitzner and J. Spurný. On quantification of weak sequential completeness. *J. Funct. Anal.*, 260(10):2986–2996, 2011. MR2774062

DEPARTMENT OF MATHEMATICAL ANALYSIS, FACULTY OF MATHEMATICS AND PHYSICS, CHARLES UNIVERSITY, SOKOLOVSKÁ 83, 186 75, PRAHA 8, CZECH REPUBLIC
E-mail address: kalenda@karlin.mff.cuni.cz

DEPARTMENT OF MATHEMATICAL ANALYSIS, FACULTY OF MATHEMATICS AND PHYSICS, CHARLES UNIVERSITY, SOKOLOVSKÁ 83, 186 75, PRAHA 8, CZECH REPUBLIC
E-mail address: spurny@karlin.mff.cuni.cz